SAT0271

RELATIONSHIP BETWEEN ANTI-MDA5 ANTIBODIES AND CANCER: RETROSPECTIVE ANALYSIS OF AN INTERNATIONAL AND MULTIDISCIPLINARY COHORT

Ludovico De Stefano¹, Santos Castañeda², Ellen De Langhe³, Jörg Distler⁴, Johannes Knitza⁴, Ilaria Cavazzana^{5,6}, Alain Meyer^{5,7}, Jorge Rojas-Serrano⁸, Eugenio Arrigoni⁹, Federica Furini¹⁰, Marcello Govoni¹⁰, Paola Parrocchini¹¹, Giovanni Zanframundo¹, José Antonio P. Da Silva¹², Elisa Pedrollo¹³, Florenzo Iannone¹⁴, Carlomaurizio Montecucco¹, Lorenzo Cavagna¹, The AENEAS collaborative group. ¹Rheumatology, Policlinico S. Matteo Foundation, European Reference Network ReCONNET, pavia, Italy; ²Rheumatology, Hospital Universitario de la Princesa, madrid, Spain; ³Rheumatology, University Hospital, leuven, Belgium; ⁴Internal Medicine, University Hospital, erlangen, Germany; ⁵European Reference Network ReCONNET, pavia, Italy; ⁶Rheumatology, AOU Spedali Civili, brescia, Italy; ⁷Hopitaux Universitaires de Strasbourg, strasbourg, France; ⁸ILD and Rheumatology Unit, Instituto Nacional de Enfermedades Respiratorias, mexico city, Mexico; ⁹Rheumatology Unit, Ospedale Guglielmo da Saliceto, piacenza, Italy; ¹⁰Rheumatology, AOU S. Anna, ferrara, Italy; ¹¹ISOD Immunologia e Ter. Cell., AOU Careggi, firenze, Italy; ¹²12. Rheumatology, Centro Hospitalar e Universitario de Coimbra, coimbra, Portugal; ¹³Rheumatology, Policlinico S. Matteo Foundation, pavia, Italy; ¹⁴Rheumatology Unit, Department of Emergency and Organ Transplantations, Policlinico, bari, Italy

Background: Myositis specific antibodies play a central role in the assessment of idiopathic inflammatory myopathies (IIMs) as their presence may indicate specific disease subsets/manifestations or cancer risk. At present, anti-MDA5 antibodies have been associated with the occurrence of rapidly progressive interstitial lung disease (RP-ILD), a life-threatening condition, generally refractory to treatment.

Objectives: To describe the relationship between malignancy and anti-MDA5 antibodies in IIMs.

Methods: Patients with anti-MDA5+ IIMs and a history of cancer were included in this study. Data were retrospectively collected. Because of the lack of RP-ILD shared definitions, we defined RP-ILD as an admission to the Intensive Care Unit (ICU) for IIMs related respiratory insufficiency.

Results: In our cohort (n=133, 67% females) we identified 17 patients (13%) with a history of cancer (8 males [47%], 9 females [53%]; median age at IIMs onset 62 years [IQR 53-67], at cancer diagnosis 56 years [IQR 51-66]). All but 1 patient had solid tumours, 5 (29%) had breast cancer. In 9 patients (53%) cancer diagnosis ranged $\pm$ 3 years from IIMs onset; in 3 patients the delay exceeded 120 months. Three patients (18%) developed RP-ILD respectively 82, 66 and 0 months after the diagnosis of cancer. Four (24%) patients died respectively 0 (neoplasia), 0 (RP-ILD in ICU, diagnosis of cancer after an autopsy), 7 (neoplasia) and 66 (infection after RP-ILD in ICU) months after cancer diagnosis. The 13 surviving patients had a median rheumatological follow-up of 47 months (IQR 19-84) and an oncological follow-up of 57 months (IQR 10-130). Interestingly, in the overall cohort 66% of patients without cancer (76 out of 116) had less than 3 years of follow-up. Of the 22 RP-ILD patients without neoplasia 14 died, 11 during ICU admission. In the latter patients, we could not rule out an occult neoplasia due to the emergency of the situation. Only one patient, with a 18 month history of IIMs before ICU admission, underwent autopsy and was reported negative for cancer.

Figure 1: temporal relationship between anti-MDA5 related IIMs and cancer diagnosis, comprehensive of neoplasia site and outcome; legend: IIMs, Idiopathic Inflammatory Myopathies; Mo= months

Conclusion: This is the first study addressed to tackle the relationship between cancer and anti-MDA5+ IIMs. Although in some cases no clear relationship between IIMs and cancer can be proved because of the time gap, in others the timing of occurrence of cancer and IIMs was very close, suggesting a link between these manifestations. Finally, we emphasize the importance of an accurate neoplasm screening in anti-MDA5+ RP-ILD, because treatment refractoriness could suggest the paraneoplastic nature of the manifestation.

REFERENCES

- [1] Barsotti S, et al. Rheumatol Int 2014
- [2] Lu X, et al. PLoS One 2014

Disclosure of Interests: Ludovico De Stefano: None declared, Santos Castañeda Consultant for: Amgen, BMS, Pfizer, Lilly, MSD, Roche, Sanofi, UCB, Ellen De Langhe: None declared, Jörg Distler: None declared, Johannes Knitza: None declared, Ilaria Cavazzana: None declared, Alain Meyer: None declared, Jorge Rojas-Serrano: None declared, Eugenio Arrigoni: None declared, Federica Furini: None declared, Marcello Govoni Paid instructor for: Pfizer, Roche, Speakers bureau: Pfizer, Abbvie, MSD, Roche, Eli-Lilly, Celgene, Sanofi, Janssen, Paola Parrocchini: None declared, Giovanni Zanframundo: None declared, José Antonio P. da Silva: None declared, Elisa Pedrollo: None declared, Florenzo Iannone Consultant for: F Iannone has received consultancy fees and/or speaker honoraria from Pfizer, AbbVie, MSD, BMS, Novartis, Lilly, UCB outside this work, Speakers bureau: F Iannone has received consultancy fees and/or speaker honoraria from Pfizer, AbbVie, MSD, BMS, Novartis, Lilly, UCB outside this work, Carlomaurizio Montecucco Speakers bureau: AbbVie, Bristol-Myers Squibb, Celgene, Sanofi, Genzyme, Lilly, MSD, Pfizer, UCB, Lorenzo Cavagna: None declared

DOI: 10.1136/annrheumdis-2019-eular.4565

SAT0272

RITUXIMAB SIGNIFICANTLY DIMINISHES CD86+ B CELLS IN SYSTEMIC SCLEROSIS

Barbara Dreo¹, Barbara Prieti^{2,3}, Selina Kofler^{2,3}, Harald Sourij^{2,3}, Angelika Lackner¹, Florentine Fürst-Moazedi¹, Monica D'orazio¹, Martin Stradner¹, Winfried Graninger¹, Hans-Peter Brezinsek¹. ¹Medical University of Graz, Division of Rheumatology and Immunology, Graz, Austria; ²CBmed GmbH – Center for Biomarker Research in Medicine, Graz, Austria; ³Medical University of Graz, Division of Endocrinology and Diabetology, Graz, Austria

Background: Rituximab (RTX) is a monoclonal antibody that targets the CD20 surface marker. This results in a reduction of CD20+ immune cell populations, foremost B cells. B cell depletion via RTX was found to be beneficial for patients suffering from systemic sclerosis (SSc), leading to an improvement of skin fibrosis and autoimmunity (1,2). However, little is known about the influence of RTX on specific B subsets in SSc.

Objectives: The purpose of this study was to further characterize the effect of RTX on B cell populations in patients with SSc.

Methods: Peripheral blood samples from 37 patients suffering from SSc (mean age: 54 years $\pm$ 1.64 SEM, female ratio: 0.78) and 10 age-matched healthy participants were drawn over a sampling period of 2 years. 20 SSc patients in this study group were at the time under RTX treatment and the modified Rodnan Skin score (mRSS) was measured before and after treatment start. The percentage of CD19⁺, CD20⁺ lymphatic cells and CD19⁺CD20⁺ B cells co-expressing either CD5, CD24, CD27 or CD86 on their surface was done by staining freshly isolated PBMCs. A quantitative flow cytometric bead-based assay (QuantiBRITE PE kit from Becton Dickinson) was used for the estimation of CD19 antibodies bound per cell. All cytometric measurements were performed using a standardized BD LSRFortessa platform.

Results: RTX induced a significant decrease in mRSS from 19.7 ± 2.8 to 8.1 ± 1.7 (mean $\pm$ SEM; $p < 0.000$). In addition, CD19⁺CD20⁺ cells were significantly diminished as a result of the treatment. Thus, the frequency of CD19⁺CD20⁺ cells in the non-treatment group was $1.3\% \pm 0.3\%$ compared to $0.7\% \pm 0.2\%$ ($p = 0.048$). Within the B cell population $33.3\% \pm 3.8\%$ were positive for CD86, a checkpoint molecule for the activation of T cells during an immune response. RTX treatment significantly decreased this B cell population to $11.2\% \pm 4.7\%$ ($p = 0.039$). Quantification of the number of CD19 molecules on the surface of CD19⁺CD20⁺ B cells revealed a significantly lower number in SSc patients compared to healthy participants. The mean $\pm$ SE of molecules per cell was 6862 ± 625 and 7449 ± 569 , respectively ($p \leq 0.001$).

Conclusion: RTX treatment in SSc might not only be effective by reducing B cells but also by down regulation of the CD86 B cell surface marker on B cells. This would indicate that B cells under RTX treatment are less capable of activating T cells.

REFERENCES

- [1] Lafyatis R, Kissin E, York M, Farina G, Viger K, Fritzler MJ, et al. B Cell Depletion With Rituximab in Patients With Diffuse Cutaneous Systemic Sclerosis. 2009;60(2):578–83.

- [2] Jordan S, Distler JHW, Maurer B, Huscher D, Laar JM Van, Allanore Y, et al. Effects and safety of rituximab in systemic sclerosis: an analysis from the European Scleroderma Trial and Research (EUSTAR) group. 2014;1–7.

Acknowledgement: Work done in "CBmed" was funded by the Austrian Federal Government within the COMET K1 Centre Program, Land Steiermark and Land Wien.

Disclosure of Interests: Barbara Dreo: None declared, Barbara Prietl: None declared, Selina Kofler: None declared, Harald Sourij: None declared, Angelika Lackner: None declared, Florentine Fürst-Moazed: None declared, Monica D'Orazio: None declared, Martin Stradner Speakers bureau: Novartis, Roche, Lilly, BMS, Pfizer, Shire, Winfried Graninger: None declared, Hans-Peter Brezinssek: None declared

DOI: 10.1136/annrheumdis-2019-eular.8047

SAT0273

PREDICTIVE VALUE OF THE REVISED EUROPEAN SCLERODERMA TRIALS AND RESEARCH GROUP ACTIVITY INDEX (EUSTAR-AI)

Serena Fasano¹, Veronica Giacco¹, Antonella Riccardi¹, Valentina Messiniti¹, Alessandra Vacca², Oliver Distler³, Otylia Kowalka Bielecka⁴, Yannick Allanore⁵, Gabriele Valentini¹, European Scleroderma Trials and Research group (EUSTAR) co-authors. ¹University of Campania "Luigi Vanvitelli", Rheumatology, Naples, Italy; ²University Hospital of Cagliari, Rheumatology, Cagliari, Italy; ³University Hospital Zurich, Rheumatology, Zurich, Switzerland; ⁴Medical University of Bialystok, Rheumatology and Internal Medicine, Bialystok, Poland; ⁵Cochin Hospital, Paris Descartes University, Rheumatology, Paris, France

Background: Disease activity as measured by the European Scleroderma Study Group Activity Index (EScSG-AI) (1) has been recently found to predict the development of damage over time in an early systemic sclerosis (SSc) cohort (2). The European Scleroderma Trials and Research Group Task Force for the Development of Revised Activity Criteria for SSc recently succeeded in identifying a preliminarily revised activity index (EUSTAR-AI) that had a greater construct validity than the EScSG-AI i.e. performing better in identifying patients with active disease (3).

Objectives: To assess in patients with SSc the predictive value of the EUSTAR-AI for disease severity accrual.

Methods: SSc patients from the EUSTAR database with a disease duration from the onset of the first non-Raynaud sign/symptom ≤ 5 years were first extracted. Patients were considered for the study if they presented the following features: a) availability of items included in the EUSTAR-AI, in the EScSG-AI and in the Medsger severity scale at baseline and yearly for 2 consecutive years; b) availability of vital status at the last observation; c) availability of other disease features known to predict disease progression (male sex, clinical and serological subset).

To capture the disease activity variations over time, we calculated the EUSTAR-AI and EScSG-AI adjusted mean (area under the curve over time divided by time interval). Disease progression was based on the Medsger severity scale and included accrual of a $\Delta \geq 1$ summed severity score and of a $\Delta \geq 1$ severity in each organ systems at the 2 year follow-up visit compared with the initial visit. To explore specific determinants of disease progression, logistic regression analysis was carried out.

Results: A total of 549 patients satisfied the entry criteria. At univariate logistic regression analysis (among sex, age, clinical and serological subset, EScSG-AI and EUSTAR-AI adjusted mean and baseline severity score), EScSG-AI adjusted mean (OR 1.41 95%CI 1.20-1.67), antiScl70 antibody positivity (OR 1.72 95%CI 1.20-2.47), diffuse subset (OR:1.46 95%CI 1.01-2.10) and EUSTAR-AI adjusted mean (OR 1.41 95%CI 1.23-1.61) predicted disease severity accrual. Multivariate analysis revealed that the EUSTAR-AI adjusted mean was the best predictor of disease progression (Table1). Moreover, at multivariate analysis the EUSTAR-AI adjusted mean also predicted severity accrual of lung (OR 1.32), heart (OR 1.40), skin (OR 1.48) and peripheral vascular disease (OR 1.45).

Conclusion: The adjusted mean EUSTAR-AI has a distinct predictive value for disease progression and development of severe organ involvement in SSc and works better than EScSG-AI.

REFERENCES

- [1] Valentini G, et al. Ann Rheum Dis 2001;60:592-8.
[2] Nevskaya T, et al. Rheumatology (Oxford). 2017;56:1111-1122.
[3] Valentini G, et al. Ann Rheum Dis. 2017;76:270-276.

Table 1.. Predictive features of severity progression by multivariate regression analysis

Predictive factors	OR (IC 95%)	severity deterioration	p- value
Adjusted mean EUSTAR-AI	1.41 (1.23-1.61)	Δ summed severity score ≥ 1	<0,0001
Adjusted mean EUSTAR-AI	1.32 (1.14-1.54)	Δ lung severity score ≥ 1	0,0002
Adjusted mean EUSTAR - AI age	1.40 (1.15-1.70)	Δ heart severity score ≥ 1	0,0005 0,0223
	1.03 (1.01-1.05)		
Adjusted mean EUSTAR-AI age	1.48 (1.21-1.82)	Δ Skin severity score ≥ 1	0,0002 0,0359
	0.97 (0.95-0.99)		
Adjusted mean EUSTAR-AI	1.45 (1.24-1.70)	Δ Peripheral Vascular severity score ≥ 1	<0,0001

Disclosure of Interests: Serena Fasano: None declared, Veronica Giacco: None declared, Antonella Riccardi: None declared, Valentina Messiniti: None declared, ALESSANDRA VACCA: None declared, Oliver Distler Grant/research support from: Prof. Distler received research funding from Actelion, Bayer, Boehringer Ingelheim and Mitsubishi Tanabe to investigate potential treatments of scleroderma and its complications, Consultant for: Prof. Distler has/had consultancy relationship within the last 3 years with Actelion, AnaMar, Bayer, Boehringer Ingelheim, ChemomAb, espeR-are foundation, Genentech/Roche, GSK, Inventiva, Italfarmaco, IQVIA, Lilly, medac, MedImmune, Mitsubishi Tanabe Pharma, Pharmacyclics, Novartis, Pfizer, Sanofi, Serodapharm and UCB in the area of potential treatments of scleroderma and its complications. In addition, he had/had consultancy relationship within the last 3 years with A. Menarini, Amgen, Abbvie, GSK, Mepha, MSD, Pfizer and UCB in the field of arthritides and related disorders, Otylia Kowalka Bielecka Consultant for: "OK-B received consulting fees or other remuneration from Bayer, Boehringer Ingelheim, Inventiva, Medac, Novartis and Roche", Yannick Allanore Grant/research support from: Inventiva, F Hoffman La-Roche, Sanofi, BMS, Pfizer, Consultant for: Actelion, Bayer, BMS, Boehringer, Roche, Sanofi, Gabriele Valentini Grant/research support from: MSD, Pfizer, Consultant for: MSD, Pfizer, biogen, Speakers bureau: MSD, amgen, biogen, lilly, sanofi, phizer

DOI: 10.1136/annrheumdis-2019-eular.3485

SAT0274

THE EFFECT OF RITUXIMAB ON LUNG FUNCTION AND SKIN SCORE IN SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE. LONG-TERM OBSERVATION

Liudmila Garzanova¹, Lidia P. Ananyeva¹, Olga Koneva¹, Oxana Desinova¹, Olga Ovsyannikova¹, Mayya Starovoytova¹, Alexander Volkov², Svetlana Glukhova². ¹V.A. Nasonova Research Institute of Rheumatology, Laboratory of microcirculation and inflammation, Moscow, Russian Federation; ²V. A. Nasonova Research Institute of Rheumatology, Diagnostic department, Moscow, Russian Federation

Background: There is a large clinical experience about the efficiency of rituximab (RTX) for the treatment of systemic sclerosis (SSc). There are several studies showing the decrease in skin induration and interstitial fibrosis in the lungs as the effect of therapy. However, there are not many long-term observations.

Objectives: To describe the efficacy of RTX on lung function and skin score in patients with systemic sclerosis-associated interstitial lung disease, in long-term follow-up.

Methods: This study included 71 patients (pts) with SSc. Data were collected prospectively. The mean follow-up period was 42 mo (12-72). Mean age was 46 years (17-66), female-59 (83%), diffuse cutaneous subset of the disease had 42 (59%), Scl-70 positivity-73% of pts. Duration of the disease was 5,6±4,4 yrs. All pts received concomitant treatment with low dose prednisolone and 45% - with immunosuppressants. The following indicators were evaluated: forced vital capacity,% predicted (FVC), diffusing capacity for carbon monoxide,% predicted (DLCO) and Rodnan skin score (mRss) over a periods of 12-18 months (point 1), 24-30 months (point 2), 36-42 months (point 3), 48-54 months (point 4) and 60-72 months (point 5) after the start of therapy. The results are presented in the form of mean values, delta, median, upper and lower quartile.