Results: In point 1 (n=71) the cumulative mean dose of RTX was 1,43 ±0,6 gr. The mRss decreased from 11,3±9,6 to 8±6,6 (p = 0,000007). FVC increased from 77,4±19,9 to 82,6±20,7% (p=0.0001). DLCO remained stable (from 47,0±18,5 to 47,23±16,7%). In point 2 (n=55) the cumulative mean dose of RTX was 2,97±0,8 gr. ΔmRss = 5,4 (median 3; 25th% 0; 75th% 10). ΔFVC = 7,5% (median 8,2; 25th% -1,1; 75th% 14,4). ΔDLCO = 0,21% (median -0,2; 25th% -6,2; 75th% 6,6). In point 3 (n=36) the cumulative mean dose of RTX was 3,45±1,3 gr. ΔmRss = 5,1 (median 3,5; 25th% 0; 75th% 9). ΔFVC = 9,3% (median 7,7; 25th% 1,45; 75th% 15,3). ΔDLCO = 3,4% (median 3,6; 25th% 2,45; 75th% 7,76), p=0,02. In point 4 (n=24) the cumulative mean dose of RTX was 3,96±1,1 gr. ΔmRss = 5,3 (median 3; 25th% 0; 75th% 10). ΔFVC = 12,2% (median 7,9; 25th% 1,1; 75th% 24,2). ΔDLCO = 3,9% (median 0,45; 25th% -0,95; 75th% 6,8). In point 5 (n=17) the cumulative mean dose of RTX was 5,15±1,7 gr. It should be noted that this group included patients with initially the lowest DLCO (below 40%). ΔmRss = 7,3 (median 5; 25th% 1; 75th% 14). ΔFVC = 13,2% (median 11; 25th% 8,7; 75th% 23,4). ΔDLCO = 5,9% (median 1,6; 25th% -4,8; 75th% 13,8).

Conclusion: The results of this study confirm the data on the positive effect of RTX in patients with SSC (decrease of skin induration, increase of FVC, stabilization of DLCO). The decrease of skin score is accompanied by the improvement of lung function indicators. In our study, there was a significant increase of DLCO associated with long-term treatment (over 36 mo) and a cumulative dose of RTX over 3,45±1,3 gr. Patients with initially lower DLCO can achieve a significant improvement by the 60th month of RTX therapy. Our work shows that patients with SSC-associated interstitial lung disease are required long-term treatment with RTX (at least 3-5 years) to achieve an obvious improvement of the lung function.

REFERENCE

- [1] Jordan S, Distler JH, Maurer B, Huscher D, van Laar JM, Allanore Y. et al; EUSTAR Rituximab study group. Effects and safety of rituximab in systemic sclerosis: an analysis from the European Scleroderma Trial and Research (EUSTAR) group. *Ann Rheum Dis* 2015;74(6):1188–94.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5565

SAT0275

PERFORMANCE OF THE ANTISYNTHEASE ANTIBODIES AND THEIR INDIRECT IMMUNOFLUORESCENCE PATTERNS IN THE ANTISYNTHEASE SYNDROME DIAGNOSIS

Martín Greco¹, María Jesús García de Yébenes², Inmaculada Alarcón¹,
^{3,4}Anahy Brandy-García⁴, Iñigo Rua-Figueroa¹, Estibaliz Loza², Teresa Oton²,
 Francisco Rubiño¹, Juan Manuel Díaz¹, Félix Francisco¹, Soledad Ojeda¹,
 Loreto Carmona². ¹Hospital Universitario de Gran Canaria Dr. Negrín., Las
 Palmas, Spain; ²Instituto de Salud Musculoesquelética Inmusc., Madrid, Spain;
³Hospital Universitari Germans Trias i Pujol, Barcelona, Spain; ⁴Hospital
 Universitario Central de Asturias, Oviedo, Spain

Background: The antisynthetase syndromes (ASSD) are characterized by the presence of anti-aminoacyl transfer RNA synthetase (ARS) autoantibodies; which difficult the binding of amino acids to the transfer RNA during the protein synthesis. ARS can be detected by indirect immunofluorescence (IIF), and can be identified by immunoblot assay and ELISA (Enzyme-Linked Immunosorbent Assay) and immunoblotting. The main clinical features of the ASSD are myositis, arthritis, interstitial lung disease, Raynaud's phenomenon, mechanic hands, and fever. Two ASSD diagnosis criteria have been developed; those proposed by Connors, and the stricter criteria proposed by Solomon (1, 2).

Objectives: To evaluate the performance of the ARS and their IIF patterns in the ASSD diagnosis.

Methods: We performed an observational retrospective study in one center during the period 06/2008-06/2018. We searched all the myositis immunoblots (Euroimmun assay) requested by the Rheumatologists under suspicion of ASSD or myositis. We assessed: 1) the rate of cases with positive ARS; 2) the rate of cases with Connor's or Solomon's diagnosis criteria fulfillment; and 3) their relation with the IIF patterns (Hep-2 cells; ≥1/80) evaluated by an expert in autoimmune tests.

Results: A total of 140 myositis immunoblots were searched. Twenty-seven cases (19.3%) presented positive ARS: anti-Jo1 (n=13), anti-PL-12 (n=7), anti-PL-7 (n=1), anti-EJ (n=2), and anti-OJ (n=4). Twenty-five of these (17.9%) fulfilled Connors' criteria, and 15 (10.7%) additionally met Solomon's criteria. Thus, the fulfillment of Connor's and Solomon's

criteria in patients with a positive ARS was of 92.6% and 55.5%, respectively.

All cases (100%) with positive ARS presented positive immunofluorescence: 19 (70.4%) showed a cytoplasmic pattern (10 of them with an associated nuclear pattern) and 8 cases (29.6%) presented only a nuclear pattern. On the other hand, 99 of the 113 cases (87.6%) with negative ARS presented positive IIF: 29 (25.7%) showed a cytoplasmic pattern (21 of them with an associated nuclear pattern) and 42 cases (37.2%) presented only a nuclear pattern.

Correlating the ARS positivity, IIF pattern and the diagnosis criteria fulfillment:

- 13 of 15 cases (86.6%) with positive ARS and Solomon's criteria fulfillment presented a cytoplasmic pattern; and 2 of 15 cases (13.3%) presented only a nuclear pattern.

- 13 of 19 cases (68.4%) with positive ARS and cytoplasmic pattern fulfilled Solomon's criteria; and 6 only fulfilled those from Connors'.

Conclusion: One-fifth of the immunoblots requested by Rheumatologists presented positive ARS; almost all these cases fulfilled Connor's criteria, and more than a half fulfilled the stricter Solomon's criteria. All patients with positive ARS, and a high rate of those without ARS, presented positive IIF. The presence of a cytoplasmic pattern was considerably higher in patients with ARS positivity and in those that met Solomon's criteria.

Thus, our results suggests that in patients evaluated by a Rheumatologist, with clinical suspicion of ASSD or myositis and with ARS positivity, the probability of fulfilling Solomon's criteria is higher when the IIF presents a cytoplasmic pattern than when only a nuclear pattern is observed. Nevertheless, presenting only a nuclear pattern does not exclude the detection of ARS in the myositis immunoblot and the fulfillment of Solomon's criteria.

REFERENCES

- [1] Solomon J, et al. *Jornal brasileiro de pneumologia*. 2011;37(1):100-9.
- [2] Connors GR, et al. *Chest*. 2010;138(6):1464-74.

Disclosure of Interests: Martín Greco: None declared, María Jesús García de Yébenes: None declared, Inmaculada Alarcón: None declared, Anahy Brandy-García: None declared, Iñigo Rua-Figueroa: None declared, Estibaliz Loza Grant/research support from: Roche, MSD, Pfizer, Abbvie, BMS, UCB, Actelion, Celgene, Grunenthal and Sanofi, Teresa Oton: None declared, Francisco Rubiño: None declared, Juan Manuel Díaz: None declared, Félix Francisco: None declared, Soledad Ojeda: None declared, Loreto Carmona Grant/research support from: Abbvie, Actelion, Astellas, BMS, Eisay, Gebro Pharma, Grünenthal, Leo Pharma, Lilly, MSD, Novartis, Pfizer, Roche, Sanofi-Aventis and UCB Pharma, Paid instructor for: Novartis

DOI: 10.1136/annrheumdis-2019-eular.7182

SAT0276

STUDY OF THE EPIDEMIOLOGICAL, CLINICAL AND ANALYTICAL CHARACTERISTICS IN PATIENTS WITH SYSTEMIC SCLEROSIS AND CANCER IN VALL D'HEBRON HOSPITAL

Maria Roca Herrera¹, Alfredo Guillén del Castillo¹, Eduardo Callejas Moraga²,
 David Bernal Bello³, Jorge Hernando Cubero⁴, Natalia Iriarte Gay de Montella¹,
 Vicent Fonollosa Pla¹, Carmen Pilar Simeón-Aznar¹. ¹Vall d'Hebron Hospital,
 INTERNAL MEDICINE, Barcelona, Spain; ²Parc Taulí Hospital, INTERNAL
 MEDICINE, Sabadell, Spain; ³Fuenlabrada Hospital, INTERNAL MEDICINE,
 Fuenlabrada, Spain; ⁴Vall d'Hebron Hospital, ONCOLOGY, Barcelona, Spain

Background: Scleroderma or systemic sclerosis (SSc) is a systemic, chronic autoimmune disease characterized by great clinical heterogeneity. In recent years, studies have proven that there is a relationship between SSc and neoplasia.

SSc is associated with an increased risk of certain types of cancer, particularly lung, liver, hematological, non-melanoma skin and urothelial cancer. Despite this increase, the relative risk of developing cancer is still low in these patients. In the literature, neoplasms have been described in 3-11% of patients with SSc.

Objectives: Our objective is to analyze the epidemiological, clinical and analytical characteristics previously described as possibly linked to the development of a cancer in patients with systemic sclerosis (SSc) in the Vall d'Hebron Hospital cohort.

Methods: We analyzed 583 patients in the Vall d'Hebron Hospital cohort of SSc. The inclusion criteria were age > 18 years and the diagnosis of SSc limited, diffuse and SSc sine scleroderma. The different variables were analyzed by univariate statistical analysis with SPSS v21.

VARIABLE	NEOPLASIA (n=94)	NO NEOPLASIA (n=427)	P
General Characteristics			
SEX (M/F)	8/86 (9/91%)	51/376 (12/88%)	0.73
AGE >65 AT DIAGNOSIS	24 (26%)	102 (24%)	0.34
MORTALITY	34 (36%)	119 (28%)	0.10
SMOKERS	6 (6%)	36 (8%)	0.50
SUBTYPE (LIMITED /DIFFUSE /SINE)	65/16/13 (70/16/14%)	282/94/51 (66/22/12%)	0.54
CAPILLAROSCOPY PATTERN (SLOW /ACTIVE)	57/6 (60/6%)	236/68 (55/16%)	0.04/0.02
Clinical manifestation			
RAYNAUD'S PHENOMENON	88 (93%)	415 (97%)	0.08
SKIN ULCERS	38 (40%)	214 (50%)	0.08
TELANGIECTASIAS	70 (74%)	325 (76%)	0.73
CALCINOSIS CUTIS	22 (23%)	91 (21%)	0.65
ESOPHAGEAL INVOLVEMENT (PATHOLOGICAL MANOMETRY)	57 (61%)	292 (68%)	0.36
ARTHRALGIA	44 (47%)	236 (55%)	0.13
ARTHRITIS	17 (18%)	81 (19%)	0.84
PULMONARY HYPERTENSION	43 (46%)	184 (43%)	0.63
ILD	25 (27%)	103 (24%)	0.72
CARDIAC AFFECTATION	58 (62%)	285 (67%)	0.38
Immunology pattern			
AC SCL70 (ATA I)	11 (12%)	88 (21%)	0.02
ACA +	37 (39%)	139 (33%)	0.18
ANTI-RNA POL III +	7 (7%)	25 (6%)	0.65
ANTI PM/SCL +	9 (10%)	18 (4%)	0.04
Treatment			
CYCLOPHOSPHAMIDE	9 (10%)	36 (8%)	0.72
AZATHIOPRINE	6 (6%)	88 (21%)	0.00006
MYCOPHENOLATE	10 (11%)	61 (14%)	0.35
METHOTREXATE	0 (0%)	23 (5%)	0.68
ANTIPLATELET	40 (43%)	231 (54%)	0.09
ACEIs	32 (34%)	125 (29%)	0.36
ARBs	20 (21%)	54 (12%)	0.03
CCBs	57 (60%)	304 (71%)	0.04
CORTICOSTEROIDS	40 (42%)	172 (40%)	0.95
PROSTAGLANDINS	4 (4%)	37 (9%)	0.15

TYPES OF CANCER	(n=94)	%
BREAST CANCER	22	24%
HEMATOLOGIC MALIGNANCIES	21	23%
LUNG CANCER	13	14%
NON MELANOMA SKIN CANCER	6	6%
MELANOMA	2	2%
COLON CANCER	5	5%
ENDOMETRIAL CANCER	5	5%
GASTROESOPHAGEAL CANCER	4	4%
THYROID CANCER	4	4%
UROTHELIAL CANCER	2	2%
OVARIAN CANCER	1	1%
OTHERS	9	10%

Results: We included 521 patients with diagnosis from January 1985 to August 2016. 88.7% were women, with an average age of 53. 94 patients were diagnosed with neoplasia. The analyzed variables and their statistical significance and the types of cancer developed by our cohort patients are shown in Fig 1.

Conclusion: In the analyzed cohort, a slow pattern in capillaroscopy, positivity for Pm/Scl and the use of ARBs is associated with the development of neoplasia in patients with SSC. It has an inverse association the positivity of Scl70 (ATA I), the active pattern in capillaroscopy and the use of azathioprine and CCBs. In our cohort 94 patients with SSC have developed cancer. The most prevalent types of cancer were breast cancer, hematologic malignancies and lung cancer, as shown in the literature.

REFERENCES

- Onishi A, Sugiyama D, Kumagai S, Morinobu A. Cancer incidence in systemic sclerosis: meta-analysis of population-based cohort studies. *Arthritis Rheum* 2013; 65: 1913–21.
- Bernal-Bello D, de Tena JG, Guillén-del Castillo A, et al. Novel risk factors related to cancer in scleroderma. *Autoimmun Rev* 2017.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3772

SAT0277 THE 100 MOST INFLUENTIAL PUBLICATIONS IN SYSTEMIC SCLEROSIS: A BIBLIOMETRIC STUDY

Priyanka Iyer¹, Petar Lener¹, Jawad Bila². ¹University of Iowa Hospitals and Clinics, Rheumatology, Iowa City, United States of America; ²University of Arizona, Rheumatology, Tucson, United States of America

Background: In recent years, a large volume of important clinical and scientific papers about systemic sclerosis can be found in the most

influential journals. Bibliometric analysis is a statistical method that enables researchers to analyze citation patterns and highlight characteristics of highly cited scholarly work. It has been infrequently utilized in Rheumatology^{1,2}. Prior studies have not addressed the bibliometric literature gap in the field of systemic sclerosis.

Objectives: This study aims to identify the 100 most cited papers in systemic sclerosis and to describe their characteristics.

Methods: We sought to assess the characteristics of the top 100 most referenced citations in systemic sclerosis. We searched the Scopus and Web of Science databases to identify all articles relating to 'Scleroderma', 'limited' or 'diffuse Systemic Sclerosis' and 'CREST syndrome'. Articles were arranged in descending order of citations and were scrutinized by two independent reviewers. Selected articles were analyzed with respect to number of citations, publication year, journal characteristics, research type, focus area, authors and countries of authors of publications.

Results: Retrieved articles were published mostly in rheumatology journals (n=42) with impact factors ranging between 3.47 and 12.35 in these journals, followed by journals in general medicine, pulmonology, and experimental biomedical research. Journals with the highest number of top-cited articles included *Arthritis and Rheumatism* (n=25), followed by the *The New England Journal of Medicine* (n=9) and *Annals of the Rheumatic Diseases* (n=8). Studies conducted in USA, UK, and Italy that were published in high-impact journals had the highest citation count. Temporal analysis indicated that the majority of top-cited articles were published between 1998 and 2014 (n=65). The most productive 5-year time period was between 2000 and 2005 when 23 of the 100 most cited publications were produced. Among retrieved citations, guidelines and recommendations comprised 5% of all articles, narrative reviews 21%, randomized clinical trials 12%, observational studies 35%, non-randomized clinical studies (basic and translational research) 24%, case series 1%, and meta-analyses 2%. Our analysis shows that articles defining the classification criteria of Systemic Sclerosis were among the most cited work followed by those describing the pathophysiology of the disease. Other highly cited areas included cellular and molecular mechanisms of fibrosis (with special emphasis on the role of TGF-β) and Randomized Controlled Trials for Bosentan and Epoprostenol in the treatment of pulmonary hypertension. Regarding individual organ manifestations, Pulmonary and Cardiac complications (pulmonary hypertension, lung disease, myocardial effects) (n=21) were of highest interest, while other articles focusing on Cutaneous (ulcers, skin changes, Raynaud's phenomenon) (n=16), Renal (renal crises) (n=3) and Gastrointestinal complications (dysmotility) (n=1) were cited to a lesser extent. Interestingly, articles primarily describing hematopoietic stem cell transplant as a viable treatment option did not feature on this list.

Conclusion: Our study provides the first bibliometric overview of the most highly cited scientific papers on systemic sclerosis. These findings provide useful insights into trends of published work and may serve as a useful guide to researchers in this field.

REFERENCES

- Cheng T, Zhang G. Worldwide research productivity in the field of rheumatology from 1996 to 2010: a bibliometric analysis. *Rheumatology* 2013; 52 (9): 1630-1634.
- Tetik-Aydogdu Y, Aydogdu O, Sari Z. AB1275 Research contribution to the journal of annals of the rheumatic diseases from 2012 to 2016: a bibliometric analysis. *Annals of the Rheumatic Diseases* 2018; 77(Suppl 2): 1731.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7744

SAT0278 STUDY OF QUALITY OF LIFE IN PATIENTS WITH SYSTEMIC SCLEROSIS- A CROSS-SECTIONAL STUDY

Girish Kakade¹, Rohini Samant¹, Ashok Mahashur², Romi Shah¹, Akash Khune¹, Amol Raut¹. ¹PD Hinduja National Hospital and Medical Research Centre, Rheumatology, Mumbai, India; ²PD Hinduja National Hospital and Medical Research Centre, Pulmonology, Mumbai, India

Background: There has been no study on quality of life (QoL) in patients with systemic sclerosis (SSc) in India.

Objectives: 1) To determine the difference in QoL in patients with SSc compared to general population. 2) To determine the difference in QoL in subsets of patients with SSc between a) Limited cutaneous SSc (lcSSc) and diffuse cutaneous SSc (dcSSc); b) SSc with and without interstitial lung disease (ILD) c) SSc with disease duration of ≤ 5 yrs and > 5 yrs.