clinical practice, scientific evidence of methotrexate on the different manifestations of psoriatic disease is limited. Specifically, data of MTX use in usual care and in early PsA patients are lacking.

Objectives: To describe the response to MTX for patients with PsA and an oligoarthritis or polyarthritis phenotype in usual care in the first year after diagnosis, and its impact on different aspects of the disease, including patients not achieving minimal disease activity (MDA) despite continuous exposure to MTX

Methods: Data collected in the Dutch southwest Early Psoriatic Arthritis cohort (DEPAR) study were used. Patients with a new diagnosis of PsA who have not started disease-modifying antirheumatic drugs (DMARDs) for PsA were eligible to participate. In this analysis, only patients with a phenotype at time of diagnosis of oligoarthritis or polyarthritis (resp. 2-4 joints involved or 5 or more joints, as defined by the rheumatologist), and at least one year follow up were included. Research nurses collected clinical data and data on medication use every three months in the first year after diagnosis. The following outcomes were used to evaluate disease activity: MDA, EULAR active disease (at least one tender and swollen joint), disease activity index for PsA low disease activity (DAPSA-LDA: DAPSA≤14), and disease activity score 28 low disease activity (DAS28-LDA: DAS28≤3.2). Disease activity at six months of patients who started MTX within six months after diagnosis and have continued using MTX until one year after diagnosis was analyzed.

Results: In April 2018, 219 patients had a phenotype of oligoarthritis (n=134, 61%) or polyarthritis (n=85, 39%) at time of diagnosis, and at least one year follow up. In 183 (84%) patients, MTX monotherapy was started within six months after diagnosis. Within the first year, 23 of these used MTX intermittently, 25 switched to a different DMARD, and 45 started a different DMARD while remaining on MTX therapy. The remaining 90 patients used MTX monotherapy throughout the first year and of these we assessed disease activity at 6 months after diagnosis. At 6 months, 38 patients out of these 90 were not in MDA despite continuous MTX treatment, while 44 achieved MDA. Major differences amongst the two groups included tender joint count, physical function, evaluation of pain and global disease activity and impact on health-related quality of life scores (Table 1). Conversely, 18% of all patients started MTX monotherapy were in sustained MDA at one year while continuing MTX monotherapy.

Conclusion: The majority of patients with a new diagnosis of PsA with a phenotype of oligoarthritis or polyarthritis started MTX within a year after diagnosis. MTX is maintained as monotherapy in half of these patients,

Table 1. Disease activity and health-related quality of life at six months after diagnosis, of patients on MTX monotherapy in the first year after diagnosis

	No MDA at 6 months (n=38)	MDA at 6 months (n=44)
Swollen joint count	0 (0-1)	0 (0-0.5)
Swollen joint count > 1	8 (21)	3 (7)
Tender joint count	2 (0-4)	0 (0-0.5)
Tender joint count > 1	21 (55)	3 (7)
Leeds enthesitis index	0 (0-2)	0 (0-0)
Leeds enthesitis index > 1	10 (26)	1 (2)
PASI > 1	22 (58)	15 (34)
VAS Pain > 15	36 (95)	10 (23)
VAS Global > 20	35 (92)	7 (16)
HAQ > 0.5	27 (71)	7 (16)
PsAID-12	3.8 ± 2.2	1.0 ± 1.0
PsAQoL	7.6 ± 4.9	1.2 ± 2.0
Short Form 36		
Physical functioning	56 ± 18	86 ± 12
Physical role functioning	44 ± 22	82 ± 19
Bodily pain	49 ± 14	75 ± 14
General health	49 ± 16	69 ± 14
Vitality	49 ± 19	72 ± 17
Social functioning	68 ± 22	93 ± 12
Emotional role functioning	64 ± 27	91 ± 15
Mental health	71 ± 19	84 ± 12
Physical component scale	37 ± 5.1	49 ± 5.2
Mental component scale	46 ± 10	55 ± 6.8

Data shown as n (%), median (interquartile range) or mean ± standard deviation
MDA: minimal disease activity; MTX: methotrexate; PASI: psoriasis area and severity index; VAS: visual analogue scale; HAQ: health assessment questionnaire; PsAID: psoriatic arthritis impact of disease; PsAQoL: psoriatic arthritis specific quality of life

with 18% of them achieving sustained MDA using MTX monotherapy at one year. Interestingly, approximately half of the patients who had continued methotrexate monotherapy were not achieving MDA after six months and they report higher impact of disease on clinical and patient reported outcomes, suggesting need to consider improvement of therapy for these patients.

Disclosure of Interests: Kim Wervers: None declared, Hannah den Braanker: None declared, Jolanda Luime: None declared, Ilja Tchetverikov: None declared, Andreas Gerards: None declared, Cathelijne Appels: None declared, Wiebo van der Graaff: None declared, Hans van Groenendael: None declared, Lindy-Anne Korswagen: None declared, Josien Veris-van Dieren: None declared, Johanna Hazes: None declared, Marc Kok: None declared, Marijn Vis Grant/research support from: Novartis

DOI: 10.1136/annrheumdis-2019-eular.5371

Crystal diseases, metabolic bone diseases other than osteoporosis

SAT0402 SEL-212 PHASE 2 CLINICAL STUDY IN SYMPTOMATIC GOUT PATIENTS: IMMATOR TOLERANTIC NANOPARTICLES COMBINED WITH PEGADRICASE MITIGATES IMMUNOGENICITY AND ENABLES SUSTAINED REDUCTION OF SERUM URIC ACID LEVELS, LOW RATE OF GOUT FLARES AND MONTHLY DOSING

Stephen Smolinski¹, Alan Kivitz², Wesley Dehaan¹, Lloyd Johnston¹, Rehan Azeem¹, Kei Kishimoto¹. ¹Selecta Biosciences Inc., Watertown, United States of America; ²Altoona Center for Clinical Research, Altoona, United States of America

Background: Pegylated uricases are promising therapies for the treatment of severe chronic gout, but are limited by their immunogenicity. We have previously shown that ImmTOR tolerogenic nanoparticles (formerly known as SVP-Rapamycin) encapsulating rapamycin co-administered with pegadricase prevented the formation of anti-drug antibodies (ADAs) in a dose-dependent manner. A prior Phase 1b study of SEL-212, a novel combination product candidate consisting of pegadricase and ImmTOR, demonstrated sustained control of serum uric acid (SUA) for at least 30 days after a single dose. Here we provide data from a recently completed Phase 2 multidose clinical trial.

Objectives: To assess data on the safety, tolerability, and effects on SUA, ADAs, and gout flares of five monthly doses of SEL-212 in symptomatic gout patients treated with 0.1 or 0.15 mg/kg ImmTOR in combination with 0.2 mg/kg pegadricase.

Methods: Patients with symptomatic gout (≥1 tophus, gout flare within 6 months, and/or gouty arthropathy) and elevated SUA (≥6 mg/dL) were enrolled in SEL-212 treatment cohorts. Patients reported here received up to five monthly doses of SEL-212 (0.2 mg/kg pegadricase combined with 0.1 or 0.15 mg/kg ImmTOR). Safety, tolerability, SUA, and ADAs were monitored, and clinical data were collected.

Results: As of 17 Dec 2018, 152 patients had been dosed in the Phase 2 study. All evaluable patients receiving 0.1 or 0.15 mg/kg ImmTOR administered with 0.2 mg/kg pegadricase who achieved three months of SUA control maintained SUA control in months four and five of combination treatment. Approximately 66% of evaluable patients maintained SUA levels below 6 mg/dL at week 20 after five monthly doses of SEL-212. The sustained reduction of SUA correlated with low or no ADAs. SEL-212 was generally well tolerated and associated with a low rate of gout flare rates. Only 35% of patients treated with five doses of 0.1-0.15 mg/kg ImmTOR, and 29% of all current patients in the SEL-212 Phase 2 trial, experienced gout flares after initiation during the first month of treatment with continued reduction of gout flare rates over months two through five. This low rate of gout flares appears to be in contrast with higher incidence of gout flares reported in clinical trials involving other urate lowering therapies.

Conclusion: SEL-212 has been well-tolerated, showing substantially reduced immunogenicity, sustained control of SUA, and low rate of gout flares with repeated monthly dosing. SEL-212 has a favorable product profile to address the unmet need of patients with severe, chronic gout

including low immunogenicity allowing continued dosing, lower reported flare rates and convenient monthly dosing.

Disclosure of Interests: Stephen Smolinski Shareholder of: Selecta Biosciences, Inc., Employee of: Selecta Biosciences, Inc., Alan Kivitz Shareholder of: Novartis, Consultant for: Abbvie, Janssen, Pfizer, UCB, Genzyme, Sanofi, Regeneron, Boehringer Ingelheim, Sun Pharma Advanced Research, Flexion., Paid instructor for: Celgene, Horizon, Merck, Novartis, Pfizer, Genzyme, Sanofi, Regeneron, Speakers bureau: Celgene, Horizon, Merck and Genetech, Flexion, Wesley DeHaan Shareholder of: Selecta Biosciences, Inc., Employee of: Selecta Biosciences, Inc., Lloyd Johnston Shareholder of: Selecta Biosciences, Inc., Employee of: Selecta Biosciences, Inc., Rehan Azeem Shareholder of: Selecta Biosciences, Inc., Employee of: Selecta Biosciences, Inc., Kei Kishimoto Shareholder of: Selecta Biosciences, Inc., Employee of: Selecta Biosciences, Inc.

DOI: 10.1136/annrheumdis-2019-eular.7769

SAT0403

SEX DIFFERENCES ARE PRESENT IN CLINICAL CHARACTERISTICS, BUT NOT IN RESPONSE TO DIFFERENT URATE LOWERING THERAPIES IN PATIENTS WITH GOUT

Sophie Wanten¹, Minke Ter Stal¹, Wing-Yee Kwok², Frank van den Hoogen^{1,3}, Marcel Flendrie¹, Noortje van Herwaarden^{1,3}. ¹Sint Maartenskliniek, Department of Rheumatology, Nijmegen, Netherlands; ²Rijnstate ziekenhuis, Department of Rheumatology, Arnhem, Netherlands; ³RadboudUMC, Department of Rheumatology, Nijmegen, Netherlands

Background: Clinical characteristics of gout differ between women and men¹. Little however is known about the association between these differences and response to treatment. As women seem to have lower mean uric acid excretion compared to men^{2,3}, response to uricosuric agents might be better compared to xanthine oxidase inhibitors.

Objectives: To identify sex differences in clinical characteristics and response to urate lowering therapy (ULT) in patients with gout and in women difference in response to allopurinol or benzbromarone.

Methods: Patients with clinical diagnosis of gout, a first outpatient visit between January 2010 and March 2018 and a follow-up of at least 6 months were included in a retrospective cohort study, with ongoing recruitment, conducted in two rheumatology centres in the Netherlands. From this cohort, patients who started ULT were selected. Clinical characteristics and treatment outcomes of allopurinol and benzbromarone were compared between women and men, including drug survival (corrected for age and renal function) and cumulative incidence of achieving target serum uric acid (<0.36 mmol/l). In women, difference in cumulative incidence of achieving target serum uric acid while using allopurinol or benzbromarone was compared.

Results: From a total of 519 (105 women/414 men) patients, 513 (104 women/409 men) and 74 (18 women/56 men) patients were included in the allopurinol and/or benzbromarone group, respectively. Clinical characteristics are described in Table 1. Drug survival was similar for women and men for allopurinol (hazard ratio 1.08; 95% confidence interval (CI) 0.71-1.64) as well as for benzbromarone (hazard ratio 0.66; 95% CI 0.26-1.66) (Figure 1a and b). Cumulative incidences of achieving target serum uric acid were 66% (69/104) and 73% (300/409) after allopurinol and 83% (15/18) and 82% (46/56) after

Table 1. Clinical characteristics

	Women (n = 105)	Men (n = 414)	p-value*
Age (years), median (IQR)	73.9 (54.3-71.6)	62.9 (54.3-71.6)	<0.01
Current alcohol use, n (%)	34 (45)	275 (79)	<0.01
Comorbidities, n (%)	71 (68)	200 (48)	<0.01
Hypertension	41 (39)	76 (18)	<0.01
Renal impairment	36 (34)	80 (19)	<0.01
Diabetes mellitus			
Diuretics use, n (%)	67 (64)	137 (33)	<0.01
Joint involvement, n (%)	14 (13)	68 (17)	0.38
Monoarthritis	64 (62)	220 (62)	
Oligoarthritis	26 (25)	120 (29)	
Polyarthritis			
History or presence of tophi, n (%)	39 (37)	94 (23)	<0.01
Crystal proven gout, n (%)	89 (85)	348 (89)	0.86
Baseline serum uric acid (mmol/L), mean (SD)	0.44 (0.13) (n=93)	0.43 (0.12) (n=360)	0.73

IQR = inter quartile range

* P-values for categorical variables were calculated by chi-square analysis, for continuous variables the appropriate (non)parametric analysis was used based on Gaussian distribution.

benzbromarone for women and men, respectively. Comparison in women between response to allopurinol and benzbromarone was not statistically significant (p = 0.15).

Conclusion: Clinical characteristics of gout differ between women and men, confirming previous studies, with women being older, having more comorbidities and using diuretics more often.

Although women have shown to be more uric acid under secretors compared to men, this does not seem to result in a difference in response to a xanthine oxidase inhibitor or an uricosuric agent.

REFERENCES

- [1] Harrold LR, et al. Ann Rheum Dis. 2006 Oct;65(10):1368-72.
- [2] Park YB, et al. Clin Rheumatol 2000;19(2):142-6.
- [3] te Kampe R. et al. Ann Rheum Dis.2018;77: Suppl A1637

Disclosure of Interests: Sophie Wanten: None declared, Minke ter Stal: None declared, Wing-Yee Kwok: None declared, Frank van den Hoogen: None declared, Marcel Flendrie Grant/research support from: Research grants from Grünenthal, Menarini, Consultant for: Yes, advisory board for Grünenthal in 2017, Paid instructor for: Yes, for Menarini, Noortje van Herwaarden: None declared

A

B

DOI: 10.1136/annrheumdis-2019-eular.930

SAT0404

PRETREATMENT AND CO-ADMINISTRATION WITH METHOTREXATE IMPROVED DURABILITY OF PEGLOTICASE RESPONSE: A PROSPECTIVE, OBSERVATIONAL, PROOF-OF-CONCEPT, CASE SERIES

John Botson¹, Jeff Peterson². ¹Orthopedic Physicians Alaska, Anchorage, United States of America; ²Western Washington Arthritis Clinic, Bothell, United States of America

Background: Pegloticase is a recombinant DNA-produced porcine-like uricase enzyme, which metabolizes relatively insoluble urate to highly soluble allantoin. It is used in the treatment of refractory gout which has