failed maximal medical management, typically with xanthine oxidase inhibitors (XOI). Studies have shown a complete responder rate of 42% when defined as repeat serum uric acid (sUA) levels <6.0 mg/dL for >80% of the time during months 3 and 6 of treatment. Patients that do not maintain a low uric acid while on pegloticase therapy are presumed to have developed anti-drug antibodies (ADA) which rapidly clear the pegloticase molecule. As is routinely utilized in the treatment of other rheumatologic diseases, coadministration of immunomodulatory medications, such as methotrexate, could potentially temper the development of these ADAs (as defined by maintenance of sUA response) in patients treated with pegloticase for refractory gouty arthropathy.

Objectives: The aim of the current case series was to identify and clinically evaluate patients in a real-world practice setting in order to investigate the utility of adding methotrexate to a pegloticase regimen to increase the durability of response.

Methods: In this prospective, proof-of-concept, observational case series, 10 sequential patients with refractory tophaceous gouty arthropathy being started on treatment with pegloticase 8 mg every two weeks (as per the label) were identified from 3 separate infusion centers. No inclusion/exclusion criteria were implemented or prescreening performed. Methotrexate (MTX) 15 mg orally once weekly and folic acid 1 mg orally once daily was started one month prior to the initial administration of pegloticase and continued throughout pegloticase treatment. Any infusion premedications, which were consistent with standard practices, were administered per the individual physician's discretion as well as management of any gout flares which occurred during the treatment course. As per standard of care, sUA was measured every two weeks, prior to each subsequent infusion. At the completion of pegloticase treatment for all patients, the number and percentage of patients able to maintain an sUA at goal <6.0 mg/dL was recorded.

Results: Ten patients ranging in age from 35-80 were identified, from 3 separate infusion centers. There were 143 total pegloticase infusions performed within the observation period. All 10 patients received at least 10 infusions (5 months), 9 patients at least 12 infusions (6 months), 5 patients at least 16 infusions (8 months), 2 patients at least 18 infusions (9 months), and 1 patient received 19 infusions. All 10 patients completed a full course of pegloticase treatment. All patients stayed on MTX 15 mg/week. There were no dose adjustments. 100% of patients were responders as defined by >80% of sUA levels being maintained at goal <6.0 mg/dL during the observation period. None of the 10 patients stopped pegloticase therapy due to increased sUA or loss of response and there were no infusion reactions in any of the 143 infusions or safety concerns identified. Gout flares did occur, primarily following the initial infusion, with less severity/prevalence with subsequent infusions. No patients discontinued treatment because of flares.

Conclusion: In this proof-of-concept case series of 10 sequential patients, pretreatment and co-administration of methotrexate 15 mg orally once weekly and folic acid 1 mg orally once daily with pegloticase resulted in a 100% maintenance of pegloticase sUA response with no infusion reactions. Although additional studies would be needed to corroborate these results, these data support a potential paradigm shift in treatment of refractory gout with pegloticase.

REFERENCE

- [1] Sundy JS, et al. JAMA. 2011;306:711

Acknowledgement: Research sponsored by the Alaska Rheumatology Alliance.

Disclosure of Interests: John Botson Shareholder of: Publicly traded stock: Horizon, Johnson and Johnson, Pfizer, Rigel, Consultant for: Celgene, Eli Lilly, Genentech, Horizon, Mallinckrodt, Novartis, Pfizer, Speakers bureau: Celgene, Eli Lilly, Horizon, Mallinckrodt, Novartis, Pfizer, Jeff Peterson Consultant for: Celgene, Genentech, Horizon, Lilly, Novartis, Regeneron (2017-2018), Speakers bureau: Abbvie, Horizon, Lilly, Novartis (2017-2018)

DOI: 10.1136/annrheumdis-2019-eular.3475

SAT0405 GOUT AND HYPERURICEMIA IN THE US: PREVALENCE AND TRENDS

Gurkirpal Singh¹, Bharathi Lingala², Alka Mithal¹. ¹ICORE, Woodside, United States of America; ²Stanford, Stanford, United States of America

Background: While the prevalence of gout and hyperuricemia is believed to be increasing worldwide, there is little population-based data in the US

after 2008. We evaluated the prevalence of gout and hyperuricemia in the US National Health and Nutrition Examination survey (NHANES) from 2007-08 to 2015-16, and compared these to previously published estimates to study whether the time-trends continue to show an increase in these rates.

Objectives: To study the prevalence of gout and hyperuricemia in the US population from 2007-08 to 2015-16.

Methods: We studied adults 20 years and older from the National Health and Nutrition Examination Survey (NHANES), beginning with the continuous NHANES 2-year survey cycles from 2007-08 to 2015-2016. The samples in each cycle were selected using a stratified, multistage, clustered probability sampling design. Persons with gout were identified from the home interview question "Has a doctor or other health professional ever told you that you had gout?" Hyperuricemia was defined as a serum urate level of greater than 6.8 mg/dl (supersaturation levels at physiological temperatures and pH), with alternate definitions of greater than 6.0 mg/dl (defined as the treat-to-target level in some treatment guidelines)(1). All statistical analyses were performed with appropriate survey procedures in SAS v9.4 (SAS Institute Inc., Cary, NC). Survey design variables and weight variables were used to account for the complexity of the NHANES survey design with stratification and clustering. Age-adjusted rates were calculated using 2010 US population numbers (direct standardization).

Results: In 2015-16, the overall prevalence of gout among US adults was 3.9%, corresponding to a total affected population of 9.2 million US adults. Gout was significantly more prevalent in men (5.2%) compared to women (2.7%), and the prevalence increased with age. Hyperuricemia (above 6.8 mg/dl) was seen in 14.6% of US population (estimated 32.5 million individuals), and was much more common in men (24.7%) compared to women (5.2%). The alternative definition of hyperuricemia (more than 6.0 mg/dl) was seen in 29.9% of individuals (estimated 66.6 million adults). Both hyperuricemia and gout were less prevalent in Mexican-Americans compared to Whites and African-Americans. No significant trends were identified in the age-adjusted prevalence of gout and hyperuricemia. Statistical comparisons between 2007-08 and 2015-16 age-adjusted rates were not significant.

Conclusion: While the age-adjusted prevalence of gout and hyperuricemia has remained unchanged in the most recent decade from 2007-08 to 2015-16, the estimated total number of persons with self-reported gout has gone up from 8.3 million to 9.2 million, reflecting the growth and increased aging of the US population. The age-adjusted prevalence of hyperuricemia has declined slightly but the total number of affected individuals is virtually identical (32.5 million in 2015-16 compared to 32.1 million in 2007-08). While the stabilization of gout and hyperuricemia prevalence rates is encouraging, our study highlights the still considerable burden of gout and hyperuricemia in the increasingly aging US population.

REFERENCE

- [1] Ruoff G, Edwards NL. Overview of Serum Uric Acid Treatment Targets in Gout: Why Less Than 6 mg/dL? Postgraduate medicine. Sep 2016;128(7):706-715.

Disclosure of Interests: Gurkirpal Singh Shareholder of: Merck, Pfizer, Grant/research support from: Horizon, Acorda, UCB, Consultant for: Horizon, Acorda, Sancio, Unipharm, Bharathi Lingala: None declared, Alka Mithal: None declared

DOI: 10.1136/annrheumdis-2019-eular.4438

SAT0406 COMPUTED TOMOGRAPHIC MAPPING OF SPINAL AND VASCULAR URATE DEPOSITION IN THE ABDOMEN WITH CORRELATION TO URIC ACID LEVEL

Juvel Lee¹, Waleed Abdellatif², Sunghan Jung¹, Ahmed Negida³, Bo Gong⁴, Savvas Nicolaou⁵. ¹Faculty of Medicine, University of British Columbia, Vancouver, Canada; ²University of British Columbia/Vancouver General Hospital, Radiology, Vancouver, Canada; ³Faculty of Medicine, Zagazig University, Zagazig, Egypt; ⁴Vancouver General Hospital, Vancouver, Canada; ⁵University of British Columbia/Vancouver General Hospital, Vancouver, Canada

Background: Gout is characterized by accumulation of monosodium urate (MSU) in joints and soft tissues. Gout prevalence is about 1-5% in the western world, predominantly in elderly men. Accurate diagnosis of gout is paramount in managing disease progression and guiding treatment. A meta-analysis in 2018 showed that Dual-Energy CT (DECT) has high sensitivity, specificity and diagnostic accuracy in MSU detection. MSU