comorbidities. Not believing in medication may impact on the success of achieving these treatment goals.

Objectives: To study which factors were associated to and predictive of beliefs about medicines during one year in patients with a recent gout attack and a need for ULT.

Methods: Baseline data from a prospective observational study were used in patients with crystal-proven gout after a recent gout flare and insufficiently treated serum urate (sUA) level ($>360 \mu\text{mol/L}$ / $>6 \text{ mg/dl}$). In these patients a treat-to-target approach was planned to meet the treatment target (sUA $<360 \mu\text{mol/L}$, or $<300 \mu\text{mol/L}$ if clinical tophi). Assessments included demographic and clinical data, baseline serum urate levels, medication, comorbidities, physical function (HAQ), and SF-36 mental (MCS) and physical component summary scores.

The Beliefs in Medicines Questionnaire (BMQ)(1) queries patient beliefs about medicines on four subscales (necessity and concerns specific for the patient, generally on overuse and harm). Respondents indicated their degree of agreement with each individual statement about medicines on a 5-point Likert scale, (1=strongly disagree to 5=strongly agree). Scores within the four subscales were summed (ranges 5-25 for necessity and concern, 4-20 for overuse and harm). Calculation of the difference between necessity and concern shows the relative importance of patients' view on taking medication, and was grouped as high and low relative to the median.

Using multivariate analysis with logistic regression, baseline variables were explored as predictors of high beliefs in the relative importance of medication after 12 months.

Results: 202 patients were included at baseline, 94.1% men, 90.3% Caucasian, mean (SD) age 56.6 (14.2) years, disease duration 8.0 (7.7) years. Mean sUA level was 494 (SD 86.8) $\mu\text{mol/L}$ at baseline, body mass index 28.9 (4.6) kg/m^2 , and physical function (HAQ) 0.36 (0.57). 18.8% (n=28) had tophi, and 30.4% (n=55) were previous users of allopurinol.

During the first year the beliefs in necessity of medication and the perceived importance of medication increased, while the concerns decreased (Table).

Table: Beliefs in Medicines questionnaire (BMQ) subscales. * $p<0.05$, ** $p<0.01$ vs. baseline

Month (n)	0 (195)	3 (158)	6 (144)	12 (126)
Necessity (SD)	17.1 (4.5)	18.2 (3.4)*	18.4 (3.4)**	18.5 (3.3)**
Concern (SD)	13.7 (4.8)	13.3 (4.5)	13.3 (4.6)*	13.1 (4.6)**
Nec – Conc (SD)	3.4 (6.2)	4.9 (5.4)*	5.1 (5.5)**	5.4 (5.5)**
Overuse (SD)	10.6 (2.8)	10.8 (2.5)	10.6 (2.8)	10.6 (2.7)
Harmful (SD)	9.4 (2.4)	9.5 (2.5)	9.6 (2.5)	9.2 (2.7)

Bivariate comparisons studying demographic, medication and clinical variables did not show consistent cross-sectional associations with beliefs in the importance of medication. However, after 12 months, bivariate comparisons revealed statistically significant differences in mental health (SF36 MCS), physical function (HAQ) and self-efficacy for symptom control.

In logistic regression analyses, adjusting for age, gender, BMI and comorbidities, prediction of high beliefs in the relative importance of medication after 12 months were independently associated with baseline self-efficacy for symptom control (OR 1.04 per unit, 95% CI 1.01-1.06).

Conclusion: During one year of treat-to-target strategy in patients with gout, perceived necessity of medication and higher importance of medication increases. Self-efficacy for symptom control at baseline was an independent predictor of high beliefs in importance of medication after 12 months. These findings support that patients increase their perception of the necessity of beliefs in medication during a treat-to-target approach with ULT.

REFERENCE

- [1] Horne R, et al. *Psychology & Health*. 1999. 14(1): 1–24.

Disclosure of Interests: Jon Anders Grønvd: None declared, Sebastian James Hekneby: None declared, Lars Fridtjof Karoliussen: None declared,

Espen A Haavardsholm Grant/research support from: research funding from Pfizer, UCB, Roche, MSD, and AbbVie., Tore K. Kvien Grant/research support from: AbbVie, BMS, MSD, Pfizer, Roche and UCB., Consultant for: AbbVie, Biogen, BMS, Boehringer Ingelheim, Celgene, Celltrion, Eli Lilly, Hospira, Merck-Serono, MSD, Novartis, Oktal, Orion Pharma, Pfizer, Roche, Sandoz, Sanofi, Mylan and UCB, Speakers bureau: AbbVie, Biogen, BMS, Boehringer Ingelheim, Celgene, Celltrion, Eli Lilly, Hospira, Merck-Serono, MSD, Novartis, Oktal, Orion Pharma, Pfizer, Roche, Sandoz, Sanofi and UCB, Hilde Berner Hammer Speakers bureau: speakers fee from AbbVie, Bristol-Myers

Squibb, Roche, UCB Pharma and Pfizer, Till Uhlig Consultant for: Grünenthal, Novartis, Speakers bureau: Grünenthal, Novartis

DOI: 10.1136/annrheumdis-2019-eular.4019

SAT0408

BODY FAT MEDIATES THE CORRELATION BETWEEN FRUCTOSE-CONTAINING BEVERAGES CONSUMPTION AND SERUM URIC ACID IN MALE GOUT PATIENTS

LI Qianhua, Lijuan Yang, Yan-Hui Xu, Chao Deng, Jinjian Liang, Jianzi Lin, Lefeng Chen, [Lie Dai](#). *Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Department of Rheumatology, Guangzhou, China*

Background: Fructose-containing beverages (FCB) consumption has been recognized as a promoter of hyperuricemia and gout in recent years. FCB consumption also contributes to the development of obesity which is an important risk factor of gout. However, data about the associations among FCB consumption, obesity and the level of serum uric acid (sUA) is limited.

Objectives: To investigate the associations among FCB consumption, obesity and sUA level in gout patients.

Methods: Consecutive gout patients who fulfilled the 2016 ACR/EULAR classification criteria were recruited. Demographic information, clinical characteristics and comorbidities of gout were collected. A 10-items food frequency questionnaire was developed which included alcohol, red meat, animal offal, seafood, hotpot, slow-cooking soup, FCB, tea, coffee and milk/milk products. Patients were asked to report the average consumption frequency over the one year prior to the first gout attack. Body composition was assessed by bioelectric impedance analysis and overfat was defined by body fat percentage (BF%) as $\geq 25\%$ for men and $\geq 35\%$ for women.

Results: 1) Among 331 recruited gout patients, 96.1% were male, so only male patients were further analyzed. The median age of male patients was 37.5 (30, 49) years and median sUA was 9.5 (7.8, 10.5) mg/dl with 18.9% presented tophi. The mean BF% was $25.6 \pm 6.2\%$ and 54.7% male patients were overfat. The prevalence of hypertension, diabetes, dyslipidemia, metabolic syndrome and fatty liver were 34.6%, 8.5%, 62.9%, 44.3% and 46.9% respectively. 2) There were no significant difference of food intake frequency between overfat patients and normal fat patients. Spearman correlation analysis results showed that FCB consumption was positively correlated with BF% ($r=0.176$, $P=0.002$), while milk/milk products consumption was negatively correlated with BF% ($r=-0.117$, $P=0.038$). After adjusted by age, duration, family history, eGFR, hypertension, diabetes mellitus, dyslipidemia, metabolic syndrome, fatty liver and the other nine dietary factors, multivariable linear regression showed that FCB consumption was positively correlated with BF% (compared with $<1 \text{ time/w}$, 3-4 times/w: coefficient $\beta=2.4619$, 95%CI 0.3580~4.5657, $P=0.022$; 4 times/w: coefficient $\beta=2.6696$, 95%CI 0.8266~4.5726, $P=0.0049$), and BF% was positively correlated with sUA (coefficient $\beta=0.0784$, 95%CI 0.0328~0.1239, $P<0.001$). However, FCB consumption was not correlated with sUA after adjusted by above confounders and BF% ($P=0.079$). 3) Further mediation analysis were performed to evaluate whether BF% mediated the correlation between FCB consumption and sUA. After adjusted by the confounding factors mentioned above, the total effect of FCB consumption on sUA was significant (compared with $<1 \text{ time/w}$, $>4 \text{ times/w}$: coefficient $\beta=0.8813$, 95%CI 0.1457~1.6169, $P=0.019$). BF% produced indirect effect on the correlation between FCB consumption and sUA (compared with $<1 \text{ time/w}$, $>4 \text{ times/w}$: indirect effect coefficient $\beta=0.2115$, 95%CI 0.0453~0.4032).

Figure 1 Linear regression analysis of the correlation between food intake frequency and BF% and sUA in male gout patients

Food intake frequency	All patients (n=338)	Correlation with BF%				Correlation with sUA			
		Coefficient β	LCI	UCI	P	Coefficient β	LCI	UCI	P
Alcohol									
None	141(44.3)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-20g/d	89(26)	-1.0466	-2.5604	0.4711	0.1760	-0.3635	-0.9208	0.2429	0.2408
21-40g/d	39(12.3)	0.2762	-1.8168	2.3692	0.7953	0.2723	-0.5348	1.0793	0.5072
>40g/d	49(15.4)	0.7476	-1.3134	2.8086	0.4758	0.3947	-0.4607	1.1910	0.4082
Red meat									
0-100g/d	102(32.1)	0.0	NA	NA	NA	0.0	NA	NA	NA
101-200g/d	149(46.8)	-0.9870	-2.4907	0.5567	0.2126	-0.2361	-0.8253	0.3530	0.4307
201-300g/d	52(16.4)	-0.6439	-2.3878	1.0900	0.1349	0.3138	-0.4343	1.0519	0.4079
>300g/d	15(4.7)	0.9134	-2.2809	4.1076	0.5740	0.1041	-1.1282	1.3364	0.8680
Animal oil									
<1 times/w	227(71.4)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	65(20.4)	0.5580	-1.2227	2.3388	0.5378	-0.5776	-1.2646	0.1095	0.0991
3-4 times/w	15(4.7)	-0.8508	-0.9278	2.2261	0.5867	-0.7521	-1.5994	0.4946	0.2131
>4 times/w	11(3.3)	-2.5650	-5.1940	1.0028	0.1650	-10.1541	-0.2499	2.5551	0.3060
Seafish									
<1 times/w	215(67.6)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	59(16.6)	0.7886	-0.3244	2.4915	0.5672	-0.1258	-0.7203	0.5297	0.6587
3-4 times/w	25(7.8)	-0.9334	-3.3759	1.5091	0.4525	-0.0335	-0.9762	0.9092	0.9443
>4 times/w	19(6)	-2.5593	-5.3741	0.2555	0.0746	-0.1390	-1.2305	0.9525	0.8023
Veget									
<1 times/w	287(90.3)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	23(7.2)	0.7567	-1.7281	3.2415	0.5493	-0.1055	-1.0642	0.8532	0.8286
3-4 times/w	4(1.3)	-1.0208	-6.7962	4.7943	0.7248	0.1551	-2.0547	2.3650	0.8902
>4 times/w	4(1.3)	-1.0445	-6.9190	4.8299	0.7246	-0.0671	-2.3326	2.1983	0.9535
Slow-cooking soup									
<1 times/w	129(40.6)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	99(31.1)	-0.1611	-1.6797	1.3576	0.8348	-0.3682	-0.9538	0.2174	0.2169
3-4 times/w	52(16.4)	-1.1055	-3.0452	0.8342	0.2628	-0.1888	-0.9383	0.5608	0.6705
>4 times/w	38(11.9)	0.7027	-1.4212	2.8265	0.5354	-0.3409	-1.1604	0.7786	0.4135
Fructose containing beverages									
<1 time/w	168(52.8)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	44(13.8)	1.4198	-0.5328	3.3725	0.1534	0.0229	-0.7318	0.7795	0.9504
3-4 times/w	41(12.9)	2.4619	0.3580	4.5657	0.0220	-0.0888	-0.9077	0.7301	0.8311
>4 times/w	95(20.4)	2.8996	0.8266	4.9726	0.0049	0.6695	-0.021	1.4023	0.0730
Tea									
<1 times/w	147(46.2)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	34(10.7)	-0.5081	-2.7650	1.6789	0.6308	0.0261	-0.8309	0.8832	0.9522
3-4 times/w	14(4.4)	0.9783	-2.2133	4.1700	0.5407	-0.1891	-1.4205	1.0423	0.7626
>4 times/w	123(38.7)	-0.1466	-1.5524	1.2592	0.5975	0.2479	-0.2942	0.7900	0.3688
Coffee									
<1 times/w	293(92.1)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	15(4.7)	-0.9200	-4.9628	1.1228	0.1322	-0.3609	-1.2162	0.2244	0.1130
3-4 times/w	10(3.1)	-0.7054	-10.4630	11.0761	0.9121	10.1822	3.1249	5.4884	0.5894
>4 times/w	9(2.8)	-1.6474	-6.7000	2.4052	0.4243	0.2745	-1.2899	1.8388	0.7801
Milk and milk products									
<1 times/w	173(54.4)	0.0	NA	NA	NA	0.0	NA	NA	NA
1-2 times/w	95(29.4)	-0.5368	-1.9924	0.9188	0.4885	0.3462	-0.2155	0.9080	0.2760
3-4 times/w	27(8.5)	-1.7837	-4.0981	0.5407	0.320	-0.2811	-1.1810	0.6189	0.5991
>4 times/w	23(7.2)	0.9840	-2.4735	2.5519	0.965	0.0809	-0.8857	1.0436	0.8575

Multivariate, adjusted for potential confounding factors including age, duration of gout, family history, eGFR, comorbidities (including hypertension (yes/no), diabetes mellitus (yes/no), dyslipidemia (yes/no), metabolic syndrome (yes/no), and fatty liver (yes/no)) and the other dietary factors.

Multivariate, adjusted for the confounding factors mentioned above and BF%.

LCI, lower limit of confidence interval; UCI, upper limit of confidence interval; NA, not applicable; time/w, the frequency of dietary intake per week; BF%, body fat percentage; sUA, serum uric acid.

Conclusion: Our data show that body fat mediates the correlation between FCB consumption and sUA in male gout patients, which imply the importance of body composition assessment.

Acknowledgement: The present study was supported by Guangdong Natural Science Foundation, China (Grant no. 2014A030310086) to Qian-Hua Li.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5774

SAT0409

P2X7R PROMOTE THE ATTACK OF ACUTE GOUTY ARTHRITIS IN RATS FROM CLINICAL TO PATHOLOGY

Xiaojuan Dai¹, Li Xiangpei², Jinhui Tao², Xuan Fang², Yuan Xia², Li Xiaomei².

¹The First Affiliated Hospital of University of Science and Technology of China, Rheumatology and Immunology, Hefei, China; ²The First Affiliated Hospital of University of Science and Technology of China, Rheumatology and Immunology, Hefei, China

Background: ATP may be the second causative signal for the onset of gout, which acts on P2X7R to regulate the development of acute gouty arthritis (1). P2X7R is the most relevant to the inflammatory response in the P2X family and is also an ATP-gated cation channel (2).

Objectives: The aim of this study was to activate P2X7R signaling pathway through changes in extracellular ATP concentrations, leading to the development of acute gouty arthritis and the production of pro-inflammatory.

Methods: 120 male Sprague-Dawley rats were randomly divided into 3 groups: After establishment of acute gouty arthritis model, rats were given P2X7R agonist ATP, P2X7R inhibitor BBG and PBS, respectively. The clinical manifestations of the ankle joints were evaluated at 6h, 12h, 24h, 48h and 72h before the rats were sacrificed, and rat ankle synovial slices for H&E staining. IL-1 β , IL-6 and TNF- in the serum of rats were detected by ELISA kits.

Results: 1. P2X7R regulates the development of acute gouty arthritis: At 12h, the clinical scores of ATP group were significantly higher than those of BBG group and control group (P=0.001,0.042), and the control group was higher than BBG group (P=0.034). At 12h and 24h, the swelling index of ATP group was the most obvious than other two groups (P=0.000, 0.001; P=0.000, 0.003), followed by control group (P=0.009, 0.001); Furthermore, there was a large infiltration of

inflammatory cells in the synovial tissue of the right ankle joint of rats, at 12h, and 24h, the infiltration of mononuclear cell in ATP group was significantly higher than that in BBG group and control group (P=0.000, 0.007; P=0.000, 0.001); The neutrophils infiltration in ATP group was the highest among the three groups at 24h (P=0.001, 0.04), and the control group was higher than BBG group (P=0.04). 2. P2X7R regulates pro-inflammatory cytokines production: At 24h, the level of IL-1 β in ATP group was significantly higher than BBG group and control group (P=0.001, 0.003); At 6h, 12h and 24h, higher level of IL-6 in ATP group compared with BBG group and control group (P=0.004, 0.04; P=0.000, 0.002; P=0.001, 0.012); The level of TNF- was obviously higher in ATP group than in BBG group and control group at 6h and 24h (P=0.007, 0.011; P=0.001, 0.018), and BBG group was lower than control group, but without statistically significant (P>0.05).

Conclusion: Activation of P2X7R can significantly promote the attack of acute gouty arthritis and the production of IL-1 β , IL-6 and TNF-, suggesting that P2X7R affects the development of acute gouty arthritis and regulates the secretion of pro-inflammatory cytokines.

REFERENCE

- [1] Tao JH, Zhang Y, Li XP. P2X7R: a potential key regulator of acute gouty arthritis. *Semin Arthritis Rheum*, 2013, 43(3): 376-380. (2) Hechler B, Gachet C. Purinergic receptors in thrombosis and inflammation[J]. *Arterioscler Thromb Vasc Biol*, 2015, 35:2307-2315

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7147

SAT0410

RADIOGRAPHIC EROSION SCORE IMPROVED WITH TARGETED URATE-LOWERING THERAPY IN A PROSPECTIVE GOUT VIETNAMESE COHORT

Thomas Bardin¹, Quang Dinh Nguyen², Khoi Minh Tran², Nghia Hieu Le², Duc Minh Do³, Valérie Bousson⁴, Pascal Richette¹, Matthieu Resche-Rigon⁵. ¹Hôpital Lariboisière, Rheumatology, Paris, France; ²Vien Gut Medical Center, French-Vietnamese research center on gout and chronic diseases, HO Chi Minh City, Vietnam; ³University of Medicine and Pharmacy, Molecular Biology, Ho Chi Minh City, Vietnam; ⁴Hôpital Lariboisière, Radiology, Paris, France; ⁵Hôpital Saint Louis, Biostatistics, Paris, France

Background Urate deposition in joints of patients with neglected gout leads to destructive arthropathy, with subchondral bone erosion, bone construction, and late joint space narrowing. Gouty erosions are believed to improve under urate lowering drugs (ULDs), but their course has been little studied. We therefore performed a systematic prospective study of a Vietnamese cohort started on allopurinol at inclusion.

Objectives To characterize radiographic outcome of gouty erosions under serum urate-targeted ULD

Methods Included patients had crystal-proven gout, eGFR > 60 ml/min and were not taking ULDs at inclusion. Allopurinol was progressively increased to reach EULAR- recommended uricemia targets and patients were prospectively followed-up, in particular with sequential hand and foot antero-posterior radiographs and MTP and knee ultrasound (US) scans. Joint erosions were scored according to a validated index1, double contour thickness was evaluated by a 4 point-score. One index tophus per patient (hand or foot) was measured by US scan.

Results 100 patients (99M) with a mean age of 45.9 + 9.9 years, and median gout duration of 8 years (range 0-33) were included. 91 had clinical tophi. Mean baseline uricemia was 492 μ mol/L. Median final allopurinol dose was 520 mg/day. 71 patients had evidence of urate arthropathy at baseline, 68 of the feet and 23 of the hands (all but 3 with foot involvement). 1st MTPs were predominantly involved (44 and 35 patients on the right and left sides respectively). Urate arthropathy at baseline correlated with gout duration (p=0.01), clinical tophi (p=0.0007), maximum double contour thickness and index tophus US measurement (p=0.03) by univariate analysis and with clinical tophi by multivariable analysis (p=0.03). Uricemia target <300 μ mol/L was obtained in 63.6, 67.8, 70.6, and 61.0 % of patients at 3, 6, 12 and 24 months respectively. Median radiological follow-up was 12 months (range 3-49). Mean radiological erosion score1 was 12.1 at baseline (n=71), with a decrease of -0.7 at 6 months (n=38,