research support from: Celgene, Frances Humby: None declared, Costantino Pit-
zalis Grant/research support from: Celgene

DOI: 10.1136/annrheumdis-2019-eular.3634

OP0114

MACHINE LEARNING TOOLS IDENTIFY PATIENT CLUSTERS AND SWOLLEN AND TENDER JOINT CORRELATION PATTERNS IN A LARGE DATABASE FROM THE SECUKINUMAB PSORIATIC ARTHRITIS CLINICAL DEVELOPMENT PROGRAM

Matthias Kormaksson¹, Effie Pournara², Gregory Ligozio¹, Luminita Pricop¹, Ken Abrams¹, Bruce Kirkham³, Kristian Reich⁴, Iain McInnes⁵. ¹Novartis Pharmaceutical Corporation, New Jersey, United States of America; ²Novartis Pharma AG, Basel, Switzerland; ³Guy's and St Thomas' NHS Foundation Trust, London, United Kingdom; ⁴University Medical Center Hamburg-Eppendorf, Skin Inflammation Center, Hamburg, and Dermatologikum, Berlin, Germany; ⁵University of Glasgow, Glasgow, United Kingdom

Background: Identifying patient phenotypes using machine-learning (ML) techniques amidst the variability and heterogeneity of the clinical manifestations of psoriatic arthritis (PsA) could be the first critical step towards better understanding of the disease eventually leading to individualized medicine.¹

Objectives: To identify distinct clusters of patients with PsA based on patients' tender joint (TJ) and swollen joint (SJ) counts and correlation patterns among TJ and SJ counts at baseline as captured in the secukinumab FUTURE trials program.

Methods: Pairwise correlations were explored among 76 SJ and 78 TJ measurements of >2,700 patients with PsA across 5 phase III studies with ~425,000 data entries at baseline and were visualized using heatmaps. Due to high correlations between SJs and corresponding TJs, a composite variable "swollen/tender joint count" was constructed for each joint. Hierarchical clustering was then performed on the composite using "1-correlation" as the dissimilarity metric and Ward's agglomeration method for pairwise grouping of joints. A dendrogram was used to visualize and assess the resulting joint groupings.

Results: The hierarchical clustering algorithm grouped the 78 individual joints into distinct and natural clusters (Figure 1A). At higher level of the dendrogram, the algorithm grouped separately all foot, larger (jaws, clavicles, ankles, hips, wrists, knees, shoulders, elbows), and hand joints. Cutting the dendrogram at 15 clusters separated all the joints into distinct groups; hand joints (distal and proximal phalanges, metacarpals and thumbs), and foot joints (distal and proximal phalanges, metatarsals and big toes). Similar clustering algorithms were explored to identify patient clusters at baseline with distinct swelling and tenderness patterns across the identified joint groups. High correlation between swelling/tenderness of the left and swelling/tenderness of the corresponding right joint was observed across all individual joints (Figure 1B); a high correlation was also observed between swelling and tenderness at all individual joints. More localized patterns showed that there is a gradual decrease in correlation (from highest to lowest) among TJs and SJs in adjacent vs non-adjacent fingers, which is evident from

grey-scale patterns (Figure 1C). Specifically, a gradual decrease in correlation between the swelling of 2nd distal interphalangeal joint and the tenderness of the 2nd-5th distal phalanges was noted.

Conclusion: Machine learning methodology confirmed a natural grouping of joints in patients with psoriatic arthritis based on baseline swelling and tenderness and revealed complex correlation patterns. Additional cluster analyses have demonstrated distinct patient clusters across the identified joint groups. Further investigating potential associations of other disease manifestations such as skin and nail involvement to define additional phenotypes may explain differences in disease pathogenesis and treatment outcomes.

REFERENCE:

[1] Grys BT, et al., J Cell Biol. 2017; 216(1): 65-71.

Disclosure of Interests: Matthias Kormaksson Shareholder of: Novartis, Employee of: Novartis, Effie Pournara Shareholder of: Novartis, Employee of: Novartis, Gregory Ligozio Shareholder of: Novartis, Employee of: Novartis, Luminita Pricop Shareholder of: Novartis, Employee of: Novartis, Ken Abrams Shareholder of: Novartis, Employee of: Novartis, Bruce Kirkham Grant/research support from: Abbvie, Janssen, Lilly, Novartis, Roche, UCB, Consultant for: Abbvie, Janssen, Lilly, Novartis, Roche, UCB, Speakers bureau: Abbvie, Janssen, Lilly, Novartis, Roche, UCB, Kristian Reich Consultant for: Abbvie, Affibody, Amgen, Biogen, Boehringer Ingelheim Pharma, Celgene, Centocor, Covagen, Forward Pharma, Fresenius Medical Care, GlaxoSmithKline, Janssen-Cilag, Kyowa Kirin, Leo, Lilly, Medac, Merck Sharp & Dohme Corp., Novartis, Miltenyi Biotec, Ocean Pharma, Pfizer, Regeneron, Samsung Bioepis, Sanofi, Takeda, UCB Pharma, Valeant, Xenoport; Speakers bureau: Abbvie, Affibody, Amgen, Biogen, Boehringer Ingelheim Pharma, Celgene, Centocor, Covagen, Forward Pharma, Fresenius Medical Care, GlaxoSmithKline, Janssen-Cilag, Kyowa Kirin, Leo, Lilly, Medac, Merck Sharp & Dohme Corp., Novartis, Miltenyi Biotec, Ocean Pharma, Pfizer, Regeneron, Samsung Bioepis, Sanofi, Takeda, UCB Pharma, Valeant, Xenoport, Iain McInnes Grant/research support from: AstraZeneca, Celgene, Compugen, Novartis, Roche, UCB Pharma, Consultant for: Abbvie, Celgene, Galvani, Lilly, Novartis, Pfizer, UCB Pharma

DOI: 10.1136/annrheumdis-2019-eular.1910

OP0115

GENERAL AND SEX-SPECIFIC PREDICTORS OF PSA AMONG PATIENTS WITH PSORIASIS

Alexis Ogdie¹, Jose Scher², Shiyu Vanessa Wang¹, Daniel Shin¹, David Margolis¹, Junko Takeshita¹, Hyon Choi³, Thorvardur Jon Love⁴, Christopher T. Ritchlin¹, Joel Gelfand¹, Joseph F. Merola⁵. ¹University of Pennsylvania, Philadelphia, PA, United States of America; ²New York University, New York, NY, United States of America; ³Massachusetts General Hospital, Boston, MA, United States of America; ⁴University of Iceland, Reykjavik, Iceland; ⁵University of Rochester, Rochester, NY, United States of America; ⁶Brigham Women's Hospital, Boston, MA, United States of America

Background: Risk prediction models in electronic health record (EHR) databases may assist in early identification of patients with psoriasis likely to develop psoriatic arthritis (PsA).¹ A better understanding of potential predictors and whether stratification by sex would be needed in building such algorithms is required.¹

Objectives: Examine general and sex-specific predictors of PsA in an EHR dataset among patients with psoriasis

Methods: A retrospective cohort study was performed within the OptumInsights EHR Database (United States) between 2006-2017. Patients with two or more ICD codes for psoriasis and ages 16-90 were identified. The outcome was PsA (defined by a single ICD code). Potential predictors, in particular comorbidities and infections, were also identified using ICD codes. Hazard ratios were calculated using Cox proportional hazards models between individual predictors and development of incident PsA in univariate models and those that were significant (p<0.1) were entered into a multivariable model. A final model was achieved using automated stepwise regression. Separate models were developed for each sex as some predictors (e.g., polycystic ovarian syndrome, prostatitis) are sex-specific.

Results: Among 215,386 patients with psoriasis, mean age was 50 (SD 15.6) and 55% were female. At index date (one year after date of first psoriasis code), 4.6% and 4.2% of patients had been prescribed a biologic therapy or oral therapy in the past year. Mean follow up time was 5.6 years (SD 2.8) and 4,288 patients developed incident PsA (incidence 3.5 cases/1,000 person years). Previously identified predictors were significant in univariate models (depression, fatigue, inflammatory bowel disease, uveitis, hyperlipidemia, fracture; data not shown due to space restrictions) but several new predictors were also identified (diabetes, hidradenitis suppurativa, celiac disease, irritable bowel syndrome, sepsis, post-traumatic stress disorder, anxiety, anemia) (Table). Automated regression identified subsets of these factors in multivariable models; these models differed by sex.

Conclusion: Predictors of developing PsA differed by sex but obesity, depression, and fatigue were statistically significant predictors in both groups. Infections were also associated with development of PsA but the type of infection differed by sex.

Figure 1. Summary of Results

A. Dendrogram of the hierarchical clustering of 78 joints; bottom-most "leaves" correspond to the individual joints. Cutting the dendrogram at 15 clusters (grey dashed horizontal line) results in the distinct joint groups ("branches") labeled below the dendrogram.
B. Correlation heatmaps depicting pairwise correlations of all 76 swollen joints and 78 tender joints.
C. Pairwise correlations between swollen finger joints of the right hand and corresponding tender joints; darker/lighter grey corresponds to higher/lower correlations.
TEN: tender; SWO: swollen; R: right; MCP: metacarpal; PIP: proximal; DIP: distal phalanges; numbers 1-5: fingers 1 (thumb) through 5 (little)