SAT0531

COLLAGEN IMAGING USING DUAL-ENERGY COMPUTED TOMOGRAPHY CAN DETECT SOFT-TISSUE INVOLVEMENT OF THE INTERVERTEBRAL DISC IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS

Torsten Diekhoff¹, Kay-Geert Hermann¹, Fabian Proft², Mikhail Protopopov², Marcus R. Makowski¹, Denis Poddubnyy², Arthritis Imaging Research Group - Berlin. ¹Charité – Universitätsmedizin Berlin, Department of Radiology, CCM, Berlin, Germany; ²Charité – Universitätsmedizin Berlin, Department of Rheumatology, CBF, Berlin, Germany

Background: Based on the unique physical properties of collagen, dual-energy computed tomography (DECT) is able to detect pathological changes in fibrous structures. This has been demonstrated for the Achilles tendon (1) and traumatic disc injury in elderly patients (2).

Objectives: The purpose of this study was to assess the diagnostic potential of DECT collagen imaging for the detection of intervertebral disc involvement in patients with axial spondyloarthritis (axSpA) in correlation with osseous lesions in MRI.

Methods: Sixteen patients with suspicion of or known axSpA and back pain were included. All patients underwent a 1.5-Tesla-MRI including T1 and STIR sequences and an ultra-low-dose DECT of the spine using two sequential helical acquisitions. DECT images were reconstructed in 120 kV-equivalent standard CT images and collagen maps with 1.0 mm slice thickness in sagittal reconstructions. MRI and CT images (D1 to S1) were scored for transdiscal ankylosis, Andersson lesions, syndesmophytes, spondylitis anterior and fatty corner lesions (when applicable) as well as for degenerative findings. DECT images were scored for a loss of collagen density affecting the nucleus pulposus or the annulus fibrosus. Sensitivity (SE) and specificity (SP) values were calculated. All analyses were performed on the level of the disco vertebral units.

Results: Twelve male and four female patients with a mean age of 46.6 ± 11.6 years were included. Eleven were finally diagnosed with axSpA, 3 with degenerative spine disease and 2 with diffuse idiopathic skeletal hyperostosis (DISH). MRI detected 73 lesions in 274 discovevertebral units (61 SpA related, 12 degenerative; 26 affecting the disc and 35 affecting the vertebral corner) and CT 68, whereas DECT identified pathologically decreased collagen content in 60 discs (35 in the nucleus and 25 in the annulus). DECT showed high diagnostic accuracy when compared to SpA-related and degenerative changes (MRI and CT combined; SE: 67%; SP 99.5%) and for SpA-lesions only (SE 72%; SP 94.2%), however, was less susceptible for degeneration (SE 60%; SP 81.9%). When comparing disc lesions in SpA as detected by MRI (transdiscal ankylosis and Andersson lesions) with nucleus-pulposus changes in DECT, it showed an SE of 88.5% and an SP of 95.1%. However, the diagnostic accuracy was inferior when comparing anterior spondylitis and fatty corner lesions in MRI with annulus fibrosus changes in DECT (SE 51.4% and SP 97%), although the SE was markedly higher for active inflammatory corner lesion (92.3%) than for fatty marrow lesions (23.3%). The mean radiation exposure was 8.1 ± 3.4 mSv.

Conclusion: DECT is able to demonstrate soft-tissue involvement of the disc, especially for transdiscal ankylosis, Andersson lesions and active anterior spondylitis. This can be achieved with considerably low radiation exposure. While measuring the collagen density, it provides additional information to conventional CT and MR images. Thus, it has high potential to develop into a useful tool to further enhance our understanding of pathologic soft-tissue changes in axSpA patients outside the bone and bone marrow.

REFERENCES

- [1] Mallinson PI, Stevens C, Reisinger C, Nicolaou S, Munk PL, Ouellette H. Achilles tendinopathy and partial tear diagnosis using dual-energy computed tomography collagen material decomposition application. J Comput Assist Tomogr.
- [2] Pumberger M, Fuchs M, Engelhard N, et al. Disk injury in patients with vertebral fractures-a prospective diagnostic accuracy study using dual-energy computed tomography. Eur Radiol. 2019.

Disclosure of Interests: Torsten Diekhoff Paid instructor for: MSD, AbbVie, Novartis, Canon, but less than 10.000€, Kay-Geert Hermann Paid instructor for: MSD, AbbVie, Novartis, but less than 10.000€, Fabian Proft Grant/research support from: Novartis, Consultant for: yes but less than 10.000, Paid instructor for: yes but less than 10.000, Speakers bureau: yes but less than 10.000, Mikhail Protopopov: None declared, Marcus R Makowski: None declared, Denis Poddubnyy Grant/research support from: AbbVie, Merck Sharp & Dohme, Novartis, Consultant for: AbbVie, Bristol-Myers Squibb, Janssen, Merck Sharp & Dohme, Novartis, Pfizer, UCB Pharma, Speakers bureau: AbbVie, Bristol-Myers Squibb, Janssen, Merck Sharp & Dohme, Novartis, Pfizer, Roche, UCB Pharma
DOI: 10.1136/annrheumdis-2019-eular.6602

SAT0532

MRI DETECTS SUBCLINICAL PLANTAR FASCIA AND ACHILLES TENDON ENTHESTITIS IN PATIENTS WITH PSORIATIC ARTHRITIS, PSORIASIS AND ANKYLOSING SPONDYLITIS

Nienke J. Kleinrensink^{1,2,3}, Emmerik Leijten^{1,2}, Wouter Foppen³, Suzanne C. Diepstraten³, Nanette L. Vincken^{1,2}, Pim de Jong³, Timothy R. Radstake^{1,2}. ¹University Medical Center Utrecht, Department of Rheumatology and Clinical Immunology, Utrecht, Netherlands; ²University Medical Center Utrecht, Laboratory of Translational Immunology, Utrecht, Netherlands; ³University Medical Center Utrecht, Department of Radiology, Utrecht, Netherlands

Background: Enthesitis is thought to be a key feature driving the development of spondyloarthritis, although head-to-head studies in patients with psoriasis (Pso) versus psoriatic arthritis (PsA) are scarce. In PsA and ankylosing spondylitis (AS) patients the plantar fascia and Achilles tendon are among the enthesal structures most frequently affected clinically. MRI allows for objective detection of enthesitis in patients with rheumatic diseases.

Objectives: To assess the frequency of MRI-detected enthesitis at the plantar fascia and Achilles tendon in Pso, PsA and AS patients.

Methods: The cross-sectional observational 'EXTEND-UP' study* included patients with Pso (diagnosed by a dermatologist, PsA excluded by rheumatologist), PsA (fulfilling CASPAR criteria) and AS (fulfilling ASAS criteria). With the exception of two PsA patients, no study participants received DMARDs. Study participants were included regardless of the presence/absence of clinically-suspect enthesitis (nail blanching pressure on enthesitis). Clinicians were blinded to the MRI-scan results. 3 Tesla MRI-scans of both ankles were performed and evaluated by one observer with over four years of experience in musculoskeletal MRIs, who was blinded to diagnosis and clinical examination. MRI-defined enthesitis was scored following the OMERACT MRI definitions for PsA.

Results: This study evaluated a total of 79 ankles from 13 Pso patients, 15 PsA patients and 12 AS patients. The MRI-scan of one ankle was not assessable for enthesitis because of suboptimal fat suppression. Overall, MRI-defined enthesitis either at the plantar fascia and/or Achilles tendon was detected in 30% (9/30) of PsA ankles, compared to 8% (2/26) of Pso ankles ($p < 0.05$, Fisher's exact test). Specifically, MRI-detected enthesitis at the plantar fascia(s) occurred in 0% (0/13) of Pso patients, 20% (3/15) of PsA patients and 8% (1/12) of AS patients. MRI-detected enthesitis at the Achilles tendon(s) occurred in 8% (1/13) of Pso patients, 33% (5/15) of PsA patients and 0% (0/12) of AS patients. Agreement between clinically-suspect versus MRI-detected enthesitis was moderate ($\kappa = 0.41$) regarding plantar fascia enthesitis (table 1). There was a poor concordance ($\kappa = 0.12$) between clinically-suspect versus MRI-detected Achilles tendon enthesitis, where 8 out of 9 Achilles tendons that were judged as normal in physical examination, did show inflammation on MRI (table 1).

Conclusion: This study shows a substantial incidence of MRI-detected plantar fascia and Achilles tendon enthesitis in PsA, occurring more frequently compared to Pso. The clear discrepancy between clinically-suspect enthesitis and MRI-defined enthesitis, even at prototypical sites, warrants caution before using clinical evaluation of enthesitis as part of clinical decision-making or trial outcome measures.

Table 1. MRI-detected enthesitis versus clinical examination

Clinical examination for enthesitis	Plantar fascia MRI		κ-value
	No enthesitis (n = 74)	Enthesitis (n = 5)	
Normal, no (%)	72 (97)	3 (60)	0.41
Clinically suspect, no (%)	2 (3)	2 (40)	
Clinical examination for enthesitis	Achilles tendon MRI		κ-value
	No enthesitis (n = 70)	Enthesitis (n = 9)	
Normal, no (%)	68 (97)	8 (89)	0.12
Clinically suspect, no (%)	2 (3)	1 (11)	

*=This investigator-initiated study was funded by Janssen Inc.

Figure 1. Plantar fascia (A & B) and Achilles tendon (C & D) with increased T2 signal reflecting edema (arrowhead) and contrast enhancement (arrow) around the entheses.

Disclosure of Interests: Nienke J. Kleinrensink: None declared, Emmerik Leijten: None declared, Wouter Foppen Grant/research support from: WF has received research grants from NovoNordisk and Pfizer. All were paid to the institution.

Paid instructor for: WF has received paid instructor fees from Pfizer. All were paid to the institution., Suzanne C. Diepstraten: None declared, Nanette L. Vincken: None declared, Pim de Jong: None declared, Timothy R. Radstake: None declared

DOI: 10.1136/annrheumdis-2019-eular.2399

SAT0533 MRI ASSESSMENT OF HIP JOINTS INVOLVEMENT IN EARLY AXIAL SPONDYLOARTHRITIS PATIENTS

Ekaterina Agafonova, Tatiana Dubinina, Daria Rumiantceva, Anastasiya Demina, Shandor Erdes. *Federal State Budgetary Scientific Institution "Research Institute of Rheumatology named after V.A.Nasonova, laboratory spondylarthritis, Moscow, Russian Federation"*

Background: Based on available data from epidemiological studies in Russia different degree of hip involvement was found in 46% of ankylosing spondylitis (AS) patients, but only 7% of all cases resulted in total hip replacement procedures.

Objectives: To compare the ultrasound (US) manifestations of hip arthritis (coxitis) with the results of magnetic resonance imaging (MRI) of the hip joints (HJ) in patients with early axial spondyloarthritis (AxSpa).

Methods: We examined 39 patients (mean age 28 ± 5.92 years) with a diagnosis of AxSpa (ASAS criteria 2009), with prescription inflammatory back pain no more than 5 years; age <45 years with US of coxitis. The median duration of the disease was 21 [3-60] months. BASDAI 4.0 ± 2.0 , BASFI $2.0 [0.4; 3.5]$. All patients underwent clinical, radiological, ultrasound, and MRI scans of HJ. For ultrasound coxitis was considered an increase in cervical-capsular distance (CCD) more than 7 mm. Coxitis is found in 27 men and 12 women. The CCD median was 7.5 [7.0; 8.2] mm.

Results: All patients in addition to ultrasound signs of coxitis had MRI signs of inflammation of the HJ. Pain in the HJ was determined in 33 (85%) patients. The relationship between the level of pain and the amount of effusion in HJ was not detected. RCC significantly correlated with laboratory parameters of disease activity, such as ESR ($r = 0.34$) $p < 0.05$ and CRP ($r = 0.3$), $p < 0.05$. X-ray examination revealed radiological changes in 7 patients (BASRI hip II-III). The following inflammatory changes (IC) were identified in the study population using MRI: synovitis

was established in 36 (92%) patients, bone marrow edema (BME) – in 1 (3%), and combination of synovitis with BME was observed in 2 patients (5%).

Conclusion: Ultrasound allows you to detect changes in H in the early stages of the disease. It is possible to use ultrasound as a screening method to determine the presence of synovitis in the TBS. MRI HJ with AxSpa, allows to determine whether the patient has inflammatory changes, including in the absence of radiological changes in these joints. In 15% of cases, coxitis is asymptomatic at early stages, which requires a thorough examination of patients with AxSpa. In addition to clinical and radiological examination, all patients were subjected to MRI with T1 and STIR modes.

Disclosure of Interests: Ekaterina Agafonova: None declared, Tatiana Dubinina: None declared, Daria Rumiantceva: None declared, Anastasiya Demina: None declared, Shandor Erdes Consultant for: Development of studies concepts., Speakers bureau: Educational meetings organized or supported by companies.

DOI: 10.1136/annrheumdis-2019-eular.3522

SAT0534 MAGNETIC RESONANCE IMAGING MARKERS IMPROVE THE PREDICTION MODEL FOR TOTAL KNEE REPLACEMENT OVER 13 YEARS IN OLDER ADULTS

Benny Antony¹, Ishanka Munugoda¹, Dawn Aitken¹, Petr Otahal¹, Michelle Lorimer², Stephen Graves³, Johanne Martel-Pelletier⁴, Jean-Pierre Pelletier⁴, Flavia Cicuttini⁵, Graeme Jones¹. ¹University of Tasmania, Menzies Institute for Medical Research, Hobart, Australia; ²Australian Orthopaedic Association National Joint Replacement Registry (AOANJRR), Adelaide, Australia; ³South Australian Health and Medical Research Institute (SAHMRI), Adelaide, Australia; ⁴Osteoarthritis Research Unit, University of Montreal Hospital Research Centre, Montreal, Canada; ⁵Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Australia

Background: Quantitative and semi-quantitative measurement of structural changes on MRI have been widely adopted in knee osteoarthritis (OA) research. However, few long-term studies describe the independent association of these structural factors and total knee replacement (TKR) in older adults.

Objectives: To describe whether cartilage defects, bone marrow lesions (BMLs), effusion-synovitis, and meniscal pathologies at baseline are associated with TKR over 13 years and to estimate the additive effect of these measures for risk prediction of TKR.

Methods: 1082 participants (mean age 62.8 years, 50% female) were randomly recruited from Tasmania and followed over 13.3 years. A 1.5T MRI scan of the right knee was acquired at baseline (n=930). Cartilage defects (grade 0-4), BMLs (grade 0-3), effusion-synovitis (grade 0-3), meniscal tears (grade 0-3) and meniscal extrusion (grade 0-2) was scored at baseline using T1-weighted and T2-weighted MRI. WOMAC knee pain was recorded from questionnaires and radiographic OA (ROA) was assessed by OARS scale. The incidence of primary (first-time) TKR was determined by data linkage to the Australian Orthopaedic Association National Joint Replacement Registry (AOANJRR). Log binomial regression were used to estimate the risk of TKR associated with baseline MRI measures, adjusting for age, sex, BMI, ROA, and pain. Receiver operating characteristic (ROC) analyses were used and area under the curves (AUCs) were compared.

Results: After adjustment for age, sex, BMI, and ROA status, baseline cartilage defects (grade 2, RR=6.71; grade 3, RR=11.17; grade 4, RR=13.2; all $p < 0.01$) were consistently and significantly associated with TKR over 13 years. These associations remained significant after further adjustment for other MRI pathologies. BMLs (grade 1, RR=2.74; grade 2, RR=2.64; grade 3 RR=4.01, all $p < 0.05$ except grade 2), grade 3 effusion-synovitis (RR=2.74, $p < 0.05$) and suprapatellar effusion-synovitis area were associated with TKR over 13 years. However, these associations of BMLs and effusion-synovitis with TKR were largely dependent on cartilage defects. Those who had TKR all had a meniscal tear at baseline, with 96% of them having a grade 3 tear.

Compared to the baseline model with age, sex and BMI (area under the ROC curve [AUC = 0.62]), Model 1 with the addition of ROA and WOMAC pain performed better (AUC =0.77). Addition of cartilage defects to Model 1 resulted in a significant increase of AUC to 0.84. However, addition of BMLs and effusion-synovitis to the model 1 resulted in a non-significant small increase in AUC (0.78 and 0.76) only. The combination of all MRI pathologies to Model 1 resulted in a significant increase of AUC to 0.82.

Conclusion: Baseline knee MRI structural pathology markers can predict TKR over the long-term, suggesting that MRI structural markers are good predictors of rapid knee OA progression in the general population.