AB0088

MECHANICAL STIMULUS INDUCED BY CHIROPRACTIC MANIPULATION REDUCES CARTILAGE, SUBCHONDRAL BONE DAMAGE AND SYNOVIAL INFLAMMATION IN AN EXPERIMENTAL MODEL OF OSTEOARTHRITIS

Francisco Miguel Conesa-Buendía^{1,2}, Aránzazu Mediero-Muñoz², Ricardo Fujikawa¹, Paula Gratal Viñuales², Francisca Mulero³, Juan Pablo Medina Giménez², Gabriel Herrero-Beaumont^{2,4}, Raquel Largo-Carazo^{2,4}, Arantxa Ortega DE Mues¹. ¹Madrid College of Chiropractic/RCU Maria Cristina El Escorial, Madrid, Spain; ²IIS-Fundación Jiménez Díaz, Rheumatology, Madrid, Spain; ³Centro Nacional De Investigaciones Oncológicas (CNIO), Madrid, Spain; ⁴Universidad Autónoma De Madrid, Madrid, Spain

Background: Osteoarthritis (OA) is a degenerative joint disease characterized by cartilage degradation, although both subchondral bone deterioration and synovial inflammation are also hallmarks of the disease. Chiropractic manipulation (CM) is a therapeutic approach focused on the diagnosis, treatment and prevention of musculoskeletal disorders. It is essentially manual, allowing the chiropractor to restore the normal range of motion and function of the joints, muscles, and ligaments. Clinical evidences suggest that CM might exert positive effects in OA patients.

Objectives: The aim of this study was to evaluate the effects of CM on cartilage, subchondral bone and synovitis state in an OA rabbit model.

Methods: Ten (4 months old) male New Zealand rabbits underwent knee surgery to induce OA by transection of anterior cruciate ligament. One week after the surgery, CM was performed using the chiropractic adjusting instrument ActivatorV as follows: Force 2 setting was applied onto the tibial tubercle of the right hind limb (true manipulation, TM-OA group), at an angle of approximately 90°, from medial to lateral, whereas the corresponding left hind limb received a false manipulation (FM-OA group) consisting of ActivatorV firing in the air and slightly touching the tibial tubercle. These procedures were repeated 3 times a week for 8 weeks. Three healthy animals were used as control. Following sacrifice, µCT and histological damage evaluation (Mankin score) were done in femur and tibiae. RANKL/OPG protein expressions were studied by immunohistochemistry in tibia samples. Sinovitis was assessed by Krenn score and immunohistochemistry for macrophages (RAM11) and angiogenesis (CD31) were done. Protein expression of VEGF, COX2, TNFα, IL-1b and MMP3 were determined by Western Blot.

Results: In the OA rabbits, subchondral BMD decreased in relation to control, been partially reversed in the tibiae of TM-OA group. When subchondral trabecular bone structural parameters were analyzed by microCT, a significant decrease of bone volume/trabecular volume (BV/TV), trabecular number (Tb.N) and trabecular thickness (Tb.Th) was observed in the OA rabbits, while trabecular separation (Tb.S) increased compared to control animals. TM-OA group showed a significant improvement of these parameters compare to FM-OA group. FM-OA joints had higher Mankin score (cartilage damage) than control joints, and TM decreased Mankin score compare to FM-OA ($p < 0.05$). The study of RANKL and OPG in cartilage and subchondral bone showed that the significant increase in the RANKL/OPG ratio observed in both tissues respect to controls, was partially reversed in TM-OA group. OA synovial membranes presented a total Krenn synovitis score higher than healthy animals; however, TM-OA rabbits exhibited scores lower than FM-OA ($p < 0.05$). RAM11 immunostaining revealed lower expression in TM-OA synovial membranes compared to FM-OA ($p < 0.05$). Finally, the significant increase in proinflammatory cytokines (COX2, TNFα and IL-1b) and MMP3 were observed in the synovial membranes of OA rabbits respect to controls, being partially reversed in TM-OA group. Likewise, VEGF and CD31 expression was higher in FM-OA synovium compared to TM-OA.

Conclusion: These results suggest that mechanical stimulus induced by CM may retard the pathologic progression of OA. The beneficial effects of CM might be associated with an improvement in bone and cartilage damage and also inflammatory state.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.1193

AB0089

IN VITRO EFFECT OF FASCIOLA HEPATICA EXTRACT ON SYNOVIAL FIBROBLAST OF RHEUMATOID ARTHRITIS PATIENTS

Suelen Dalmolin^{1,2}, Renata Ternus Pedo^{1,2}, Mirian Farinon^{1,2}, Jordana Miranda de Souza Silva^{1,2}, Vanessa Hax³, Martin Cancela⁴, Henrique Bunselmeyer Ferreira⁴, Rafaela Cavalheiro Do Espírito Santo^{1,2}, Fabiany Gonçalves⁵, Ricardo Xavier^{2,3} on behalf of Laboratório de Doenças Autoimunes. ¹Laboratório de Doenças Autoimunes, Porto Alegre, Brazil; ²Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil; ³Hospital de Clínicas de Porto Alegre, Serviço de Reumatologia, Porto Alegre, Brazil; ⁴Universidade Federal do Rio Grande do Sul, Departamento de Biologia Molecular e Biotecnologia, Porto Alegre, Brazil; ⁵Erasmus University Medical Center, Department of Internal Medicine, Rotterdam, Netherlands

Background: Synovial fibroblasts (FLS) have an aggressive and invasive profile and play a major role in RA[1]. Available therapeutic options are effective for

induction and maintenance of disease remission[2], but not all patients respond to treatment and there is no cure. So new therapeutic options need to be explored. Studies have shown that products secreted by helminths contain components with anti-inflammatory properties[3-4], capable of suppressing Th1 immune response[5] and the production of inflammatory cytokines[6]. One example is the immunomodulating properties[7] of the extract of *Fasciola hepatica*.

Objectives: To evaluate the effects of *F. hepatica* extract in FLS from RA patients.

Methods: Firstly, the cultures of FLS collected from the synovial fluid of RA patients were exposed at different concentration of *F. hepatica* extract (60µg/mL, 80µg/mL and 100µg/mL) and analyzed after 24h, 48h and 72h by MTT cell proliferation assay. FLS controls were exposed to standard culture medium. The effect of extract was also evaluated through adhesion and wound healing assay, and TNF-α production after IFNγ stimulation. Statistical analysis was performed with ANOVA or T Test and the $p < 0.05$ were considered statistically significant.

Results: *F. hepatica* extract decreased the cell proliferation of FLS at concentration of 100µg/ml after 48h (83.8% ± 5.0 extract vs 100.0% ± 0.0 control; $p < 0.05$), and at concentrations of 80 µg/ml (88.4% ± 3.0 extract vs 100.0% ± 0.0 control; $p < 0.05$) and 100 µg/ml (89.8% extract vs 3.8 extract vs 100.0% ± 0.0 control; $p < 0.05$) after 72h, when compared with control group. Based on these results, the concentration of 100µg/ml and the time of 48h were chosen for the following tests. The treatment with extract showed a decreased of FLS adhesion (92.0 cells ± 5.8 extract vs 116.3 cells ± 7.9 control; $p < 0.05$) and migratory potential (40.2 inch/h ± 13.9 extract vs 58.7 inch/h ± 6.5 control). Moreover, after treatment there was a trend of decreased TNF levels (7.0 pg/mL ± 0.5 extract vs 8.1 pg/mL ± 0.7 control).

Conclusion: Taken together, our results suggest that extract of *F. hepatica* is able to reduce the aggressive and invasive profile of FLS. However, further analyses are needed for a better understanding the mechanisms of the effect of *F. hepatica* extract on FLS and its different components, such as cystatin.

REFERENCES

- [1] Bottini N, et al. Nat Rev Rheumatol. 2013;9(1):24–33.
- [2] Tanaka Y. Korean J Intern Med. 2016;31(2): 210–218.
- [3] Hamilton C, et al. Infect. Immun. 2009;(77):2488–98.
- [4] Wang S, et al. Parasites & Vectors. 2016;(9):6.
- [5] Donnelly S. Infection and immunity. 2005;(73):166–73.
- [6] Ravidà A, et al. Molecular & cellular proteomics: MCP. 2016;15(3139–3153).
- [7] Cancela M, et al. Parasitology. 2017;144(13):1695–707.

Acknowledgement: FIPE/HCPA, Serviço de Reumatologia/HCPA

Disclosure of Interests: Suelen Dalmolin: None declared, Renata Ternus Pedo: None declared, Mirian Farinon: None declared, Jordana Miranda de Souza Silva: None declared, Vanessa Hax: None declared, Martin Cancela: None declared, Henrique Bunselmeyer Ferreira: None declared, Rafaela Cavalheiro do Espírito Santo: None declared, Fabiany Gonçalves: None declared, Ricardo Xavier Consultant for: Abbvie, Pfizer, Novartis, Janssen, Lilly, Roche

DOI: 10.1136/annrheumdis-2019-eular.5907

AB0090

INHERITED DEFICIT OF PROTEOGLYCAN MIMICKING SEPTIC ARTHRITIS

Angelo Florio¹, Riccardo Papa¹, Roberta Caorsi¹, Alessandro Consolaro¹, Roberto Gastaldi², Angelo Ravelli², Marco Gattorno¹, Paolo Picco¹, Tuula Rinne³. ¹IRCCS Istituto Giannina Gaslini, Clinica Pediatrica e Reumatologia, Genova, Italy; ²IRCCS Istituto Giannina Gaslini, Clinica Pediatrica ed Endocrinologia, Genova, Italy; ³Radboud university medical center, Genome Diagnostics Nijmegen – Department of Human Genetics, Nijmegen, Netherlands

Background: Aggrecanopathies (AP) are a heterogeneous group of skeletal disorders caused by *ACAN* gene mutations leading to dysfunction of a proteoglycan called aggrecan that plays a pivotal role in the organization of the extracellular matrix of the growth-plate cartilage. Clinically, patients with AP display bone dysplasias ranging from severe spondyloepimetaphyseal dysplasia to familial cases of osteochondritis dissecans (OCD) associated with short stature and typical facial dysmorphisms¹.

Objectives: Herein we report a patient with AP referred to our Institute for an inflammatory articular involvement mimicking a septic arthritis.

Methods: Molecular analysis of the *ACAN* gene was performed using Sanger sequencing method.

Results: A 14-year-old boy displayed pain and swelling at the right elbow. Echo-scan revealed an effusion in both coronoid and olecranon recess. Acute phase reactants were negative. A month later, since non-steroidal anti-inflammatory drugs were administered without benefit, the patient was admitted in our Institute. On physical examination, acute arthritis at the right elbow was noted: it appeared painful, warm, and not erythematous. Laboratory test showed slight elevation of acute phase reactant (C reactive protein 1.7 mg/dl, normal value <0.5).