AB0096

ANALYSIS OF THE GENE EXPRESSION OF URAT-1, ABCG2, GLUT-9, OCT3, IL1B, TLR4 AND ALPK1 IN MONONUCLEAR CELLS OF PATIENTS WITH INTER-CRITICAL GOUT

Natsuko Paniagua Díaz¹, Laura Sanchez Chapul¹, Carlos Aguilar Salinas², Alicia Ortega Mora¹, Ambar López Macay¹. ¹Instituto Nacional de Rehabilitación, Neurociencias, Mexico City, Mexico; ²Instituto Nacional de Ciencias Médicas y Nutrición Salvador Subirán, Endocrinología y Metabolismo, Mexico City, Mexico

Background: Gout is a common type of arthritis that has increased in the world due to environmental and genetic factors that associate it with an altered metabolic state. In Mexico, overweight, obesity, diabetes and metabolic syndrome have increased in children and adult population, These alterations are associated with the development of gout in adults.

The gout is characterized by an inflammatory chronic process caused by the deposition of monosodium urate crystals (CUMs) in the joints. The recognition of these crystals by macrophages and neutrophils in the joint activates innate immunity. However, high levels of uric acid (UA) in intercritical phase could activate the expression of urate transporters and inflammation genes in other cell types such as monocytes and peripheral blood lymphocytes.

Objectives: The aim of this project was to analyze the changes in gene expression of SLC22A12, SLC2A9, SLC22A3, ABCG2, ALPK1, IL1 β , TLR4 in peripheral blood mononuclear cells (PBMC) of Mexican patients with inter-critical gouty arthritis and controls without hyperuricemia (HU) comparing it with their metabolic profile.

Methods: The determination of the biochemical parameters was performed by a serum - blood chemical test for uric acid, glucose, triglycerides, cholesterol, HDL, LDL and creatinine. through the use of UniCel Dx C 600, Synchron Clinical System.

The total RNA of the PBMC cells was amplified by qRT-PCR using the GoTaq® 1-Step RT-qPCR System real kit (Promega) and the Qiagen Rotor Gene Q kit, using specific primers. The gene expression analysis was performed by the Δ CT and $\Delta\Delta$ CT method and using GAPDH as a reference gene to compare the expression of urate transporter genes and inflammatory genes in the study groups.

The expression results were associated to clinical and biochemical parameters such as levels of uric acid, triglycerides, and weight. All statistical analyzes were performed with the SPSS program (IBM 22 version for Windows, SPSS, Inc., Chicago, IL, EU), using student's t-test or U Mann-Whitney test, for the normal and non-normal data, when $p < 0.05$ or $p < 0.01$.

Results: In this study we compared 24 patients with inter-critical gout and 26 controls without HU. It was found that the expression of ABCG2 was higher in the patients and that the expression of OCT3 was higher in overweight people, regardless of whether they had gout or not compared with normal weight people. The expression of ALPK1 was higher among patients and controls while the expression of TLR4 and IL1 β was not different. The patients had a higher mean in the levels of UA, triglycerides, weight, age and hypertension, but lower HDL than the controls.

Conclusion: Patients with inter-critical gout maintain metabolic alterations associated with the metabolic syndrome although some of them are under treatment. They are also characterized by having overexpression of ABCG2 and ALPK1. These conditions could favor a new attack of gout.

REFERENCES

- [1] Dalbeth, N.; Merriman, T. R.; Stamp, L. K., Gout. *Lancet* 2016, 388 (10055), 2039-2052.
- [2] Merriman, T. R., An update on the genetic architecture of hyperuricemia and gout. *Arthritis research & therapy* 2015, 17, 98
- [3] Ko, A. M.; Tu, H. P.; Liu, T. T.; Chang, J. G.; Yuo, C. Y.; Chiang, S. L.; Chang, S. J.; Liu, Y. F.; Ko, A. M.; Lee, C. H.; Lee, C. P.; Chang, C. M.; Tsai, S. F.; Ko, Y. C., ALPK1 genetic regulation and risk in relation to gout. *International journal of epidemiology* 2013, 42 (2), 466-74.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3901

AB0097

NOVEL EX VIVO MODEL OF SEPTIC ARTHRITIS DEVELOPED TO IDENTIFY BIOMARKERS RELEASED UPON ARTICULAR INFECTION

Kathryn McCall, Caroline Atherton, Neal L. Millar, Carl Goodyear, Tom Evans, Iain McInnes. *University of Glasgow, Institute of Infection, Immunity, and Inflammation, Glasgow, United Kingdom*

Background: Septic arthritis (SA) is most commonly caused by infectious agents such as *Staphylococcus aureus*. Due to the morbidity and mortality associated with the disease, joint infections are treated as a medical emergency and

therefore early diagnosis is essential. Unfortunately, no reliable biomarkers are available to assist diagnosis, beyond serum CRP and synovial fluid culture.

Objectives: To generate a novel *ex vivo* model of septic arthritis comprised of cartilage matrix, neutrophils and SA-associated *Staph. aureus*, to identify biomarkers released upon an inflammatory host response to infection.

Methods: Cartilage explants biopsied from human femoral heads were compared to chondrocyte disks differentiated from mesenchymal stem cells (MSC). MSC differentiation into chondrocytes under defined media and growth factor conditions was confirmed by IHC staining for collagen II on the disks. Explants and disks were infected for 48h with 10⁶ cfu *Staph. aureus* (SA patient derived). Structural damage was measured by glycosaminoglycan (GAG) release. Neutrophils were purified from health donor blood samples using the StemCell whole blood isolation kit, and 10⁶ cell were added in the final 4h of bacterial infection. Disks were stained with eosin to visualise microscopic damage, CMFDA for viability and PI to identify cell death.

Results: Co-culture of MSC-derived chondrocyte disks with neutrophils or *Staph. aureus* alone resulted in microscopically identifiable structural damage to the disk matrix. The addition of neutrophils in the final 4h of the *Staph. aureus* co-cultures resulted in enhanced structural damage. Assessment of cellular viability revealed that both neutrophils and *Staph. aureus* alone induced cell death throughout the disk whilst the combination of both resulted in total abrogation of viable chondrocytes. Subsequent evaluation of GAG release, however, showed no clear differentiation between the conditions and as such this model was judged not to be sufficient for biomarker development. In comparison, femoral head cartilage explants studies demonstrated that *Staph. aureus* infection in this explant model resulted in significant GAG release after 24h and 48h, which was consistent between donors.

Conclusion: Cartilage explants provide a superior and more physiologically relevant cartilage matrix model for the *ex vivo* analysis of septic arthritis. *Staph. aureus* infection resulted in reproducible damage to cartilage matrices that was not emulated in an MSC-derived chondrocyte disk system. These findings suggest that the femoral head cartilage explants are the optimal matrix model to identify biomarkers for joint infections.

Disclosure of Interests: Kathryn McCall: None declared, Caroline Atherton: None declared, Neal L. Millar: None declared, Carl Goodyear Grant/research support from: AstraZeneca, BMS, Celgene, Janssen, MedAnnex, Pfizer and UCB, Speakers bureau: Abbvie, Tom Evans: None declared, Iain McInnes Grant/research support from: AstraZeneca, Celgene, Compugen, Novartis, Roche, UCB Pharma, Consultant for: AbbVie, Celgene, Galvani, Lilly, Novartis, Pfizer, UCB Pharma

DOI: 10.1136/annrheumdis-2019-eular.7501

AB0098

GENE EXPRESSION AND FUNCTIONAL COMPARISON BETWEEN MESENCHYMAL STEM CELLS FROM LATERAL AND MEDIAL CONDYLES OF KNEE OSTEOARTHRITIS PATIENTS

Clara Sanjurjo Rodriguez^{1,2}, Thomas Baboolal^{2,3,4}, Agata Burska², Frederique Ponche², Jehan El-Jawhari², Joseph Aderinto⁴, Owen Wall⁴, Hemant Pandit^{2,3,4}, Dennis McGonagle^{2,3,4}, Elena Jones². ¹Universidad da Coruña; ²INIBIC-CICA, CIBER-BBN, A Coruña, Spain; ³Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, United Kingdom; ⁴NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds, United Kingdom; ⁵Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom

Background: Osteoarthritis (OA) is the most common degenerative joint disorder, mainly affecting the weight-bearing joints, and is the leading cause of physical disability worldwide. Despite the identified risk factors, the exact pathogenesis of osteoarthritis remains unclear (1). Formation of mesenchymal stem cells' (MSC) clusters and their aberrant osteogenic differentiation in OA subchondral bone (SB) has been proposed as a key contributor to progression of OA in animal models (2) and in human hip OA (3). MSCs have a crucial role in joint repair but it remains unknown how OA severity affects MSC numbers or characteristics. Knee OA provides a good model to determine this as in knee OA, distribution of damage is usually asymmetrical and tends to be more severe in the medial (Med) compared to the lateral (Lat) compartment (4).

Objectives: The aim of this study was to determine whether there were differences in MSC numbers, topography and gene expression (GE) between Med and Lat femoral condyles of patients with knee OA.

Methods: Condyles were obtained from OA patients that underwent total knee replacement (n=16; UK Ethics Committee approval, 14/YH/0087). Decalcified samples were histologically evaluated for cartilage damage, bone sclerosis and CD271+ MSC distribution (2). MSCs were extracted from SB and sorted using the CD271+CD45- phenotype for GE analysis. Colony forming unit (CFU-F) (3), tri-lineage differentiation (5) and population doubling (PD) assays were performed. MSC numbers, topography and GE were compared between condyles.

Results: Med condyles presented significantly higher ($p < 0.05$) degree of cartilage damage (median OARSI score 20, range 15-20) compared to Lat condyles

(score 3, range 1.5-15), and sclerotic SB area was higher in the Med condyles (64.3 ± 7.0 vs $27.6 \pm 7.4\%$, $p < 0.05$). CD271+ MSCs were found in bone-lining and perivascular locations, and accumulated in the areas of vascular invasion into the articular cartilage in both, Med and Lat condyles. No differences in the MSC numbers were found using CFU-F assay (1.4 ± 0.7 and $1.6 \pm 0.8\%$ of total cells in Lat and Med condyles, respectively), or by flow cytometry (4.4 ± 3.9 and $5.9 \pm 6.8\%$, respectively). MSCs had similar PD and trilineage capacities *in vitro*. Out of 95 genes tested, 3 genes were significantly upregulated in Med condyle CD271+ MSCs: GREM1 (lateral MSCs below detection), PTHLH (2.4-fold, $p = 0.02$) and STMN2 (10.5-fold, $p = 0.02$), all implicated in osteogenic differentiation and mineralisation (6, 7, 8).

Conclusion: Upregulation of ossification-related genes in Med condyle MSCs suggest their potential contribution to sclerotic plate formation and OA severity. Further work is needed to establish if biomechanical or biological stimulation of these MSCs can result in GE modulation in preference of cartilage tissue formation and subchondral bone restoration.

REFERENCES

- [1] Conaghan. Nat Rev Rheumatol (2013).
- [2] Zhen. Nat Med (2013);
- [3] Campbell. Arthritis Rheumatol (2016);
- [4] Bae. Cartilage (2010).
- [5] Altaie. Cytotherapy (2018);
- [6] Hu. Mol Med Rep (2017);
- [7] García-Martín. Mol Endocrinol (2014);
- [8] Chiellini. Biochem Biophys Res Commun (2008)

Acknowledgement: CSR funded by fellowship Xunta de Galicia (Consell de Cul, Educ e Ordenación Univ). TB and DM part-supported by NIHR Leeds Musculoskeletal Biomed Research Centre. TB part-supported by FOREUM. JE part-funded by AO foundation start-up grant.

Disclosure of Interests: Clara Sanjurjo Rodriguez: None declared, Thomas Baboolal: None declared, Agata Burska: None declared, Frederique Ponchel: None declared, Jehan El-Jawhari: None declared, Joseph Aderinto: None declared, Owen Wall: None declared, Hemant Pandit Grant/research support from: GSK, Consultant for: For education for Bristol Myers Squibb, Dennis McGonagle Consultant for: Lilly, Novartis UCB, Speakers bureau: Lilly, Novartis UCB, Elena Jones: None declared

DOI: 10.1136/annrheumdis-2019-eular.2649

AB0099

COCULTURE OF SYNOVIAL MEMBRANE AND CARTILAGE: A POTENTIAL HUMAN EX VIVO CARTILAGE DEGRADATION MODEL

Johanna Mucke¹, Victoria Warzynski¹, Ellen Bleck¹, Thomas Pauly², Tim Claßen², Matthias Schneider¹, Stefan Vordenbäumen¹. ¹Poliklinik für Rheumatologie & Hiller Research Centre for Rheumatology, Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany; ²Dept. of Orthopedic Surgery, St. Elisabeth-Hospital, Rheinisches Rheumazentrum, Meerbusch-Lank, Germany

Background: Cartilage degradation is a hallmark in both osteoarthritis (OA) and rheumatoid arthritis (RA). Ex vivo cartilage models offer the potential to study disease processes but are surprisingly rarely used.

Objectives: Here we investigate the effect of synovial tissue and stimulation with TNF- α on cartilage degradation in a physiologically relevant coculture-model of synovial membrane and cartilage.

Methods: Cartilage and synovial membrane were collected from patients with RA and OA who underwent joint replacement. Cartilage was cultivated either alone or with synovial membrane in medium with or without TNF α as a stimulant for up to 28 days. The explants were harvested at different time points and Safranin-O-staining was performed to visualize proteoglycan (PG) content. A scoring system from 0 (no PG content) to 3 (high PG content) was applied as a semi-quantitative measure for evaluation. Linear mixed modeling adjusted for gender and age was performed to identify factors influencing PG content.

Results: Samples of 18 patients (9 OA, 9 RA) were collected. Factors associated with proteoglycan loss in cartilage were the time in culture ($p = 0.02$), stimulation with TNF- α ($p = 0.0007$) and coculture with synovial membrane ($p = 0.00002$) in univariate analysis with TNF- α ($p = 0.0009$) and coculture ($p = 0.001$) maintaining the level of significance in multivariate analysis. (Figure 1)

Conclusion: Exposure to synovial tissue and TNF stimulation drive cartilage proteoglycan loss in an ex vivo cartilage degradation model. This model may prospectively serve to investigate pathophysiologic pathways of cartilage degradation in both OA and RA.

REFERENCES

None

Figure 1. Proteoglycan content at d38. PG, proteoglycan.

Disclosure of Interests: Johanna Mucke: None declared, Victoria Warzynski: None declared, Ellen Bleck: None declared, Thomas Pauly: None declared, Tim Claßen: None declared, Matthias Schneider Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study., Speakers bureau: Chugai, Stefan Vordenbäumen: None declared

DOI: 10.1136/annrheumdis-2019-eular.2159

AB0100

COMPARATIVE ANALYSIS OF THE EFFECTIVENESS AND TOLERANCE OF INTRA-ARTICULAR INJECTION OF PLATELET RICH PLASMA VS HYALURONIC ACID AND NON-ARTHROSCOPIC JOINT LAVAGE IN PATIENTS WITH KNEE OSTEOARTHRITIS

Rafaela Ortega Castro¹, Rocio Segura¹, Jerusalem Calvo Gutierrez¹, Rocio Espino-Garcia², Maria del Carmen Castro Villegas¹, Alejandro Escudero Contreras¹, Miguel A. Caracuel-Ruiz¹, Ladehesa Pineda Lourdes¹, Desiree Ruiz-Vilchez¹, Montserrat Romero-Gómez¹, Pilar Carreto-Font³, Eduardo Collantes Estevez¹, Font Ugalde Pilar¹. ¹Reina Sofía University Hospital/ Maimónides Institute of Biomedical Research of Córdoba (IMIBIC); ²Clinical Management Unit of Rheumatology, Córdoba, Spain; ³University of Cordoba/Reina Sofía Hospital, Córdoba, Spain; ³Tudela Duero Clinic/East Area, Valladolid, Spain

Background: Knee Osteoarthritis (KOA) is a common condition associated with pain and morbidity. The impact of this pathology makes it necessary to develop new procedures for joint regeneration. Regenerative treatments such as platelet-rich plasma (PRP) are being proposed as a safe alternative, capable of regenerating damaged tissues and improving the quality of life, reducing the need to surgical procedures (1-4).

Objectives: To compare the effectiveness of PRP treatment vs Hyaluronic Acid (HA) and Non-Arthroscopic Joint Lavage (NAJL) in patients with KOA.

Methods: It was performed a prospective, 3-months observational study of 51 patients with KOA, from the OA clinic of the Reina Sofía Hospital of Córdoba. Patients were treated according to clinical practice in 1 of the 3 treatment arms (PRP, HA, NAJL). Pain and disability were evaluated using Visual Analogue Scale (VAS) of joint pain, WOMAC and OARSI. We compared the baseline data using simple ANOVA and chi-squared tests and the intra- and inter-group differences before and after treatment (0, 1 and 3 months) using the mixed ANOVA test. All comparisons were bilateral considering $p \leq 0.05$ as a significant result.

Results: 51 patients were enrolled, of which 52.9% were female. The average age ($\pm$ SD) was 58.47 ± 7.48 years. The average Body Mass Index (BMI) was 30.64 ± 4.25 Kg/m². Characteristics at baseline were comparable among the 3 treatment groups (Table 1). An overall improvement was observed during the follow-up in the PRP group in terms of pain and disability with respect to the baseline, with a VAS MD of -3.00 [95% CI, -4.79 to -1.21] in the first month and -3.08 [CI 95%, -4.79 to -1.38] at 3 months ($p < 0.001$), a MD WOMAC of -18.83 [95% CI, -30.62 to -7.05] ($p = 0.001$) in the first month, and a MD of OARSI of -19.32 [95% CI, -34.26 to -4.37], ($p = 0.008$) in the first month and -18.94 [95% CI, -36.81 to -1.07] at 3 months ($p = 0.035$). In addition, the PRP group was superior to the HA group, showing a significant improvement in the VAS at 1 and 3 months ($p < 0.005$) and compared to the NAJL group, PRP showed better results in the WOMAC throughout the follow-up ($p < 0.01$) and in the OARSI at 3 months ($p = 0.013$).

Conclusion: Based on our results, the intra-articular infiltration with PRP could therefore constitute a safe and effective treatment option in KOA patients, with short-medium term results superior in terms of pain and disability to conventional treatment with HA and NAJL.