AB0143 PERIODONTITIS AND SALIVA ANTIBODIES TO CITRULLINATED PEPTIDES IN RHEUMATOID ARTHRITIS

Anna Svård¹, Stefan Renvert², Johan Berglund³, Maria Söderlin⁴. ¹Center for clinical research Dalarna, Uppsala University, Falun, Sweden; ²University of Kristianstad, Kristianstad, Sweden; ³Blekinge Institute of Technology, Karlskrona, Sweden; ⁴Department of Clinical Sciences, Section of Rheumatology, Lund University, Lund, Sweden

Background: Mucosal immunity, involving antibodies against citrullinated peptides (ACPA), is a prevailing hypothesis concerning the pathogenesis of rheumatoid arthritis (RA). The oral mucosa is a possible site that is involved. Several epidemiological studies have found an association between RA and periodontitis, and periodontal inflammation has been suggested as a possible pathogenic factor.

In established RA, periodontitis has been associated with ACPA positivity (1) although a recent large study (2) did not find any association between periodontitis and IgG ACPA. Previous studies only evaluate ACPA in serum, and as local mucosal immune responses are likely involved in periodontitis, saliva antibodies are of interest to investigate. Saliva IgA ACPA has in a pilot study of RA patients been associated with lower disease activity and less erosions(3), but has not previously been evaluated in relation to periodontitis.

Objectives: To evaluate if saliva or serum ACPA is associated to periodontitis, in patients with RA.

Methods: In this population-based cohort of patients with RA (n=132), we included all patients with RA older than 61 years, living in Karlskrona in southern Sweden, between October 21, 2013 and January 7, 2015.

The patients underwent a clinical examination by a rheumatologist, a dental clinical examination by a dental nurse and a panorama radiograph of the jaws and teeth. Periodontitis was defined as a distance from the cement-enamel junction (CEJ) to the bone level ≥ 5 mm, as defined by the radiographic assessment, at $\geq 30\%$ of interproximal sites.

We analysed antibodies against second generation cyclic citrullinated peptides (anti-CCP) in enzyme immunoassays in serum and saliva. Saliva samples were analysed in parallel for anti-CCP and anti-CAP, the arginine containing control peptide. Serum ACPA positivity was defined as the 99th percentile of healthy controls. In saliva, a positive test was defined as a difference in optical density (OD) for IgA anti-CCP and IgA anti-CAP (delta OD) > 0 .

Results: Saliva IgA ACPA was found in 11% of RA patients with periodontitis and in 24% of RA patients with no periodontitis (p=0.052). There was no difference between the groups concerning IgG or IgA ACPA in serum.

Table 1. Demographic and laboratory characteristics of the study participants

	Patients with RA and periodontitis N=80	Patients with RA no periodontitis N=52	P value*
	Mean or percent	Mean or percent	
Age at inclusion, years	71.4	69.0	0.034
Females%	71%	69%	0.847
Disease duration, years	12.1	12.3	0.930
DAS28 ESR	3.01	2.84	0.403
DAS28 CRP	2.81	2.80	0.936
Erosions in hands or feet %	59%	53%	0.584
Biologics currently%	21%	23%	0.806
Biologics ever%	36%	44%	0.577
Smoker (ever vs never)%	64%	60%	0.714
Serum IgG ACPA positive,%	66%	69%	0.849
Serum IgA ACPA positive,%	35%	43%	0.461
Saliva IgA ACPA positive,%	24%	11%	0.052

*Student's T-test was used for continuous variables and Fischer's exact test for categorical variables.

A logistic regression analysis adjusting for gender, age and smoking showed an Odds Ratio of 0.299 (95% CI=0.096-0.929, p=0.037) for RA patients with periodontitis to be positive for saliva IgA ACPA, compared to RA patients with no periodontitis.

Conclusion: A weak, negative, correlation was observed between periodontitis and saliva IgA ACPA, when adjusting for age, sex and smoking. No correlation was observed between periodontitis and IgA or IgG ACPA in serum. In this cohort of patients with established RA, our findings do not support the hypothesis that periodontitis leads to increased formation of IgA ACPA in saliva or serum.

REFERENCES

- [1], Dissick A, et al. J Periodontol. 2010;81(2):223-30.

- [2] Eriksson K, et al. PloS one. 2016;11(5):e0155956.
- [3] Svård A, et al. Immunobiology. 2013;218(2):232-7.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.1209

AB0144 EFFECTS OF AB501 (CERTOLIZUMAB MICE EQUIVALENT) IN ARTHRITIS INDUCED BONE LOSS

Bruno Vidal¹, Mikko A. Finnilä², Inês Lopes¹, Rita Cascão¹, Joao Eurico Fonseca^{1,3}. ¹Unidade de Investigação em Reumatologia, Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Centro Académico de Medicina de Lisboa, Lisboa, Portugal; ²Research Unit of Medical Imaging, Physics and Technology, Faculty of Medicine, University of Oulu, Oulu, Finland; ³Serviço de Reumatologia e Doenças Ósseas Metabólicas, Hospital de Santa Maria, CHLN, Centro Académico de Medicina de Lisboa, Lisboa, Portugal

Background: Rheumatoid arthritis (RA) is a chronic immune-mediated inflammatory disease, which causes local and systemic bone damage.

Objectives: The main goal of this work was to analyze, how treatment intervention with Ab501 (certolizumab mice equivalent) prevents the disturbances on bone structure and mechanics induced by arthritis.

Methods: Thirty one DBA/1 collagen-induced arthritis (CIA) mice were randomly housed in experimental groups, as follows: arthritic untreated (N=10), preventive intervention with ab501 2 days before arthritis induction (N=10) and treatment intervention upon arthritis onset with ab501 (N=11). A non-induced group (N=5) was used as a control. Mice were monitored during 70 days after disease induction for the inflammatory score, ankle perimeter and body weight. Non-induced mice were used as controls. After 70 days of disease progression mice were sacrificed and bone samples were collected for histology, micro-computed tomography (μCT) and 3-point bending analysis. In addition, blood samples were also collected for bone turnover markers quantification. Statistical differences were determined with Mann-Whitney tests using GraphPad Prism (GraphPad, California, USA). Data were expressed as median with interquartile range. Differences were considered statistically significant for p<0.05.

Results: Results showed that Ab501 administration was able to control and abrogate disease development both in preventive and early therapeutic intervention. μCT results revealed that ab501 was able to preserve bone structure when delivered before arthritis induction.

Conclusion: Ab501 preventive administration was able to control inflammation and prevent the degradative effects of arthritis on trabecular bone structure.

Acknowledgement: This work was supported by UCB in the context of an Investigator Initiated Study where UCB provided financial and product support.

Disclosure of Interests: Bruno Vidal Grant/research support from: Aspire Pfizer 2013, Mikko A. Finnilä: None declared, Inês Lopes: None declared, Rita Cascão: None declared, Joao Eurico Fonseca: None declared

DOI: 10.1136/annrheumdis-2019-eular.7025

AB0145 THE INHIBITION OF JAK PATHWAY WAS ASSOCIATED WITH REDUCTION OF AUTOPHAGY IN SYNOVIOCYTES FROM RHEUMATOID ARTHRITIS PATIENTS

Marta Vomero¹, Mattia Caliste¹, Cristiana Barbat¹, Tania Colasanti¹, Francesca Spinelli¹, Fulvia Ceccarelli¹, Carlo Perricone¹, Annacarla Finucci¹, Mariangela Speziali¹, Alessandra Ida Celia¹, Michele Bombardieri², Fabrizio Conti¹, Guido Valesini¹, Cristiano Alessandri¹. ¹Sapienza University of Rome, Internal Medicine and Medical Specialties, Rome, Italy; ²Queen Mary University of London, London, United Kingdom

Background: The pathway of Janus tyrosine kinases (JAKs) has a central role in the pathogenesis of rheumatoid arthritis (RA) by regulating multiple immune functions and cytokine production. The orally available JAK inhibitor CP-690,550, named tofacitinib, is able to inhibit Jak1, Jak2 and Jak3 and it showed a great clinical efficacy in RA patients not responding to methotrexate or TNF-inhibitors (1). Autophagy, a highly conserved mechanism involved in the degradation of intracellular components, was found to be dysregulated in several autoimmune diseases including RA (2). In fact, fibroblasts like synovioocytes (FLS) from RA patients showed a resistance to apoptosis associated with the induction of autophagy (3). Maeshima and colleagues recently demonstrated that tofacitinib suppressed proliferation, but not apoptosis, of CD4+ T cells derived from the synovium (4), but to date there are no data on the effect of inhibition of JAK pathway on autophagy.

Objectives: Since hyperactive autophagy has been associated with impaired apoptosis of RA FLS, the aim of the study was to investigate the role of tofacitinib in modulating autophagy and apoptosis in FLS from RA patients.

Methods: Primary RA FLS isolated from RA biopsies (Figure 1A) were cultured in presence of autophagy inducer rapamycin (as control condition) and tofacitinib (1μM). After 24h of culture, autophagy was evaluated both by flow cytometry and

western blot analysis. Apoptosis was analyzing by annexin V (AV) and propidium iodide (PI) apoptosis detection kit by flow cytometry.

Results: As expected tofacitinib inhibited the expression of the phosphorylated form of STAT3 ($p < 0.01$). Rapamycin caused an increase in autophagy, while the levels of autophagy marker LC3-II were reduced after treatment with tofacitinib in vitro ($p < 0.001$ and $p = 0.02$, respectively, Figure 1B). In addition, the analysis of autophagic vacuoles by specific fluorescence dye confirmed the reduction of autophagy in RA-FLS treated with tofacitinib. The percentage of annexin V-positive apoptotic cells was not influenced by the treatment with tofacitinib.

Conclusion: The results of this study elucidated a new mechanism of action of tofacitinib related to autophagy modulation and led to a better understanding of the role of autophagy in RA. This study was supported by an unconditioned Research grant from Pfizer Inc.

REFERENCES

- [1] Burmester GR, Blanco R, Charles-Schoeman C, et al. Tofacitinib (CP-690,550) in combination with methotrexate in patients with active rheumatoid arthritis with an inadequate response to tumour necrosis factor inhibitors: a randomised phase 3 trial. *Lancet* 2013; 381: 451-60.
- [2] Vomerio M, Barbatì C, Colasanti T, et al. Autophagy and Rheumatoid Arthritis: Current Knowledge and Future Perspectives. *Front Immunol*. 2018;9:1577.
- [3] Shin YJ, Han SH, Kim DS, et al. Autophagy induction and CHOP under-expression promotes survival of fibroblasts from rheumatoid arthritis patients under endoplasmic reticulum stress. *Arthritis Res Ther*. 2010; 12: R19.
- [4] Maeshima K, Yamaoka K, Kubo S, et al. The JAK inhibitor tofacitinib regulates synovitis through inhibition of interferon- γ and interleukin-17 production by human CD4⁺ T cells. *Arthritis Rheum*. 2012;64:1790-8.

A. Image of fibroblast-like synoviocytes obtained from synovial tissue of RA patients during total knee replacement and used for in vitro experiments. B. Western blot analysis of autophagy marker LC3-II after in vitro treatment with tofacitinib.

Disclosure of Interests: Marta Vomerio: None declared, Mattia Caliste : None declared, cristiana barbatì: None declared, Tania Colasanti: None declared, francesca spinelli: None declared, Fulvia Ceccarelli: None declared, Carlo Perricone Speakers bureau: BMS; Lilly, Celgene, Sanofi, Annacarla Finucci: None declared, Mariangela Speziali: None declared, Alessandra Ida Celia: None declared, Michele Bombardieri Grant/research support from: Celgene, Consultant for: Medimmune, fabrizio conti: None declared, Guido Valesini: None declared, cristiano alessandri: None declared

DOI: 10.1136/annrheumdis-2019-eular.5445

AB0146

IDENTIFICATION A MARKED INFLAMMATION MEDIATED CARDIAC DYSFUNCTION DURING THE DEVELOPMENT OF ARTHRITIS IN COLLAGEN INDUCED ARTHRITIS MICE

Fang Wang, Aishu Luo, Zihao Zhou, Wenfeng Tan. *the First Affiliated Hospital of Nanjing Medical University, Nanjing, China*

Background: Systemic inflammation is strong linked to the increased risk of cardiovascular diseases (CVD) in rheumatoid arthritis (RA)¹. Increased cardiovascular risk in RA patients can not be fully explained by the traditional risk factors, such as hypertension, obesity, diabetes, age and smoking^{2,3}. Much more work is needed to elucidate the mechanism of chronic inflammation bridging RA and cardiovascular disease.

Objectives: To investigate the cardiac changes during the development of arthritis in collagen induced arthritis (CIA) mice and to explore the potential role of inflammation on cardiac dysfunction in RA.

Methods: Arthritis severity was evaluated using clinical indices, micro-computed tomography and histopathology. Cardiac function was determined by transthoracic echocardiography at week 5, 7, 9 and 11 after immunization in mice. At week 7 (day 50), mice joints and hearts were removed for pathological study, and cardiomyocytes and cardiac fibroblasts were isolated using Langendorff perfusion method *Ex vivo* to measure expression of inflammatory and cardiac-related genes by real time PCR. The expression of key molecule in cardiac dysfunction (β -MHC) was also tested in H9c2 cardiomyocytes treated with sera derived from CIA mice or RA patients.

Results: At day 50 after immunization, cardiac function in CIA mice was prominently reduced as evidenced by decreased ejection fraction (EF) and fractional shortening (FS), increased left ventricular end-systolic volume (LVESV) and internal systolic diameter (LVIDs). Accordingly, enhanced inflammatory cells infiltration and fibrosis were identified in ventricular tissues pathologically, and increased inflammatory genes expression including TNF- α , IL-6, IL-17 and MMP3 detected in isolated ventricular cardiomyocytes and cardiac fibroblasts from CIA mice. Furthermore, H9c2 cells treated with sera from CIA mice or RA patients exhibited high levels of β -MHC.

Conclusion: Joint inflammation is associated with an obvious cardiac dysfunction and enhanced inflammation infiltration and inflammatory cytokines production in cardiomyocytes and cardiac fibroblasts during CIA development. Our data provide the direct evidence that inflammation contributes to development of cardiac diseases in RA patients.

REFERENCES

- [1] SOLOMON DH, GOODSON NJ, KATZ JN, et al.: Patterns of cardiovascular risk in rheumatoid arthritis. *Ann Rheum Dis*. 2006;65:1608-12.
- [2] JAFRI K, BARTELS CM, SHIN D, GELFAND JM, OGDIE A: Incidence and management of cardiovascular risk factors in psoriatic arthritis and rheumatoid arthritis: A population-based study. *Arthritis Care Res (Hoboken)*. 2017;69:51-7.
- [3] AN J, ALEMAO E, REYNOLDS K, et al.: Cardiovascular outcomes associated with lowering low-density lipoprotein cholesterol in rheumatoid arthritis and matched nonrheumatoid arthritis. *J Rheumatol*. 2016;43:1989-96.

Acknowledgement: This project was sponsored by the grants from the National Natural Science Foundation of China (No. 81671610, 81471610, 81471611, 81801625)

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4553

AB0147

RESEARCH PROGRESS OF MESENCHYMAL STEM CELL-DERIVED MICROVESICLES BY DELIVERING MICRORNA IN RHEUMATIC DISEASES

Na Zhang¹, Liyun Zhang², Gailian Zhang², Jin-Fang Gao³, MA Dan⁴, LI Juan⁵, Ke Xu¹. ¹Shanxi Dayi Hospital Affiliated to Shanxi Medical University, Taiyuan, China; ²Shanxi Dayi Hospital Affiliated to Shanxi Medical University, Taiyuan, China; ³Shanxi Dayi Hospital Affiliated to Shanxi Medical University, Taiyuan, China; ⁴Shanxi Dayi Hospital Affiliated to Shanxi Medical University, Taiyuan, China; ⁵Shanxi Dayi Hospital Affiliated to Shanxi Medical University, Taiyuan, China

Background: Recently, studies have shown that Mesenchymal Stem Cell-derived Microvesicles (MSC-MVs) can play a role in immune regulation and tissue repair by transporting miRNAs, and their effects are not weaker than that of their mother stem cells. Rheumatoid arthritis is an autoimmune disease characterized by chronic synovitis and progressive cartilage and bone destruction. Current treatments include non-steroidal anti-inflammatory drugs, glucocorticoids, DMARDs, and biological agents, but 30% of patients with rheumatism are ineffective, so there is an urgent need to find a new treatment.

Objectives: To review the research progress of MSC-MVs by delivering micro-RNA (miRNA) in rheumatic diseases.

Methods: Literatures were searched by using MSC-MVs, autoimmune diseases and miRNA as keywords.

Results: In the field of rheumatology, Chen C showed that after MSCT transplantation, mir-151-5p was transferred through exosomes to damaged bone marrow MSC of systemic sclerosis (SSc) mice, and the activation of IL4R/mTOR pathway was down-regulated to improve the osteoporosis of SSc mice^[1]. Exosomes of human synovial mesenchymal stem cells overexpressing miR-140-5p promoted cartilage regeneration and prevented OA in rat models^[2]. In the field of autoimmune diseases, Chen L et al. demonstrated that BMSC-derived exosomes containing miR-223 play a important role in liver protection by down-regulating NLRP3 (nod-like receptor family protein) and caspase-1 (caspase-1) in the experimental model of autoimmune hepatitis^[3]. Human BMSC-derived exosomes transfected with siFas and anti-mir-375 plasmids down-regulated Fas and miR-375 in human NOD scid gamma (NSG) mice, thereby protecting islet function and inhibiting immune rejection in islet transplantation^[4].

Conclusion: MiRNA targeting therapy with MSC-MVs as vector or the construction of MSC-MVs with overexpression of protective miRNA will provide new, accurate and promising treatment strategies for multiple rheumatic diseases.

REFERENCES

- [1] Chen, C., et al. Mesenchymal stem cell transplantation in tight-skin mice identifies miR-151-5p as a therapeutic target for systemic sclerosis. *Cell Res*, 2017. 27(4): p. 559-577. DOI:10.1038/cr.2017.11.