REFERENCE:

- [1] Bagnato G, De Andres I, Sorbara S, Verduci E, Corallo G, Ferrera A, et al. Pain threshold and intensity in rheumatic patients: correlations with the Hamilton Depression Rating scale. *Clinical rheumatology*. 2015;34(3):555-61.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7390

AB0232

TIGHT CONTROL MANAGEMENT OF EARLY RA ACHIEVES MORE PATIENTS IN REMISSION AT 6 MONTHS AND IMPROVES PAIN, FUNCTION AND FATIGUE AT 24 MONTHS COMPARED TO A CONTROL GROUP

Maria Aronsson^{1,2}, Annika Teleman¹, Stefan Bergman^{2,3}, Kristina Forslind⁴, Maria Andersson^{2,5}. **Background:** Even with modern antirheumatic treatment, many patients experience chronic pain and loss of function despite less disease activity. The understanding of how this chronic pain development can be prevented is sparse. There is furthermore little knowledge if a tight control management without specific treatment schemes and instructions for the physicians to follow could increase the remission frequency.

Objectives: To determine whether solely a tight control management of patients, without further intervention in the physician's work, can increase the remission frequency and improve patient reported outcome in early RA.

Methods: In all, 96 patients with early RA were consecutively included in a tight control study group (TCS). Inclusion criteria were early RA patients (<13 month duration) with no previous history of rheumatic disease. A control group, with so far 27 patients receiving standard care, has been analyzed. The tight control management included monthly visits to the physician the first 6 months, followed by visits at 9, 12 and 24 months. If not in remission at 6 months the patients' monthly visits could continue to 12 months. The study did not include any treatment guidelines for the physicians. Pain, Fatigue and the Patient's Global Assessment of the General Health (PtGA) were reported on a Visual Analogue Scale (VAS). The 28 tender and swollen joint score, DAS28, C-Reactive protein, Erythrocyte sedimentation rate and Health Assessment Questionnaire (HAQ) were assessed.

Results: There was no significant difference between the groups at baseline in gender, age, smoking, disease duration, disease activity, pain, HAQ, fatigue, RF and ACPA. There was a significant difference between groups in outcome, with 70% of TCS and 37% of control patients in remission (DAS28 <2.6), $p=0.010$, at 6 months, figure 1A. At 12 and 24 months there was no significant difference in the remission frequency, figure 1A. The TCS group improved more than the control group in Pain, Fatigue and HAQ during follow up, figure 1B-D. The median VAS pain at 24 months was 11 (IQR 2-22) in the TCS group and 22 (IQR 6-42) in the control group, $p=0.036$. The median VAS fatigue was 13 (IQR 2-38) in the TCS group and 33 (IQR 8-70) in the control group $p=0.028$ at 24 months. The difference in HAQ reached significant numbers solely at 6 months where the median was 0.13 (IQR 0-0.50) in the TCS group and 0.57 in the control group (IQR 0.16-0.94).

Figure 1. Disease activity (DAS28), functional disability (HAQ), pain and fatigue during 24 months follow-up. The TCS group is included in a tight control study with follow-ups every month during the first half year. The control group has received standard care. A. Median DAS28 in the separate groups over time. B. Median HAQ in the separate groups over time. C. Median VAS pain in the separate groups over time. D. Median VAS fatigue in the separate groups over time. * significant difference $p < 0.05$

Figure 1

Conclusion: Patients in tight control reached remission faster than patients receiving standard care. They also reported less pain and fatigue, despite the fact that there was no difference in remission frequency at 24 months. The results indicate that a tight control scheme could influence pain development in patients with early RA, but further studies are required.

Disclosure of Interests: Maria Aronsson: None declared, Annika Teleman Consultant for: Pfizer expert Group 2016, Sanofi expert Group 2017. No association with this study., Stefan Bergman: None declared, Kristina Forslind: None declared, Maria Andersson: None declared
DOI: 10.1136/annrheumdis-2019-eular.6938

AB0233

ASSESSMENT OF THE RESPONSE TO THERAPY WITH CSDMARDS AND BDMARDS IN PATIENTS WITH RHEUMATOID ARTHRITIS

Vladimira Boyadzhieva, Nikolay Stoilov. *University Hospital "St Ivan Rilski", Rheumatology, Sofia, Bulgaria*

Background: The significant economic burden of rheumatoid arthritis (RA) and new treatment options, consuming much of the healthcare resources, pay attention on appropriate management strategies. These strategies are due to the need for an adequate assessment of the response to treatment and changes if it is necessary. Currently used American College of Rheumatology (ACR) response criteria and disease activity score (DAS28-CRP) allows on time evaluation of the results and facilitate decision process when is the right time to switch patient's therapy.

Objectives: The aim of the study is to evaluate the effectiveness and response to the treatment with csDMARDS and/or corticosteroid and bDMARDS in patients with RA after one year follow up period using ACR 20/50/70/90 response criteria and DAS28-CRP.

Methods: A total of 220 patients with a mean age 55.05 ± 10.63 SD meeting the 1987 ACR classification criteria for RA were included in the study. The mean disease duration was 9.97 ± 5.78 SD. Patients were stratified according to treatment regimens into age-matched treatment groups: 96 on csDMARDS and/or corticosteroid (Sulfasalazine ? 15, Methylprednisolone ? 15, Chloroquine - 16, Methotrexate - 20, Methotrexate and Methylprednisolone -30) and 124 on bDMARD (Tocilizumab ? 30, Certolizumab pegol ? 16, Golimumab - 22, Etanercept - 20, Adalimumab - 20, Rituximab - 16). All participants were evaluated for ACR 20/50/70/90 response to therapy on 6th and 12th month of the follow up period, DAS28-CRP was calculated via licensed calculator. Comparison was performed by analysis variance ANOVA

Results: The majority of patients (71%) on bDMARDS achieved ACR 50 response criteria and 16% achieved ACR 70, indicating a significantly better therapeutic effect compared to the group on csDMARDS (ACR 50 - 11% ACR 70-1%) at 6th month follow up ($p < 0.001$, $p = 0.002$). Only 3% of bDMARDS patients achieved ACR 90 response on the 6th month. Most of the patients on csDMARDS (64%) reach ACR 20 response criteria after the same period of therapy. At 6th month of follow-up, best response was achieved by the patients on combined MTX and corticosteroid (CS) therapy (50%) achieved ACR 20, 27% - ACR 50 and 3% - ACR 70. At the 12th month, a significant proportion of patients on bDMARDS reached the ACR 50 (36%) and ACR 70 (52%) response criteria compared to the csDMARDS group (ACR 50 - 20%, ACR 70 - 10%) ($p = 0.0102$; $p < 0.001$). We evaluated the response to therapy according to the DAS28-CRP categories. On baseline patients on bDMARDS had higher disease activity in comparison to patients on csDMARDS ($p=0.0004$). In the follow-up to the 6th month, we observed a significant reduction in the proportion of patients (on bDMARDS) with severe disease activity - 2.4%, compared to csDMARDS group - 13.5% ($p = 0.0019$). In 26.6% of patients on bDMARDS, we found minimal disease activity, which was significantly more than 13.5% for the csDMARDS group ($p = 0.0186$). A remission was found in 5.6% of patients on bDMARDS which is significantly less than 14.6% for the group on csDMARDS. In contrast to the results obtained at the 6th month of follow-up, at 12th month we observed significantly higher percentage of patients on bDMARDS who had reached the remission category - 29.8%, in comparison with csDMARDS group - 12.5%, ($p = 0.0025$).

Conclusion: Patient treated with bDMARDS showed better response to treatment and lower disease activity than those on therapy with csDMARDS within a period of 12 months of treatment. Current management strategies should focus on switching on another medication earlier in the treatment plan when patients inadequately respond to therapy after 6 months of follow up.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4236

AB0234

AN INTEGRATED PROTEOMICS AND ANTIBODY ANALYSIS OF THE U-ACT-EARLY TRIAL TO IDENTIFY MARKERS OF TREATMENT RESPONSE AND DISEASE PROGRESSION IN EARLY RHEUMATOID ARTHRITIS

Francesco Brizzi¹, Suleiman A. Khan², Marco Prunotto¹, Jenny Devenport³, Attila Pethoe-Schramm¹, Michelle Borm⁴, Johannes Wj Bijsma⁵, Johannes W. G. Jacobs⁵, Floris Lafeber⁵, Paco Welsing⁵, Tero Aittokallio^{2,6}, Antoine Soubret¹.
¹F. Hoffmann-La Roche, Basel, Switzerland; ²Institute for Molecular Medicine Finland FIMM, University of Helsinki, Helsinki, Finland; ³Genentech, Inc., South San Francisco, United States of America; ⁴Roche Nederland BV, Woerden, Netherlands; ⁵University Medical Center Utrecht, Utrecht, Netherlands; ⁶University of Turku, Department of Mathematics and Statistics, Turku, Finland

Background: Predictive markers of treatment response in rheumatoid arthritis (RA) are necessary to stratify patients to devise personalised treatment strategies. The U-Act-Early trial [1], in which disease-modifying antirheumatic drugs (DMARD) naïve early RA patients initiated either methotrexate (MTX) or tocilizumab (TCZ) in monotherapy or in combination, offers a unique opportunity to assess the capacity of many baseline -omics markers (i.e. genomics, transcriptomics, proteomics, metabolomics, auto-antibodies) to predict clinical outcomes.

Objectives: To identify predictive markers of treatment response by developing a statistical machine learning methodology for correlating baseline clinical and -omics variables with clinical response data over time.

Methods: We developed a multi-omics approach that combines longitudinal statistical modelling of clinical data with Bayesian machine learning variable selection to identify the most predictive markers of treatment response. The time course of clinical outcomes was fit using a non-linear mixed longitudinal model, that characterise each patient of the U-Act-Early trial (n=317) by three parameters: a baseline value, an asymptotic value and a velocity (describing the speed at which the asymptote is reached). Supervised machine learning methods (such as Bayesian sparse regression as well as tree-based algorithms) were subsequently employed to find predictive markers for each of the longitudinal model parameters. We specifically focused on 85 auto-inflammatory proteins and 450 auto-antibodies markers, including 41 citrullinated-peptides antigens. The analysis was repeated for the three treatment-arms since DAS28-subscores response markers may vary across treatment strategies and clinical outcomes. We finally investigated whether predictions are enhanced by incorporating known networks of protein-protein interactions, rather than considering the predictive power of each protein independently.

Results: Preliminary results are that some proteins (e.g. sCD14, VEGF) were moderately predictive of the disease status at baseline. Specifically, the available inflammatory proteins predicted baseline DAS28, swollen and tender joints and ESR with R² values of 0.25, 0.32, 0.16 and 0 respectively. Surprisingly, these proteins did not improve the prediction of treatment-response (i.e. the velocity and asymptotic parameters) after controlling for the baseline disease status. Protein networks marginally improved prediction when compared to prediction of individual proteins. Citrullinated peptide antigens did not appear predictive of the baseline disease status nor the treatment response.

Conclusion: In our initial investigation inflammatory proteins, especially when considered as part of interaction networks, appear to be predictive of the baseline status of the disease but not of treatment-response. The most predictive marker of treatment response appears to be disease activity at baseline.

REFERENCES:

[1] Bijsma JWW, et al. Lancet. 2016;388:343-55.

Acknowledgement This study was funded by F. Hoffmann-LaRoche

Disclosure of Interests: Francesco Brizzi Employee of: F. Hoffmann-La Roche, Suleiman A. Khan: None declared, Marco Prunotto Employee of: F. Hoffmann-La Roche, Jenny Devenport Employee of: F. Hoffmann-La Roche, Attila Pethoe-Schramm Shareholder of: F. Hoffmann-La Roche, Employee of: F. Hoffmann-La Roche, Michelle Borm Employee of: An employee of Roche Nederland BV, Johannes Wj Bijsma Grant/research support from: The department of the author who included patients (JWJB) in the U-Act-Early trial received reimbursements from Roche Nederland BV. JWJB reported grants and fees from Roche, AbbVie, Bristol-Myers Squibb, Merck Sharp & Dohme, Pfizer, and UCB University Medical

Center Utrecht, Utrecht University, Consultant for: SUN Pharma, Speakers bureau: Lilly, Roche, Johannes W. G. Jacobs Grant/research support from: Roche, Consultant for: Roche, Floris Lafeber Shareholder of: Arthro-Save, Grant/research support from: FOREUM; Dutch Arthritis Society, Paco Welsing: None declared, Tero Aittokallio: None declared, Antoine Soubret Employee of: F. Hoffmann-La Roche
 DOI: 10.1136/annrheumdis-2019-eular.5049

AB0235

THE LIFETIME RISK OF KNEE AND HIP REPLACEMENT FOLLOWING A DIAGNOSIS OF RHEUMATOID ARTHRITIS: FINDINGS FROM ROUTINELY COLLECTED DATA FROM ENGLAND

Edward Burn¹, Christopher Edwards^{1,2}, David W. Murray¹, Alan Silman¹, Cyrus Cooper^{1,3}, Nigel Arden^{1,3}, Rafael Pinedo-Villanueva¹, Daniel Prieto-Alhambra^{1,4}.
¹NDORMS, University of Oxford, Oxford, United Kingdom; ²NIHR Wellcome Trust Clinical Research Facility, University Hospital Southampton, Southampton, United Kingdom; ³MRC Lifecourse Epidemiology Unit, Southampton University, Southampton, United Kingdom; ⁴GREMPAL Research Group, Idiap Jordi Gol and CIBERFes, Universitat Autònoma de Barcelona and Instituto de Salud Carlos III, Barcelona, Spain

Background: Understanding the lifetime risk of knee and hip replacement following a diagnosis of rheumatoid arthritis (RA) would help provide patients and clinicians with an indication of long-term prognosis and future health care utilisation.

Objectives: To estimate the lifetime risk of knee and hip replacement following a diagnosis of RA.

Methods: Routinely collected data from the English NHS was used to inform the analysis. Diagnosis of RA was identified using primary care records, with knee and hip replacement observed in linked hospital records. Parametric survival models were fitted for up to 15 years of follow up, with age, sex, Charlson comorbidity score, socioeconomic status, BMI, and smoking status included as explanatory variables. A decision model was used to combine and extrapolate survival models to estimate lifetime risk. Lifetime risk was estimated for individuals with average characteristics (median for continuous variables and mode for categorical ones). The partial effect of explanatory factors on lifetime risk was assessed by re-running models with participant profiles varying in the explanatory factor of interest, while holding other characteristics constant at their average.

Results: 13,961 study participants with a diagnosis of RA were included. Lifetime risk of knee replacement and hip replacement was estimated to be 22% (95%CI 16% to 29%) and 17% (11% to 26%) following a diagnosis of RA for the average patient profile (non-smoking women aged 64 with no other comorbidities, BMI of 27 and in the top socioeconomic quintile). Risks were higher for younger patients, but broadly similar for men and women. Comorbidities, socioeconomic status, BMI, and smoking status had relatively little impact on lifetime risk.

Conclusion: This study has estimated lifetime risk of knee and hip replacement following a diagnosis of RA, allowing for a better understanding of long-term prognosis and healthcare resource use. These risks are more than double those of the general UK population.

Disclosure of Interests: Edward Burn: None declared, Christopher Edwards Grant/research support from: Abbvie, BMS, Biogen, Celgene, Fresenius, Janssen, Lilly, Mundipharma, Pfizer, MSD, Novartis, Roche, Samsung, Sanofi, UCB, Consultant for: Abbvie, BMS, Biogen, Celgene, Fresenius, Janssen, Lilly, Mundipharma, Pfizer, MSD, Novartis, Roche, Samsung, Sanofi, UCB, Speakers bureau: Abbvie, BMS, Biogen, Celgene, Fresenius, Janssen, Lilly, Mundipharma, Pfizer, MSD, Novartis, Roche, Samsung, Sanofi, UCB, David W Murray Grant/research support from: Grants from Zimmer Biomet, Consultant for: Personal fees from Zimmer Biomet, Alan Silman: None declared, Cyrus Cooper Consultant for: Personal fees from Alliance for Better Bone Health, Amgen, Eli Lilly, GSK, Medtronic, Merck, Novartis, Pfizer, Roche, Servier, Takeda and UCB., Nigel Arden Grant/research support from: Grants from BIOIBERICA, and from MERCK., Consultant for: Personal fees from Flexion, from Regeneron, from Freshfields Bruckhaus Deringer, outside the submitted work., Rafael Pinedo-Villanueva: None declared, Daniel Prieto-Alhambra Grant/research support from: Grants from Amgen, UCB Biopharma and Servier outside the submitted work, Consultant for: UCB Biopharma, Speakers bureau: Amgen

DOI: 10.1136/annrheumdis-2019-eular.2432