OP0165

EULAR RECOMMENDATIONS FOR THE DIAGNOSIS AND THE MANAGEMENT OF RHEUMATIC IMMUNE-RELATED ADVERSE EVENTS DUE TO CANCER IMMUNOTHERAPY

Marie Kostine¹, Axel Finckh², Clifton Bingham³, Karen Visser⁴, Jan Leipe⁵, Hendrik Schulze-Koops⁶, Ernest Choy⁷, Karolina Benesova⁸, Timothy R. Radstake⁹, Andrew Cope¹⁰, Oliver Lambotte¹¹, Jacques-Eric Gottenberg¹², Yves Allenbach¹³, Marianne Visser¹⁴, Cindy Rusthoven¹⁵, Lone Thomasen¹⁶, Shahin Jamal¹⁷, Aurélien Marabelle¹⁸, James Larkin¹⁹, John Haanen²⁰, Leonard Calabrese²⁰, Xavier Mariette¹¹, Thierry Schaevebeke¹. ¹University Hospital of Bordeaux, Bordeaux, France; ²University Hospitals of Geneva, Geneva, Switzerland; ³Johns Hopkins University, Baltimore, United States of America; ⁴Haga Hospital, The Hague, Netherlands; ⁵University Medical Centre Mannheim, Mannheim, Germany; ⁶University of Munich, Munich, Germany; ⁷Cardiff University School of Medicine, Cardiff, United Kingdom; ⁸University Hospital Heidelberg, Heidelberg, Germany; ⁹University Medical Center Utrecht, Utrecht, Netherlands; ¹⁰King's College London, London, United Kingdom; ¹¹Le Kremlin-Bicêtre, Paris, France; ¹²University Hospital of Strasbourg, Strasbourg, France; ¹³Pitié-Salpêtrière University Hospital, Paris, France; ¹⁴EULAR PARE Patient Research Partner, Netherlands, Netherlands; ¹⁵Aarhus University Hospital, Aarhus, Denmark; ¹⁶University of British Columbia, Vancouver, Canada; ¹⁷Gustave Roussy, Paris, France; ¹⁸Royal Marsden Hospital NHS Foundation Trust, London, United Kingdom; ¹⁹Netherlands Cancer Institute, Amsterdam, Netherlands; ²⁰Cleveland Clinic, Cleveland, United States of America

Background: Rheumatic immune-related adverse events (irAEs) are increasingly recognized musculoskeletal manifestations in cancer patients receiving immune checkpoint targeted immunotherapy (1). Since they represent a spectrum of new clinical entities and a robust evidence base is lacking, a task force was convened to harmonize expert consensus regarding their identification and management due to the lack of dedicated clinical trials.

Objectives: To develop EULAR recommendations for the diagnosis and the management of rheumatic irAEs due to cancer immunotherapy, based on literature and expert opinion.

Methods: Recommendations were developed according to the 2014 EULAR Standard Operating Procedures. The task force consisted of 19 clinical experts from Europe and North America (14 rheumatologists, 2 internists and 3 oncologists), 1 clinical epidemiologist, 1 allied health professional and 2 patient representatives. During the first meeting, the group defined the focus of the task force, the target population, and formulated research questions. A systematic literature research was performed by one fellow (MK) with the help of a librarian. Based on available evidence and using a consensus procedure, recommendations were developed during a second meeting. The level of agreement was determined by an anonymous voting process.

Results: 4 overarching principles and 10 recommendations were developed. The overarching principles define the role of rheumatologists and highlight the shared decision-making process between patients, oncologists and rheumatologists. One recommendation addresses the referral process, two address the diagnosis, and five address the therapeutic strategy of cancer patients experiencing rheumatic, musculoskeletal, and systemic signs or symptoms while receiving immunotherapy. An additional recommendation was included to address pre-existing rheumatic conditions and the last focuses on the diagnostic approach before immunotherapy.

Conclusion: These recommendations provide the basis of a EULAR consensus on the diagnosis and the management of rheumatic irAEs.

REFERENCE:

- [1] Calabrese LH, Calabrese C, Cappelli LC. Rheumatic immune-related adverse events from cancer immunotherapy. *Nat Rev Rheumatol*. 2018 Oct;14(10):569-579.

Disclosure of Interests: Marie Kostine: None declared, Axel Finckh Grant/research support from: Bristol-Myers Squibb, Pfizer Inc, Consultant for: AbbVie, A2Bio, Bristol-Myers Squibb, MSD, Roche, Pfizer Inc, and UCB, Clifton Bingham Grant/research support from: BMS, Consultant for: AbbVie, BMS, Eli Lilly, Genentech/Roche, Janssen, Pfizer, Sanofi/Regeneron, Karen Visser: None declared, Jan Leipe Grant/research support from: Novartis, Pfizer; Scientific Support: Novartis, Pfizer, Consultant for: AbbVie, AstraZeneca, Bristol-Myers Squibb, Celgene, Hospira, Janssen-Cilag, LEO Pharma, Lilly, Novartis, Roche, Sanofi, Speakers bureau: AbbVie, Bristol-Myers Squibb, Celgene, Janssen-Cilag, Lilly, Novartis, MSD, Pfizer, Roche, Sanofi, UCB, Hendrik Schulze-Koops Grant/research support from: Novartis, Pfizer, Roche, Consultant for: AbbVie, Actelion, Amgen, AstraZeneca, Biogen International, BMS, Celgene, Chugai, GSK, Hospira, Janssen-Cilag, Leo Pharmaceuticals, Lilly, MSD, Medac, Merck, Novartis, Pfizer, Hexal Sandoz, Sanofi, Roche, UCB, Paid instructor for: AbbVie, Actelion, Amgen, AstraZeneca, Biogen International, BMS, Celgene, Chugai, GSK, Hospira, Janssen-Cilag, Leo Pharmaceuticals, Lilly, MSD, Medac, Merck, Novartis, Pfizer, Hexal Sandoz, Sanofi, Roche, UCB, Speakers bureau: AbbVie, Actelion,

Amgen, AstraZeneca, Biogen International, BMS, Celgene, Chugai, GSK, Hospira, Janssen-Cilag, Leo Pharmaceuticals, Lilly, MSD, Medac, Merck, Novartis, Pfizer, Hexal Sandoz, Sanofi, Roche, UCB, Ernest Choy Grant/research support from: Amgen, Bio-Cancer, Chugai Pharma, Ferring Pharmaceuticals, Novimmune, Pfizer, Roche, and Union Chimique Belge, Consultant for: AbbVie, Amgen, AstraZeneca, Biogen, BMS, Boehringer Ingelheim, Celgene, Chelsea Therapeutics, Chugai Pharma, Daiichi Sankyo, Eli Lilly, Ferring Pharmaceutical, GlaxoSmithKline, Hospira, ISIS, Jazz Pharmaceuticals, Janssen, MedImmune, Merrimack Pharmaceutical, Merck Sharp & Dohme, Napp, Novimmune, Novartis, ObsEva, Pfizer, Regeneron, Roche, R-Pharm, SynAct Pharma, Sanofi-Genzyme, Tonix and Union Chimique Belge, Speakers bureau: Amgen, BMS, Boehringer Ingelheim, Chugai Pharma, Eli Lilly, Hospira, Merck Sharp & Dohme, Novartis, Pfizer, Regeneron, Roche, Sanofi-Aventis, and Union Chimique Belge, Karolina Benesova Grant/research support from: Scientific Projects (IIT) I'm involved in as an investigator have been supported, but not initiated or designed by AbbVie and Novartis, Consultant for: Advisory board member on early disease detection for Novartis, Timothy R. Radstake: None declared, Andrew Cope: None declared, Oliver Lambotte: None declared, Jacques-Eric Gottenberg Grant/research support from: Bristol-Myers Squibb, Grant/research support from: Bristol-Myers Squibb, Consultant for: Bristol-Myers Squibb, Lilly, Pfizer, Sanofi-Genzyme, UCB Pharma, Consultant for: Bristol-Myers Squibb, Eli Lilly, UCB, Sanofi-Genzyme, Pfizer, Yves Allenbach: None declared, Marianne Visser: None declared, Cindy Rusthoven: None declared, Lone Thomasen: None declared, Shahin Jamal Consultant for: Consultant for AbbVie, Amgen, BMS, Eli Lilly, Pfizer, Janssen, Merck, UCB, Aurélien Marabelle: None declared, James Larkin: None declared, John Haanen Consultant for: Amgen, Merck, MSD, Helsinn, Tesaro and Hexal, Leonard Calabrese Consultant for: Bristol-Myers Squibb, Genentech and AstraZeneca, Xavier Mariette Grant/research support from: Servier, Consultant for: AstraZeneca, Bristol-Myers Squibb, GlaxoSmithKline, Janssen, Pfizer, UCB Pharma, Thierry Schaevebeke: None declared

DOI: 10.1136/annrheumdis-2019-eular.7372

OP0166

THE EFFICACY OF CALCINEURIN INHIBITORS IN PATIENTSWITH ADULT-ONSET STILL'S DISEASE: MULTI-CENTRE HISTORICAL COHORT STUDY

Masato Tarumi¹, Hiroyuki Nakamura¹, Yuichiro Fujieda², Hirohiko Kitagawa³, Ryo Hisada⁴, Yuka Shimizu⁴, Ikuma Nakagawa⁵, Akito Tsutsumi⁵, Takashi Kurita⁶, Hiroshi Kataoka⁷, Masaya Mukai⁷, Hideki Kasahara⁸, Atsushi Noguchi⁹, Yoshiharu Amasaki⁹, Tetsuya Horita¹, Tatsuya Atsumi². ¹Tomakomai City Hospital, Tomakomai, Japan; ²Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Rheumatology, Endocrinology and Nephrology, Sapporo, Japan; ³Kushiro Red Cross Hospital, Kushiro, Japan; ⁴Obihiro-Kosei General Hospital, Obihiro, Japan; ⁵Takikawa Municipal Hospital, Takikawa, Japan; ⁶Japanese Red Cross Kitami Hospital, Kitami, Japan; ⁷Sapporo City General Hospital, Sapporo, Japan; ⁸NTT-East Sapporo Hospital, Sapporo, Japan; ⁹Tonan Hospital, Sapporo, Japan

Background: Adult-onset Still's disease (AOSD) is a systemic inflammatory disorder generally responsive to corticosteroid (CS) therapy, whereas cases not responding CS monotherapy are frequently found in clinical practice. In such cases, methotrexate and/or biologics including TNF- α , IL-1, or IL-6 inhibitors are used [1]. However, further treatment options are required for refractory AOSD patients who are intolerant to methotrexate and/or biologics. Calcineurin inhibitors (CNI) downregulate T cell activation through inhibiting IL-2 transcription and signal transduction. Therefore, they are reasonable therapeutic medication for AOSD considering T cells and subsequently activated macrophages play a pathophysiological role in AOSD [2]. Nevertheless, only a few case series have indicated effects of CNI for difficult AOSD.

Objectives: To evaluate the efficacy of CNI in patients with AOSD.

Methods: This is a multi-centre historical cohort study comprised the consecutive patients with AOSD in accordance with Yamaguchi classification criteria. The primary endpoint was set as the time from the initiation of treatment to event defined as death of any causes or relapse of AOSD requiring an increase of CS dose. Secondary endpoints were set as persistency rate of CNI and safety. Based on the recurrent event data analysis, these endpoints were evaluated for each event. We compared the frequency of events between two groups according to the treatment that included calcineurin inhibitors (CNI+) or conventional therapy without calcineurin inhibitors (CNI-).

Results: One hundred seventy-eight patients were enrolled in this study. Mean age was 46 year-old, and median follow-up period 36 months. Seventy-one events in 56 patients were treated with therapeutic regimen including CNI (CNI+, cyclosporine: 14, tacrolimus: 60), and 176 events in 138 patients were treated with the conventional therapy excluding CNI (CNI-). CNI were used in AOSD patients with recurrent history, high ferritin level, serositis, hemophagocytic syndrome (HPS) and/or disseminated intravascular coagulation (DIC). The CNI+ group had longer event-free survival than the CNI- group (83% versus 75% at 5th

year). The hazard ratio (HR) was 0.57 (95% CI, 0.32-0.99) after adjustment of initial dose of CS and other concomitant medications. Subgroup analysis showed that CNI were more effective for AOSD patients with high ALT levels (≥ 80 IU/L) and/or severe complications such as HPS and DIC. The persistency rate of CNI was 71% at 5th year. Adverse events occurred more frequently in the CNI+ group than in the CNI- group (18% versus 8%, $p = 0.04$); however, serious adverse events did not increase in the CNI+ group (3% versus 2%). AOSD-related mortality had never been found in the observation period. One patient had a fatal course due to septic shock in the CNI+ group.

Conclusion: Our retrospective analysis suggested that CNI could be an additional option for treating AOSD with acceptable safety.

REFERENCES:

- [1] Efthimiou P, et al. Diagnosis and management of adult onset Still's disease. *Ann Rheum Dis*. 2006.
- [2] Chen DY, et al. Predominance of Th1 cytokine in peripheral blood and pathological tissues of patients with active untreated adult onset Still's disease *Ann Rheum Dis*. 2004.

Disclosure of Interests: Masato Tarumi: None declared, Hiroyuki Nakamura: None declared, Yuichiro Fujieda: None declared, Hirohiko Kitagawa: None declared, Ryo Hisada: None declared, Yuka Shimizu: None declared, Ikuma Nakagawa: None declared, Akito Tsutsumi: None declared, Takashi Kurita: None declared, Hiroshi Kataoka: None declared, Masaya Mukai: None declared, Hideki Kasahara: None declared, Atsushi Noguchi: None declared, Yoshiharu Amasaki: None declared, Tetsuya Horita: None declared, Tatsuya Atsumi Grant/research support from: Astellas Pharma Inc., Takeda Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., Chugai Pharmaceutical Co., Ltd., Daiichi Sankyo Co. Ltd., Otsuka Pharmaceutical Co., Ltd. and Pfizer Inc., Alexion Inc., Bayer Yakuhin, Ltd., Otsuka Pharmaceutical Co., Ltd., Chugai Pharmaceutical Co., Ltd., Takeda Pharmaceutical Co., Ltd., Eisai Co., Ltd., Bristol-Myers Squibb Co., Daiichi Sankyo Co. Ltd., Mitsubishi Tanabe Pharma Co., Asahi Kasei Pharma Co., Consultant for: ONO PHARMACEUTICAL CO., LTD Sanofi K.K., Daiichi Sankyo Co., Ltd., Pfizer Inc., Speakers bureau: Mitsubishi Tanabe Pharma Co., Chugai Pharmaceutical Co., Ltd., Astellas Pharma Inc., Takeda Pharmaceutical Co., Ltd., Pfizer Inc., Daiichi Sankyo Co. Ltd., Bristol-Myers Squibb Co., Eli Lilly Japan K.K.

DOI: 10.1136/annrheumdis-2019-eular.3182

OP0167 B LYMPHOCYTES DIRECTLY CONTRIBUTE TO TISSUE FIBROSIS IN IGG4-RELATED DISEASE

Emanuel Della Torre^{1,2}, Elena Rigamonti¹, Cory Perugino², Massimo Falconi³, John H. Stone², Angelo Manfredi¹, Shiv Pillai². ¹San Raffaele Scientific Institute, Unit of Immunology, Rheumatology, Allergy and Rare Diseases, Milan, Italy; ²Ragon Institute of MGH, MIT, HARVARD, Cambridge, United States of America; ³San Raffaele Scientific Institute, Pancreas Center, Milan, Italy

Background: IgG4-related disease (IgG4-RD) is a fibro-inflammatory condition marked by rapid clinical improvement after selective depletion of B lymphocytes with rituximab. This feature suggests that B cells might participate in fibrogenesis and wound healing (1-3).

Objectives: In the present work we aimed to demonstrate that B lymphocytes contribute directly to tissue fibrosis in IgG4-RD.

Methods: Total circulating CD19+ B-lymphocytes, naïve B cells, memory B cells, or plasmablasts from IgG4-RD patients were cultivated with human fibroblasts. Pro-fibrotic soluble factors and collagen production in the co-cultures were assessed by ELISA and Luminex assays. RNA-sequencing and quantitative RT-PCR were used to assess fibroblast activation in the presence of B cells, as well as the induction of pro-fibrotic pathways in B cell subsets. Relevant pro-fibrotic and inflammatory molecules were confirmed in vitro by functional experiments and on IgG4-RD tissue sections by multi-color immunofluorescence studies.

Results: B cells from IgG4-RD patients (i) produced the pro-fibrotic molecule PDGF-B and stimulated collagen production by fibroblasts; (ii) expressed enzymes implicated in extracellular matrix remodeling such as LOXL2; (iii) produced the chemotactic factors CCL-4, CCL-5, and CCL-11; and (iv) induced the production of these same chemokines by activated fibroblasts. Plasmablasts expressed sets of genes implicated in fibroblast activation and proliferation, and therefore represent cells with intrinsic pro-fibrotic properties

Conclusion: We have demonstrated that B cells, contribute directly to tissue fibrosis in IgG4-RD. These unanticipated pro-fibrotic properties of B lymphocytes, particularly of plasmablasts, might be relevant for fibrogenesis in other fibro-inflammatory disorders and for wound healing processes in physiological conditions.

REFERENCES:

- [1] Della-Torre E, Lanzillotta M, Doglioni C. Immunology of IgG4-related disease. *Clin Exp Immunol* 2015;181:191-206.
- [2] Carruthers MN, Topazian MD, Khosroshahi A, Witzig TE, Wallace ZS, Hart PA, et al. Rituximab for IgG4-related disease: a prospective, open-label trial. *Ann Rheum Dis*. 2015;74:1171-7.
- [3] Della-Torre E, Feeney E, Deshpande V, Mattoo H, Mahajan V, Kulikova M, et al. B-cell depletion attenuates serological biomarkers of fibrosis and myofibroblast activation in IgG4-related disease. *Ann Rheum Dis* 2015;74:2236-43.

Disclosure of Interests: Emanuel Della Torre: None declared, Elena Rigamonti: None declared, Cory Perugino: None declared, Massimo Falconi: None declared, John H. Stone Grant/research support from: F. Hoffmann-La Roche, Genentech, Xencor, Consultant for: Chugai, F. Hoffmann-La Roche, Genentech, Xencor, Angelo Manfredi: None declared, Shiv Pillai: None declared

DOI: 10.1136/annrheumdis-2019-eular.1741

OP0168 DERIVATION AND VALIDATION OF A NEW STILL ACTIVITY SCORE (SAS)

Umut Kalyoncu¹, Timuçin Kaşifoğlu², Ahmet Omma³, Cemal Bes⁴, Muhammet Çınar⁵, Hakan Emmungil⁶, Orhan Küçükşahin⁷, Servet Akar⁸, Kenan Aksu⁹, Fatih Yıldız¹⁰, Nilüfer Alpay Kanitez¹¹, Abdulsamet Erden¹, Emre Bilgin¹, Sezin Turan⁶, Ediz Dalkılıç¹², Selime Erzurum¹², Kadir Mutlu Hayran¹³. ¹Hacettepe University Medical School Internal Medicine, Rheumatology, Ankara, Turkey; ²Eskişehir Osmangazi University Faculty of Medicine, Rheumatology, Ankara, Turkey; ³Ankara Numune Training and Research Hospital, Rheumatology, Ankara, Turkey; ⁴University of Health Sciences, Bakırköy Sadi Konuk Research and Training Hospital, Rheumatology, Ankara, Turkey; ⁵Gulhane Training and Research Hospital, Rheumatology, Ankara, Turkey; ⁶Trakya University Medical School, Rheumatology, Edirne, Turkey; ⁷Liv Hospital Ankara, Rheumatology, Ankara, Turkey; ⁸Izmir Katip Çelebi University, Rheumatology, İzmir, Turkey; ⁹Ege University Medical Faculty, Rheumatology, Ankara, Turkey; ¹⁰Van Research and Training Hospital, Rheumatology, Ankara, Turkey; ¹¹Koc University Medical School, Rheumatology, Ankara, Turkey; ¹²Uludağ University Medical School, Rheumatology, Ankara, Turkey; ¹³Hacettepe University Medical School, Preventive Oncology, Ankara, Turkey

Background: Adult-onset Still's disease (AOSD) is a rare, chronic, multisystem and auto-inflammatory disorder. Although several classification criteria exist for the diagnosis of AOSD, there is no valid method for the assessment of disease activity for these patients.

Objectives: To determine a simple method for the assessment of AOSD activity.

Methods: We conducted a cross-sectional, multicenter study for assessment of disease activity in AOSD patients. All AOSD patients were fulfilled the Yamaguchi criteria. Age, sex, disease duration and current symptoms were recorded. Erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), leukocyte count, and ferritin levels were also recorded. Visual analog scale (VAS) (0-10 cm) and likert scale (as remission, low, moderate, severe, and more severe disease activity) were used for physician's (PhGA) and patient's (PtGA) global assessment of disease activity. Patients were enrolled to "derivation cohort" (n=125) and subsequently "validation cohort" (n=72). Ordinal logistic regression model was used to derive Still Activity score (SAS).

Results: Overall, 197 (66.5% female) AOSD patients were enrolled. Mean age was 39 (13 years and median disease duration was 2 (0-30) years. Cross-sectional frequency of AOSD findings were as follow; fever 69 (35%), rash 55 (27.9%), arthritis 57 (28.9%), arthralgia 112 (56.9%), sore throat 58 (29.4%), myalgia 80 (40.6%), lymphadenopathy 25 (12.7%), splenomegaly 37 (18.8%), hepatomegaly 23 (11.7%), abdominal pain 8 (4.1%), pleuritic pain 8 (4.1%), pericarditis 4 (2%), hemophagocytic syndrome 4 (2%). Mean (SD) ESR (mm/h), CRP (mg/dl), ferritin and leukocyte count were 32.5 (31), 43.4 (65.3), 1169 (3088) and 10356 (4821), respectively. Mean PtGA and PhGA was 3.86 (3.29), and 3.05 (3.06). SAS derived for the assessment of AOSD disease activity is given in Table 1.