AB0297

HOW OFTEN IS THE COMBINATION OF NSAID-INDUCED LESIONS OF THE UPPER AND LOWER GI TRACT?

Anna Balabantseva, Andrey Karateev. V. A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: Non-steroidal anti-inflammatory drugs (NSAIDs) can damage all parts of the gastrointestinal tract (GIT). However, the frequency of the combination of lesions in different parts of the gastrointestinal tract has not been studied.

Objectives: To assess the frequency of the combination of NSAID-induced lesions of the upper GI tract, small and large intestine.

Methods: The study group consisted of 112 patients with rheumatic diseases (62.5% of women, 56.2 ± 14.6 years), who regularly took NSAIDs. All patients underwent endoscopy of the upper GI tract and colon. Capsule endoscopy was performed in 35 patients with signs of NSAID-gastropathy.

Results: NSAID-gastropathy (erosions and/or gastric or duodenum ulcers) were detected in 43.8%, NSAID enteropathy (hemorrhages, erosions and ulcers of the small intestine) in 68.6%, NSAIDs-colopathy (hemorrhages, erosions and ulcers of the colon) - 14.3% patients. The combination of NSAID-gastro- and colonopathy was detected in 28.6% of patients (odds ratio 12.2, 95% CI 2.619-56.84), the combination of NSAID-gastro, entero- and colopathy - in 10 patients (20.4% of the number of patients with NSAID-gastropathy). There was a significant risk association for the development of NSAID-induced lesions in all parts of the GI tract with a diagnosis of spondyloarthritis, the presence of abdominal pain, small intestine bacterial overgrowth, as well as in the polymorphism of the CYP2C19 gene (CYP2C19 * 17 * 1/* 17 allele).

Conclusion: The combined NSAID-induced damage of various parts of the GI tract is a frequent and serious pathology requiring complex diagnostics and the combined use of prophylactic agents with a different mechanism of action.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6153

AB0298

COMPLICATIONS OF RHEUMATOID ARTHRITIS AND ALLIED DISORDERS— A STATISTICAL COMORBIDITY STUDY OF 234 AUTOPSY PATIENTS

Miklós Bély¹, Ágnes Apáthy². ¹Hospital of the Order of the Brothers Saint John of God in Budapest, Department of Pathology, Budapest, Hungary; ²St. Margaret Clinic, Budapest, Department of Rheumatology, Budapest, Hungary

Background: Complications of rheumatoid arthritis (RA) may modify the clinical course and symptoms of allied disorders leading to missed diagnosis or late recognition of associated diseases.

Objectives: The aim of this study was to determine the possible role of classic complications of RA: systemic autoimmune vasculitis (AV), AA amyloidosis (AAa), lethal cardiac insufficiency (CI) caused by endo-, myo- or pancarditis, furthermore lethal septic infection (SI) on prevalence and mortality of coexistent associated diseases: atherosclerosis (Ath), hypertension (HT), type 2 diabetes mellitus (DM) and tuberculosis (Tb) with miliary dissemination (mTb).

Methods: 234 non- selected autopsy patients with RA were studied. RA was confirmed clinically according to the criteria of the ARA.

The presence of AV, AAa, CI and SI was determined at autopsy and confirmed by a detailed review of extensive histological material. The prevalence and mortality of associated diseases Ath, HT, DM, Tb or mTb were analyzed retrospectively, reviewing the clinical and pathological reports.

The link between AV, AAa, CI or SI and Ath, HT, DM, Tb or mTb was analyzed by (χ^2) test.

Results: RA was complicated by AV in 43 (18.4%), by AAa in 48 (20.5%), by CI with lethal outcome in 15 (6.4%), and by lethal SI in 33 (14.1%) of 234 patients.

RA associated with severe Ath in 106 (45.3%), with HT in 41 (17.5%), with DM in 41 (17.5%), and with Tb in 28 (11.9%), with active mTb in 9 (3.8%) of 234 patients.

As a basic disease Ath led to death in 61 (26.1% of 106), HT in 2 (0.9% of 41), DM in none (0% of 41), and Tb with mTb in 3 (1.3% of 28) of 234 patients. Tb without miliary dissemination was not lethal in our patient population.

The statistical links ("p" values of significance) between complications of RA and prevalence or mortality of allied disorders are summarized in Table.

(*asterisk indicates a negative value of associations' coefficient with inverse relationship).

Complications of RA	AV n=43 of 234	AAa n=48 of 234	CI n=15 of 234	SI n=33 of 234
Allied disorders in RA				
Ath n=106 of 234	$\chi^2=6.43^*$, p<0.01	$\chi^2=10.04^*$, p<0.001	$\chi^2=0.92^*$, p<0.33	$\chi^2=5.03^*$, p<0.02
Ath lethal n=61 of 106	$\chi^2=15.92^*$, p<0.00	$\chi^2=11.04^*$, p<0.00	$\chi^2=4.29^*$, p<0.03	$\chi^2=12.01^*$, p<0.00
HT 41 of 234	$\chi^2=3.30^*$, p<0.07	$\chi^2=0.03^*$, p<0.86	$\chi^2=0.00^*$, p<0.92	$\chi^2=1.27^*$, p<0.25
HT lethal n=2 of 41	$\chi^2=0.05^*$, p<0.80	$\chi^2=0.02$, p<0.87	$\chi^2=1.16^*$, p<0.28	$\chi^2=0.19^*$, p<0.65
DM 41 of 234	$\chi^2=0.04$, p<0.8	$\chi^2=0.84^*$, p<0.35	$\chi^2=0.92$, p<0.33	$\chi^2=0.01$, p<0.91
Tb 28 of 234	$\chi^2=4.02$, p<0.04	$\chi^2=2.61^*$, p<0.10	$\chi^2=0.33$, p<0.56	$\chi^2=0.37$, p<0.54
Tb lethal n=3 of 28	$\chi^2=0.00^*$, p<0.93	$\chi^2=0.02^*$, p<0.86	$\chi^2=0.53^*$, p<0.46	$\chi^2=0.01^*$, p<0.89
mTb 9 of 28	$\chi^2=4.24$, p<0.03	$\chi^2=0.08$, p<0.77	$\chi^2=0.01$, p<0.91	$\chi^2=0.05$, p<0.82
mTb lethal n=3 of 9	$\chi^2=0.00^*$, p<0.93	$\chi^2=0.02^*$, p<0.86	$\chi^2=0.53^*$, p<0.46	$\chi^2=0.01^*$, p<0.89

Conclusion: The inverse correlations between AV, AAa, CI, SI and Ath, HT, DM, Tb or mTb indicate that the prevalence and mortality of allied disorders were not influenced basically by the complications of RA.

The consequently inverse and (in most cases) significant correlations between prevalence of AV, AAa, CI, SI and the prevalence and mortality of Ath show that these are independent entities in RA. The AV, AAa, CI and SI are the most important complications of RA, and are characterized severe forms of disease, mostly involving younger patients, with an earlier onset (without pronounced Ath); while Ath is basically an age dependent phenomenon, characteristically present in RA patients with advanced age. RA patients with Ath may represent a special group of RA, characterized by lower incidence of AV, AAa, CI or SI, and a better prognosis.

The positive and significant relations AV to Tb or mTb suggest an increased risk of Tb, e.g. the presence of AV may promote Tb or endogenous exacerbation and miliary dissemination of Tb.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.1273

AB0299

COXITIS DURING CHRONIC INFLAMMATORY RHEUMATISM

Hedia Ben Abla, Sonia Rekik, Samia Jammali, Soumaya Boussaid, Hela Sahli, Elhem Cheour, Mohamed Elleuch. *la rabta, tunis, Tunisia*

Background: The occurrence of coxitis in chronic inflammatory rheumatism is a prognostic factor of great importance given the functional repercussions of the involvement of the coxofemoral joint.

Objectives: The aim of this work is to evaluate the prevalence of this lesion in chronic inflammatory rheumatism (PR/JIA/SpA) and to describe the evolution of this disorder according to the pathology.

Methods: Eighty-two patients with chronic inflammatory rheumatism were enrolled. These patients were followed by the rheumatology department rabta since 2004 until March 2018. The clinical demographics, activity scores, treatment background (all our patients in this series were under biotherapy) and the need PTH were identified.

Results: The eighty-two patients were distributed as follows: 36.58% (n = 30) patients with RA, 36.58% (n = 30) SpA and 26.83 (n = 22) JIA. The average age was 48.3 years, 47.6 years and 32.17 years. For the rheumatoid arthritis population 26.67% (n = 8) of the patients had a coxite of which 62.5% (n = 5) was bilateral and 37.5% (n = 3) unilateral. The mean time to discovery of coxite was 11.7 years and the evolution was in 50% (n = 4) to worsening requiring in 3 cases a bilateral PTH and a case of synoviorrhesis.

In the SpA group, 23.33% (n = 7) of patients had coxitis, of which 57.14% (n = 4) were bilateral and 32.86% (n = 3) unilateral. The average time to discovery was 8.22 years. PTH was performed in 71.43% (n = 5) of patients.

For the JIA population, 45.45% (n = 10) of the patients had a coxite of which 80% (n = 8) was bilateral and 20% unilateral. The average time to discovery of this lesion was 14 years, with 60% of cases progressing to an aggravation requiring 40% of the cases for synoviorrhesis and 60% for PTH.

Conclusion: SpA is the most common chronic inflammatory rheumatism with coxofemoral involvement with the most pejorative evolution despite the biological treatment