OP0174

PARTIAL KNEE REPLACEMENT IS ASSOCIATED WITH A LOWER RISK OF VENOUS THROMBOEMBOLISM AND OPIOID USE THAN TOTAL KNEE REPLACEMENT BUT INCREASED RISK OF LONG-TERM REVISION: A MULTINATIONAL, MULTI-DATABASE, PROPENSITY SCORE-MATCHED, COHORT ANALYSIS INCLUDING OVER 280,000 PATIENTS

Daniel Prieto-Alhambra¹, Edward Burn¹, James Weaver², Anthony G Sena², Henry Morgan Stewart³, Patrick Ryan². ¹Centre for Statistics in Medicine, NDORMS, University of Oxford, Oxford, United Kingdom; ²Janssen Research and Development, Titusville, United States of America; ³IQVIA, London, United Kingdom

Background: Knee replacement is one of the few effective treatments for severe knee osteoarthritis. A number of surgical interventions are available, including two main types: partial or total knee replacement.

Objectives: We aimed to assess the outcomes of partial compared to total knee replacement.

Methods: We conducted a multi-database propensity score-matched cohort study. Data was obtained from 4 US claims databases (IBM MarketScan[®] Commercial Database (CCAE), IBM MarketScan[®] Medicare Supplemental Database (MDCR), Optum[®] de-identified Clinformatics[®] Datamart, Extended - Date of Death (Optum), and Pharmetrics), and 1 UK primary care electronic medical record database (THIN).

All people aged 40 years or older at the time of knee replacement surgery were included. Data were mapped to a common data model (OMOP), and processed using analytical tools developed by the OHDSI community.

Participants were followed up from the date of their first knee replacement and for up to 5 years. Outcomes included short-term (60-day) post-operative complications (infection, venous thromboembolism, mortality, readmission), opioid use in the 3-12 months post-surgery as a proxy for persistent pain, and 5-year revision risk.

Propensity score matching (up to 1:10) was used to control for all available confounders, and negative control outcomes and calibration to minimize the impact of residual confounding. Separate Cox regression models were fitted for each outcome to estimate calibrated HR (cHR) and 95% confidence intervals related to partial, using total knee replacement as a reference group.

Results: A total of 32,379 and 250,377 patients receiving partial and total knee replacement respectively were matched. Short-term risk of venous thromboembolism was consistently lower in the former, with cHR ranging from 0.47 [0.32-0.71] in MDCR to 0.76 [0.59-0.99] in CCAC. No differences were observed in the risk of other complications. The use of opioids in months 3 to 12 after surgery was also lower in subjects receiving partial knee replacement, with cHR ranging from 0.70 [0.57-0.90] in THIN to 0.86 [0.78-0.96] in Optum.

Conversely, risk of 5-year revision was consistently higher amongst those undergoing partial compared to total knee replacement: cHR ranged from 1.51 [1.24-1.88] in CCAC to 2.16 [1.63-3.15] in MDCR.

Conclusion: Partial knee replacement is associated with a 25% to 50% reduction in short-term risk of venous thromboembolism, and a 15%-30% lower risk of persistent pain post-surgery. In contrast, partial knee replacement results in a 50% to >100% excess risk of revision in the longer term. This information should be used to individualize surgical indications for patients with severe knee osteoarthritis.

Disclosure of Interests: Daniel Prieto-Alhambra Grant/research support from: Grants from Amgen, UCB Biopharma and Servier outside the submitted work, Consultant for: UCB Biopharma, Speakers bureau: Amgen, Edward Burn: None declared, James Weaver Shareholder of: Shareholder in Johnson & Johnson, Employee of: Full-time employee of Janssen Research & Development, a pharmaceutical company of Johnson & Johnson, Anthony G Sena Shareholder of: Shareholder in Johnson & Johnson, Employee of: Full-time employee of Janssen Research & Development, a pharmaceutical company of Johnson & Johnson, Henry Morgan Stewart Employee of: Full-time employee of IQVIA, Patrick Ryan Shareholder of: Shareholder in Johnson & Johnson, Employee of: Full-time employee of Janssen Research & Development, a pharmaceutical company of Johnson & Johnson.

DOI: 10.1136/annrheumdis-2019-eular.886

OP0175

METHOTREXATE IN PATIENTS WITH HAND OSTEOARTHRITIS REFRACTORY TO ANALGESICS: A RANDOMISED, DOUBLE-BLIND, PLACEBO CONTROLLED TRIAL

Stéphanie Ferrero¹, Ruth Wittoek², Edem Allado³, Eric Fontas⁴, Coralie Cruzel⁴, Liana Euler-Ziegler¹, Damien Loeuille³, Christian Roux¹. ¹University of Nice cote d'azur/CHU Nice, Department of rheumatology, Nice, France; ²Ghent, Ghent, Belgium; ³Nancy, Nancy, France; ⁴Nice, Nice, France

Background: To date, no studies have been published on the effect of Methotrexate (MTX) in hand osteoarthritis (HOA).

Objectives: The aim of our study is to examine the effect of methotrexate on pain in symptomatic erosive HOA, as well as its functional and structural effect using MRI and radiography.

Methods: This prospective, single-center, randomized, double-blind, placebo-controlled study followed patients with HOA over 12 months. Patients were randomized to Methotrexate 10 mg per week or placebo.

The primary endpoint was pain assessment at 3 months, and secondary endpoints were clinical (Visual Analogue Scale "VAS" pain at 12 months), and radiographic (radiographic anatomical score, then the Ghent University Score System "GUSS" and MRI).

All subjects aged between 45 and 85 years, who had apparent NSAIDs and pain traitement failure with an VAS greater than 4/10, and at least one erosive joint were included.

Results: Out of 71 evaluated patients with symptomatic erosive HOA, and 64 matched the inclusion criteria and were randomized. Among them, 32 patients were randomized in the placebo group and 32 in the Methotrexate group. There was no significant difference between the two groups on the primary endpoint at 3 months, with an average decrease in VAS in the placebo group of 8.36 mm (+/- 25.15), and 17.47 mm (+/- 28.37) in the treatment group (p = 0.1797). At 12 months, no difference was found in the evolution of pain between the two groups (p = 0.6045) as well as in the functional evolution. In subjects taking Methotrexate, erosive joints move more to a remodeling phase undergoing treatment (27%) than placebo subjects (15%) (p = 0.0278). They also seem to evolve less towards erosion when their joints present a space loss (29% become erosive under placebo, against 8% in the Methotrexate group) (p = 0.0879). No predictor of erosive outcome was found in patients with non-erosive hand osteoarthritis.

Conclusion: Our study does not show superior efficacy of MTX over placebo over pain and function in subjects with erosive HOA. Our results suggest a possible structural effect on the radiological evolution of HOA under Methotrexate.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4373

OP0176

HEALTH-RELATED QUALITY OF LIFE IN HAND OSTEOARTHRITIS PATIENTS FROM THE GENERAL POPULATION AND THE OUTPATIENT CLINIC

Marieke Loeft¹, Wendy Damman¹, Renée de Mutser², Frits Rosendaal², Margreet Kloppenburg^{1,2}. ¹Leiden University Medical Center (LUMC), Rheumatology, Leiden, Netherlands; ²Leiden University Medical Center (LUMC), Epidemiology, Leiden, Netherlands

Background: In osteoarthritis (OA) patients health-related quality of life (HRQoL) is decreased. Whether patients with OA seeking care in a rheumatology outpatient clinic experience more impact on HRQoL than those in the general population is unknown.

Objectives: To investigate the impact of hand OA on physical and mental HRQoL in the general population, and to investigate the difference in impact between patients who have, and who have not been referred to a medical specialist.

Methods: In the population-based Netherlands Epidemiology of Obesity (NEO) study, middle-aged participants were recruited from the greater area of Leiden. In the Hand Osteoarthritis in Secondary care (HOSTAS) study, patients with a rheumatologist's diagnosis of primary hand OA were recruited from the outpatient clinic at the Leiden University Medical Center, a secondary and tertiary referral center. In both cohorts, hand and knee OA was defined by the ACR clinical classification criteria. Patients with fibromyalgia or inflammatory rheumatic conditions were excluded. For the current analyses, only patients classified with hand OA alone were included. HRQoL was measured with the SF-36 questionnaire; we calculated standardized scores on a scale of 0 to 100 with subsequent application of a norm-based transformation (mean 50, standard deviation 10). Linear regression analyses, corrected for age, sex, education, ethnicity and BMI, were used to study cross-sectional associations between OA and HRQoL. Data are presented as regression coefficients with 95% confidence intervals (CI). Because a suitable reference group without OA was lacking in the HOSTAS study, these patients were compared to the normative value of 50. Previous research concluded a minimal clinically important difference of 2 points on the SF-36 scale, which was used to assess clinical relevance of differences in scores.

Results: Of the 6,334 NEO participants 8% were classified with only hand OA and 4% were classified concurrent hand and knee OA. The HOSTAS cohort consisted of a total of 538 patients with hand OA, of whom 57% fulfilled the ACR criteria for only hand OA and 32% was defined with concurrent hand and knee OA. In the population-based NEO study, mean PCS was reduced with -2.4 (-3.6; -1.3) in participants with only hand OA, compared to participants without hand or knee OA (table 1). The subscales bodily pain and physical functioning were affected the most with mean differences of -3.4 (-4.6; -2.2) and -2.1 (-3.0; -1.1). Mental HRQoL was not reduced in participants with only hand OA, compared to participants without OA. In the population-based cohort 14% of participants with hand OA reported to have visited a medical specialist for OA. Participants with hand OA that have been referred to a medical specialist showed a lower physical HRQoL with a