OP0179

DO DISTINCT PAIN PHENOTYPES HAVE DIFFERENT RISK OF KNEE REPLACEMENT: A 13.7-YEAR FOLLOW-UP STUDY?

Feng Pan¹, Jing Tian¹, Ishanka Munugoda¹, Stephen Graves², Michelle Lorimer³, Flavia Cicuttini⁴, Graeme Jones¹. ¹University of Tasmania, Menzies Institute for Medical Research, Hobart, Australia; ²Australian Orthopaedic Association National Joint Replacement Registry (AOANJRR), Adelaide, Australia; ³South Australian Health and Medical Research Institute (SAHMRI), Adelaide, Australia; ⁴Monash University Medical School, Department of Epidemiology and Preventive Medicine, Melbourne, Australia

Background: Pain is the main impetus for osteoarthritis (OA) patients to seek healthcare including joint replacement. The pain experience is very heterogeneous and affected by factors across multiple domains—peripheral, psychological, and neurological, which suggests the existence of homogeneous subgroups/phenotypes within OA patients with pain. We recently identified three pain phenotypes using a wide spectrum of factors including main pain dimensions (structural damage on MRI, body mass index (BMI), comorbidities, psychological and multi-site pain (a surrogate of neurological factor)).

Objectives: To examine whether the risk of knee replacement (KR) varied when these three pain subgroups were compared.

Methods: 1099 participants (mean age 63 years; range 51–81 years) from a population-based cohort study were recruited at baseline. 875, 768 and 563 participants attended years 2.6, 5.1 and 10.7 follow-up, respectively. Demographic, psychological, lifestyle and comorbidities data were obtained at baseline. T1-weighted or T2-weighted MRI of the right knee was performed to measure knee structural pathology—cartilage defects, bone marrow lesions (BMLs) and effusion-synovitis at baseline. Knee pain was assessed using the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) pain questionnaire at each time-point. Presence of pain (yes/no) in the neck, back, hands, shoulders, hips, knees and feet was also assessed by questionnaire at each time-point. KR up to 13.7 years after recruitment was identified by linking to the Australian Orthopaedic Association National Joint Replacement Registry. Latent class analysis was used to differentiate pain phenotypes by considering sex, BMI, emotional problems, education level, comorbidities, number of painful sites and knee structural damage on MRI. Log-binomial regression was used to evaluate the association between three pain phenotypes and risk of KR.

Results: 963 participants were included in the analysis (BMI 27.7 kg/m², 50% female). Three distinct phenotypes were identified: Class 1: high prevalence of emotional problems and low prevalence of structural damage (25%); Class 2: high prevalence of structural damage and low prevalence of emotional problems (20%); Class 3: low prevalence of emotional problems and low prevalence of structural damage (55%). Participants in Class 1 had greater pain severity than those in Class 2 and 3. During a follow-up of 13.7 years, There were 46 right and 51 left TKR's in 79 participants. In multivariable analyses, participants in Class 1 and 2 had higher risk of requiring KR on the right [Class 1 vs. Class 3: relative risk (RR) 6.66, 95% confidence interval (CI) 2.17–20.49; Class 2 vs. Class 3: RR 14.14, 95%CI 4.85–41.22], left (Class 1 vs. Class 3: RR 2.75, 95%CI 1.22–6.19; Class 2 vs. Class 3: RR 5.46, 95%CI 2.64–11.32), and any knee joints (Class 1 vs. Class 3: RR 3.65, 95%CI 1.88–7.09; Class 2 vs. Class 3: RR 6.81, 95%CI 3.68, 12.60) compared to Class 3, but the associations were stronger with Class 2 than Class 1.

Conclusion: Participants with distinct pain phenotype groups have different risks of KR. This suggests that the identified phenotypes reflect distinct clinical subgroups with different prognoses. As expected the highest risk of KR was found in those with the most structural damage (Class 2). However, those participants with low structural damage have an increased risk of KR when they have high compared to low emotional problems, indicating that selection of subjects for KR could be improved by screening out those in Class 1.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6588

OP0180

LOW-DOSE PREDNISOLONE IN PATIENTS WITH HAND OSTEOARTHRITIS (HOPE): RESULTS FROM A RANDOMISED DOUBLE-BLIND PLACEBO-CONTROLLED TRIAL

Féline Kroon¹, Marion Kortekaas¹, Annelies Boonen^{2,3}, Stefan Böhringer⁴, Monique Reijnen⁵, Frits Rosendaal⁶, Naghmeh Riyazi⁷, Miriam Starmans⁸, Frankien Turkstra⁹, Jende van Zeven¹⁰, Cornelia Allaart¹, Margreet Kloppenburg^{1,6}. ¹LUMC, Rheumatology, Leiden, Netherlands; ²Maastricht University Medical Center, Rheumatology, Maastricht, Netherlands; ³Care and Public Health Research Institute, Maastricht, Netherlands; ⁴LUMC, Medical Statistics, Leiden, Netherlands; ⁵LUMC, Radiology, Leiden, Netherlands; ⁶LUMC, Clinical Epidemiology, Leiden, Netherlands; ⁷Haga Hospital, Rheumatology, The Hague, Netherlands; ⁸Zuyderland Medical Center, Rheumatology, Heerlen, Netherlands; ⁹Amsterdam rheumatology and immunology center | Reade, Amsterdam, Netherlands; ¹⁰Sint Franciscus Vlietland Group, Rheumatology, Rotterdam, Netherlands

Background: Hand osteoarthritis (OA) is a prevalent joint disease with high disease-burden in need for effective therapeutic options. Studies have shown that synovial inflammation is often present in hand OA and a main determinant of pain and radiographic disease progression.

Objectives: To investigate the efficacy and safety of short-term low-dose prednisolone in patients with painful hand OA.

Methods: This randomised, double-blind, placebo-controlled trial enrolled patients with painful hand OA, fulfilling American College of Rheumatology criteria, and signs of synovial inflammation. Patients with ≥ 4 interphalangeal joints (IPJ) with osteoarthritic nodes, ≥ 1 IPJ with soft swelling or erythema and ≥ 1 IPJ with positive power Doppler signal (PDS) or synovitis grade ≥ 2 on ultrasound, were eligible. Key exclusion criteria were chronic inflammatory rheumatic diseases, psoriasis, using immune modulating drugs within 90 days before baseline, and predominant thumb base pain. Eligible patients with visual analogue scale (VAS) finger pain ≥ 30 mm, flaring ≥ 20 mm upon non-steroidal anti-inflammatory drug washout, were randomised to receive prednisolone 10 mg daily for 6 weeks or placebo, followed by a two-week tapering scheme and 6 weeks without study medication. Outcomes were assessed at 2, 4, 6, 8 and 14 weeks. Primary endpoint was VAS finger pain at week 6 in intention-to-treat analysis. Secondary clinical endpoints included fulfilment of OMERACT-OARSI responder criteria, Australian/Canadian Hand OA Index (AUSCAN) pain/function, Functional Index for Hand OA (FIHOA), VAS patient global assessment, Short-Form 36 and grip strength. Imaging endpoints included ultrasound synovitis and PDS.

Table. Baseline characteristics and results of secondary endpoints at week 6 of HOPE study.

	Mean (SD) at baseline		Mean (SD) change from baseline		Adjusted mean between-group difference (95% CI)*	p-value
	Prednisolone	Placebo	Prednisolone	Placebo		
Pain						
AUSCAN pain, 0-20	11.3 (3.3)	10.2 (3.1)	-4.7 (3.5)	-1.1 (3.1)	-3.5 (-4.9 to -2.1)	<0.001
Function						
AUSCAN function, 0-36	18.6 (7.8)	19.0 (7.1)	-6.5 (7.4)	-2.7 (4.7)	-3.7 (-6.2 to -1.1)	0.01
FIHOA, 0-30	12.4 (5.4)	11.0 (4.7)	-2.6 (5.1)	-0.5 (4.0)	-2.1 (-4.0 to -0.2)	0.03
SF-36 physical component scale*	44.6 (7.9)	46.2 (6.8)	3.1 (6.7)	-0.3 (6.3)	3.5 (0.8 to 6.2)	0.01
VAS patient global assessment, 0-100	55.5 (21.7)	55.7 (22.0)	-23.6 (23.5)	-8.0 (25.3)	-15.4 (-25.6 to -5.2)	0.003
Grip strength, kg	20.4 (11.3)	20.4 (11.8)	3.5 (4.1)	2.2 (5.5)	1.2 (-0.8 to 3.2)	0.24
Ultrasound (sum scores)						
Synovitis, 0-90	16.4 (6.3)	17.8 (6.3)	-2.8 (4.7)	-0.3 (5.0)	-2.5 (-4.5 to -0.5)	0.02
Power Doppler Signal, 0-90	5.3 (4.1)	7.0 (4.3)	-1.7 (4.3)	-1.3 (4.2)	-0.4 (-2.2 to 1.4)	0.68

*Adjusted for baseline value and study center. *Norm-based scores with a standardized mean of 50 and SD of 10 using age- and sex-specific Dutch population-based norms. AUSCAN=Australian/Canadian Hand Osteoarthritis Index. FIHOA=Functional Index for Hand Osteoarthritis. SF-36=Short Form-36. VAS=visual analogue scale.

Results: Of 92 patients (mean (SD) age 63.9 (8.8), 79% women) randomised to prednisolone (n=46) or placebo (n=46), 42 patients in each group completed the study. Baseline characteristics were well-balanced between the groups. The mean (SD) change from baseline to week 6 in VAS finger pain was -21.5 (21.7) in the prednisolone and -5.2 (24.3) in the placebo group, with a mean between-group difference of -16.5 (95% confidence interval (CI) -26.1 to -6.9; figure). At week 6, 33 (72%) patients in the prednisolone versus 15 (33%) in the placebo group fulfilled OARSI responder criteria (odds ratio 5.3, 95% CI 2.0 to 13.6, p=0.001). In analogy with the primary endpoint, prednisolone was superior to placebo in most other secondary clinical endpoints (table). Ultrasound synovitis significantly improved at week 6 in the prednisolone compared to the placebo group, while no difference was observed in PDS (table). After tapering, between-group

differences disappeared. Adverse events were mostly mild and comparable between groups.

Conclusion: Six-week treatment with low-dose oral prednisolone led to a substantial improvement of symptoms in patients with painful hand OA and signs of inflammation. This trial provides evidence that local inflammation is a suitable target for drug-treatment in hand OA.

Acknowledgement: The HOPE study was supported by a grant from the Dutch arthritis foundation (ReumaNederland).

Disclosure of Interests: Féline Kroon: None declared, Marion Kortekaas: None declared, Annelies Boonen: None declared, Stefan Böhlinger: None declared, Monique Reijnders: Grant/research support from: Funding from the Dutch Arthritis Foundation.

The funding source had no role in the design and conduct of the study., Frits Rosendaal: None declared, Naghmeh Riyazi: None declared, Miriam Starmans: None declared, Frankien Turkstra: None declared, Jende van Zeben: None declared, Cornelia Allaart: None declared, Margreet Kloppenburg: Grant/research support from: Pfizer, IMI-APPROACH (Grant Agreement n° 115770), Consultant for: GlaxoSmithKline, Merck-Serono, Abbvie, Levicept, Pfizer

DOI: 10.1136/annrheumdis-2019-eular.997

THURSDAY, 13 JUNE 2019

Systemic sclerosis, myositis and related syndromes – etiology, pathogenesis and animal models

OP0181

MOLECULAR CHARACTERIZATION AND STRATIFICATION OF IDIOPATHIC INFLAMMATORY MYOPATHIES: ON THE BASIS OF SKELETAL MUSCLE TRANSCRIPTOME STUDY

Alfred Chen^{1,2}, Xin Lu², Jing-Zhe Shang^{3,4}, Ai-Ping Wu^{3,4}, Gen-Hong Cheng^{3,4}, Ning Lu³, Qinglin Peng², Guo-Chun Wang². ¹Graduate School of Peking Union Medical College, Beijing, China; ²China-Japan Friendship Hospital, Department of Rheumatology, Beijing, China; ³Center for Systems Medicine, Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China; ⁴Suzhou Institute of Systems Medicine, Suzhou, China

Background: Idiopathic inflammatory myopathies (IIM) are a group of complicated heterogenous autoimmune diseases, to date, little is known about its skeletal muscle transcriptomic features.

Objectives: Here, we performed skeletal muscle RNA-sequencing (RNA-seq) to discover the global muscle transcriptional signature of IIM based on myositis-specific antibodies (MSA) profiles and investigate the potential molecular pathway of IIM.

Methods: Muscle specimen were taken from 60 patients with IIM, 6 patients with non-IIM myopathies and 9 healthy controls. The serum and PBMC samples were also obtained from the IIM patients at the time of muscle biopsy. For RNA-seq, IIM was dissected into eight groups based on their MSA profiles: MSA and ANA negative (n = 4), anti-synthetase antibodies positive (n = 13), -MDA5 positive (n = 6), -NXP-2 positive (n = 7), -TIF-1γ positive (n = 10), -SRP or anti-HMGCR positive (n = 6), -Mi-2 positive (n = 7), MSA negative but anti-Ro-52 positive (n = 7). RNA from muscle specimen were extracted according to manufactory guide and sequenced using Illumina HiSeq2500. Quantitative real-time reverse transcription-polymerase chain reaction (qRT-PCR) was performed on sequencing cohorts and expanding cohorts to validate the results of RNA-seq. Immunohistochemistry was also performed on muscle biopsy to determine the MxA expression in different MSA subgroups.

Results: To define the global muscle signature of IIM, all IIM samples were compared to NC and total of 1637 transcripts were differentially expressed (log2 Fold Change > 1, Padj < 0.05). Unsupervised hierarchical clustering of these differentially expressed transcripts (DETs) revealed a prevalent interferon (IFN) signature and showed that 68 interferon-stimulated genes (ISGs) were significantly up-regulated in IIM. Then these 68 ISGs were used to cluster different MSA subgroups and distinct ISG expression was found. The mRNA expression levels of several ISGs (MX1, IFIH1, LAMP3, CMPK2, HERC6) in sequencing cohorts and expanding cohorts also confirmed the diverse ISG expression between different MSA subgroups. An IFN signature scoring system was established to quantify the IFN activity and subsequently IIM could be classified into IFN-Dominant, IFN-Moderate and IFN-Weak respectively based on the IFN intensity and different MSA subgroup. Moreover, the IFN-Dominant group showed much higher MxA expression on muscle biopsy than the IFN-Moderate and IFN-Weak group by immunohistochemistry.

Conclusion: We revealed a prominent IFN signature and MSA-based ISG expression heterogeneity in IIM through muscle transcriptomics. Preliminary

results showed that the IFN muscle signature may play a predominant role in some subgroups but not all IIM groups in the pathogenesis of IIM.

REFERENCES:

- [1] Miller FW, Lamb JA, Schmidt J, Nagaraju K. *Nat Rev Rheumatol*. 2018 Apr 20;14(5):255-268.
- [2] Mariampillai K, Granger B, Amelin D, Guiguet M, Hachulla E, Maurier F, Meyer A. *JAMA Neurol*. 2018 Dec 1;75(12):1528-1537.
- [3] Banchereau R, Hong S, Cantarel B, Baldwin N, Baisch J, Edens M, Cepika AM. *Cell*. 2016 Apr 21;165(3):551-65.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.854

OP0182

SYNTHETIC PEPTIDES TARGETING CD206 INHIBIT PATHOGENIC MACROPHAGES IN SYSTEMIC SCLEROSIS

Henry Lopez¹, Kimti Kumar¹, George Martin², Jesse Jaynes², James Stanway¹, Bahja Ahmed Abdi¹, Nikita Arumalla¹, Sarah Karrar¹, Sian Morris¹, Christopher Denton¹, David Abraham¹, Xu Shiwen¹, Richard Stratton¹. ¹UCL Division of Medicine Royal Free Campus, Centre for Rheumatology and Connective Tissue Diseases, London, United Kingdom; ²Riptide Bioscience, Vallejo, United States of America

Background: In systemic sclerosis, activated macrophages having M2-like properties are believed to contribute to the fibrotic pathology by secreting profibrotic factors such as TGFβ. Certain synthetic peptides (10-12mers) studied here coopt the mechanism used by macrophages to detect bacteria by binding to certain regions of CD206, a receptor highly expressed in the M2 macrophages, and repolarise them to an M1 phenotype.

Objectives: We aimed to determine the macrophage activation signature in SSc, assess CD206 as a biomarker of ongoing fibrotic activity, and measure the effect of the peptides on macrophage activation and macrophage-fibroblast cross-talk. Methods: sCD206 was assayed in plasma (n=50 healthy, 50 limited SSc, 50 diffuse SSc), and suction blister fluid (BF) obtained over active lesions by ELISA. sSIGLEC (monocyte interferon signature) and IL-31 (Th2 signature) were assayed for comparison. Cell surface CD206 and DAMP receptor P2X7, were assayed by flow cytometry. The macrophage activation signature was investigated further by qPCR for inflammatory M1, as well as M2-like gene expression (IFNγ, Arg1, CD2106). Macrophage-fibroblast cross talk was assessed using media transfer following heat activation to SSc dermal fibroblasts assayed by qPCR for collagen I. The macrophage secretome was determined by Luminex and ELISA.

Results: CD206 was significantly elevated in SSc BF (SSc median sCD206 42, HC 31 pg/ml, p<0.041). Plasma sCD206 was raised in diffuse versus limited SSc (P<0.0103) and was correlated with ESR (R=0.364, p<0.009) and sSIGLEC (R=0.244, p<0.05), but not disease duration or IL-31. By flow cytometry both CD206 and P2X7 were highly expressed by SSc macrophages (mean fluorescence SSc, 776.1 SD=409.1, 724.4 SD=455.3 vs HC 632.2 SD=73.7, 472.9 SD=25.4), correlating with modified Rodnan Skin Score (mRSS) (p<0.05, r=0.51). Double positive P2X7 and CD206 cells were seen in a subgroup with higher mRSS. Four RP peptides (RP182, 185, 832 & 837) were assessed for effect on growth of SSc and control macrophages. In general, the RP peptides selectively inhibited SSc macrophage growth, with RP185 and RP832 reaching significance (p<0.05) for 1 and 10μM concentrations at 96 hours. Pro-fibrotic cross-talk leading to elevated fibroblast collagen I, was characteristic of 4 out of 9 SSc macrophage cell lines examined. A mixed activation signature was identified by qPCR of the fibroblast-stimulating macrophages (elevated Arginase1, CD206 and IFNγ) but not the non-fibroblast inducing cell lines. Pre-treatment of the pro-fibrotic SSc macrophages with RP832, but not the others, eliminated the pro-fibrotic cross-talk (fibroblasts treated with SSc macrophage media mean collagen I, 1995 (SEM 149), RP832 treated macrophages 1551 (SEM 133) p<0.038 relative expression units, paired t test), and suppressed the macrophage activation signature (reduced mean Arg1 p<0.030, CD206 p<0.058, IFNγ p=0.058, paired t test). Secretome analysis confirmed the mixed activation signature of SSc macrophages including elevated total TGFβ, PDGF-BB, sCD40L as well as inflammatory factors IL-6, IL-1β, TNFα and type I cytokine IFNγ. Treatment with RP 832c reduced TGFβ in 6 out of 9 SSc macrophage cell lines but this did not reach statistical significance.

Conclusion: A macrophage activation signature is identified in SSc patients, in proportion to the severity of disease. A mixed activation signature with inflammatory as well as pro-fibrotic M2-like properties is seen. RP peptides, which target the cells via CD206, reduce the activation signature and inhibit pro-fibrotic cross talk in these cells.

Disclosure of Interests: : Henry Lopez Shareholder of: Murigenics, Riptide Bioscience, Employee of: Pfizer, Kimti Kumar: None declared, George Martin