Disclosure of Interests: None declared
DOI: 10.1136/annrheumdis-2019-eular.7489

AB0523 MENOPAUSE IN AN EGYPTIAN COHORT OF SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS: EFFECT OF THE DISEASE

Samah A. Elbakry, Safaa A. Hussein, Rasha M. Hamouda, Marina W. Naguib.
Internal Medicine Department, Division of Rheumatology, Ain Shams University, Cairo, Egypt

Background: Systemic lupus erythematosus (SLE) is a chronic systemic autoimmune disease that mainly affects females in the reproductive age¹. Although SLE generally emerges during reproductive ages, it was found that lupus patients experience menopause at younger age than the general population; however, whether the occurrence of menopause at a younger age in lupus patients results from the gonadotoxic effects of Cyclophosphamide treatment or from an autoimmune-mediated ovarian injury is debated².

Objectives: To identify menopause characteristics in an Egyptian cohort of women with SLE with effect of the disease on menopausal symptoms and the characteristics of disease activity and disease damage in perimenopausal and post-menopausal patients.

Methods: In this cross-sectional observational study, data of 120 consecutive SLE female patients who fulfilled the 2012 ACR/SLICC criteria³, above the age of 35. Disease activity was assessed by using the SLE disease activity index (SLEDAI)⁴, and accumulated damage was assessed by Systemic Lupus International Collaborative Clinics/American College of Rheumatology Damage Index (SLICC/ACR DI)⁵. Laboratory assessment was done to all patients including follicle stimulating hormone (FSH) and luteinizing hormone (LH) blood levels.

Results: The mean age of the patients was 45.067 ± 8.211 years and median disease duration was 5.074 ± 5.567 years (min 0.08 – max 21 years). It was found that the mean of total SLEDAI score was 4.667 ± 3.537 and mean of total SLICC/ACR damage index was 0.633 ± 0.819 . Of 120 patients 20% had premature menopause, 29.17% had natural menopause and 50.83% were still menstruating (table 1). It was found that the age at menopause (either natural or premature menopause) ranged from 26 to 54 years with the mean 45.170 ± 7.278 years. There was a statistically significant negative correlation between LH and total SLEDAI ($r = -0.178$, $P = 0.052$). It was also found a statistically significant positive correlation between FSH and cumulative Cyclophosphamide dose ($r = 0.440$, $P < 0.001$) and between LH and cumulative Cyclophosphamide dose ($r = 0.214$, $P = 0.046$) (table 2).

Table 1. Menstrual status:

	Positive	
	N	%
Premature menopause	24	20.00
Natural menopause	35	29.17
Menstruating	61	50.83

Conclusion: SLE has a reproductive and hormonal impact on female patients, either because of disease activity or due to pharmacological adverse effects. The mean age at menopause (either natural or premature menopause) is 45 years. High LH is associated with lower disease activity. High cumulative Cyclophosphamide dose is associated with high FSH and LH.

REFERENCES

- [1] Mills JA. Systemic lupus erythematosus. *New Engl J Med* 1994; 330: 1871–1879.
- [2] Sammaritano LR. Menopause in patients with autoimmune diseases. *Autoimmun Rev*. 2012; 11(6-7).
- [3] Petri M., Orbai AM., Alarcón GS, et al. Derivation and Validation of the Systemic Lupus International Collaborating Clinics Classification Criteria for Systemic Lupus Erythematosus. *Arthritis Rheum*. 2012; 64: 2677-86.
- [4] Bombardier C, Gladman D, Urowitz M, et al. Derivation of the SLEDAI. A disease activity index for lupus patients. The Committee on Prognosis Studies in SLE. *Arthritis Rheum* 1992; 35, 630–640.
- [5] Gladman D, Ginzler E, Goldsmith C, et al. The development and initial validation of the Systemic Lupus International Collaborating Clinics/American College of Rheumatology damage index for systemic lupus erythematosus. *Arthritis Rheum*. 1996; 39: 363-369.

Table 2. Correlation study between FSH & LH and Cyclophosphamide:

Correlations	FSH (mIU/ml)		LH (mIU/ml)	
	P-value	R	P-value	R
Cumulative Cyclophosphamide (mg)	0.440	< 0.001*	0.214	0.046*

Disclosure of Interests: None declared
DOI: 10.1136/annrheumdis-2019-eular.1226

AB0524 SERUM BAFF AND APRIL AS CANDIDATE BIOMARKERS IN SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): A PROSPECTIVE FOLLOW-UP STUDY

Selma Sarı¹, Suzan Çınar², Bahar Artım-Esen³, Ahmet Gül³, Mahdume Lale Ocal³, Gunnur Deniz², Murat İnanc³. ¹Istanbul University, Istanbul Faculty of Medicine, Internal Medicine, Istanbul, Turkey; ²Istanbul University, Aziz Sancar Institute of Experimental Medicine, Immunology, Istanbul, Turkey; ³Istanbul University, Istanbul Faculty of Medicine, Rheumatology, Istanbul, Turkey

Background: BAFF and APRIL are cytokines involved in B cell development and they take place in the pathogenesis of SLE.

Objectives: The aim of this study was to investigate the relationship between serum BAFF/APRIL levels with clinical features and disease activity in SLE patients.

Methods: We included 79 patients with SLE (ACR criteria) and 27 healthy controls into the study. Serum BAFF and APRIL levels were assessed by ELISA. In 19 patients with active disease at the time of the assessment BAFF/APRIL levels were reassessed at least 6 months later (mean 7.8 months) and disease activity was evaluated by SLEDAI. New renal involvement was observed in 16 patients during the study and renal involvement was previously detected in 12 patients.

Results: Although both BAFF (median 0.7 vs 0.41 ng/ml) and APRIL (median 2.3 vs 1.05 ng/ml) levels were higher in patients with SLE compared to the control group ($p < 0.001$), no correlation was found between BAFF/APRIL levels and SLEDAI scores. Only 9 patients (11.4%) had both BAFF and APRIL in normal range (95% confidence interval). When patients were grouped according to disease activity as no activity (SLEDAI = 0), low disease activity and active disease, there was no difference in BAFF/APRIL levels between groups. Serum BAFF levels were higher in patients with renal disease activity (median 0.94 ng/ml vs 0.61 ng/ml, $p=0.01$), and there was a positive correlation between APRIL levels and proteinuria ($r=0.42$, $p=0.02$). There was no association between BAFF/APRIL levels and anti-dsDNA positivity but a weak inverse correlation was observed between BAFF and C3 levels ($r=0.25$, $P=0.02$). No correlation was found between BAFF/APRIL levels and renal SLEDAI score, renal histopathology activity and chronicity index scores. In the active disease group after follow-up, there was no significant change in BAFF (from 1.63 ng/ml to 1.2 ng/ml) and APRIL levels (from 2.11 ng/ml to 2.31 ng/ml).

Conclusion: BAFF/APRIL levels were found to be significantly higher in patients with SLE compared to controls, but no association with disease activity was found. BAFF levels are correlated with decreased C3 levels. These results suggest that both cytokines are involved in the pathogenesis of SLE, and that serum BAFF and APRIL levels can be valuable as a biomarker in SLE especially in patients with renal activity. Long-term studies on the effect of treatment are needed.

Disclosure of Interests: None declared
DOI: 10.1136/annrheumdis-2019-eular.4671

AB0525 ANTI-SMITH ANTIBODY IS ASSOCIATED WITH DISEASE ACTIVITY IN PATIENTS WITH NEW-ONSET SYSTEMIC LUPUS ERYTHEMATOSUS

Seungmin Jung, Juyoung Yoo, Sungsoo Ahn, Sangwon Lee, Jason Jungsik Song, Yongbeom Park. *Yonsei University College of Medicine, Internal Medicine, Seoul, Korea, Rep. of (South Korea)*

Background: Anti-Smith (Sm) antibody is a highly specific antibody for systemic lupus erythematosus (SLE). Despite the remarkable specificity of anti-Sm antibodies for SLE, the association between anti-Sm antibody level and the clinical manifestation of SLE is still unclear.

Objectives: We aimed to evaluate the association between anti-Sm antibodies and disease activity in patients with new-onset SLE.