Figure 1.

Conclusion: AQUILA study revealed high disease burden at BL across AS and PsA pts characterized by the presence of extra-articular manifestations and cardiovascular diseases. Adherence rates under real-world conditions were consistent and comparable with published literature regarding immune-mediated inflammatory diseases³. SEC demonstrated a high drug persistence in real world in particular in Bx-naïve pts. Overall, this interim analysis demonstrated that SEC is a reliable treatment in Bx-naïve as well as Bx-pretreated pts with AS or PsA in a daily routine setting.

REFERENCES

- [1] Baeten D et al. *Lancet*. 2013;382(9906):1705-13;
- [2] McInnes IB et al. *Lancet*. 2015;386(9999):1137-46;
- [3] Vangeli E et al. *Adv Ther*. 2015;32:983-1028

Disclosure of Interests: Uta Kiltz Grant/research support from: AbbVie, Chugai, Eli Lilly, Grünenthal, Janssen, MSD, Novartis, Pfizer, Roche, and UCB., Consultant for: AbbVie, Chugai, Eli Lilly, Grünenthal, Janssen, MSD, Novartis, Pfizer, Roche, and UCB. Daniel Peterlik Employee of: Novartis Pharma GmbH, Veronika Winkelmann Employee of: Novartis Pharma GmbH, Hans-Peter Tony Consultant for: Eli Lilly and Company, Speakers bureau: Eli Lilly and Company
DOI: 10.1136/annrheumdis-2019-eular.435

AB0706

DIFFERENCES IN RHEUMATOLOGISTS' TREATMENT PREFERENCES FOR ANKYLOSING SPONDYLITIS AND NON- RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS

Phil pouliot, Lynn Price. *Spherix Global Insights, Exton, United States of America*

Background: The lack of a formal indication for Non-Radiographic Axial Spondyloarthritis (nr-axSpA) has meant that many US rheumatologists fundamentally treat it as an early manifestation of Ankylosing Spondylitis (AS) and seldom differentiate their treatment approach when using biologics/small molecules.

Objectives: This study sought to gain a better understanding of the number of AS and nr-axSpA patients rheumatologists manage, recent changes made in treatment, and how AS and nr-axSpA patients are differentiated beyond radiographic evidence. The study also evaluated the differences in use of biologics/small molecules for the treatment of AS and nr-axSpA and rheumatologist projected changes in the management of these two conditions.

Methods: An independent market analytics firm administered a survey with US rheumatologists (n=104) on the treatment landscape for AS and nr-axSpA. To qualify, rheumatologists had to spend the majority of their professional time in clinical practice, manage a minimum of 50 AS or nr-axSpA patients, and have been in practice between 2 and 35 years. Respondents answered questions pertaining to their practice, the demographics of their AS/nr-axSpA patients, their treatment approaches for each type, as well as attitudinal statements. The survey was administered online in February 2018 and respondents were compensated for their participation. A similar study was also conducted in 2017 allowing for year over year comparison. The data were collected and analyzed in SPSS.

Results: Three-quarters of rheumatologists reported they treated their AS and nr-axSpA patients in a similar manner, though there was a higher use of NSAIDs and local steroids in non-radiographic disease, and more biologic/JAK use in AS. More established TNF inhibitors such as adalimumab, etanercept, and infliximab dominated biologics/small molecule preference, collectively accounting for approximately three-quarters of all biologic-treated AS and nr-axSpA patients. However, use of adalimumab, etanercept, and infliximab was expected to decline or stay flat in the next three months, while use of secukinumab, tofacitinib, golimumab IV, and both infliximab biosimilars, was anticipated to increase.

Although not indicated for either condition, 28% of respondents reported off-label use of tofacitinib for AS, and 25% for nr-axSpA, a significant increase from 2017. Projected off-label use of tofacitinib in the following three months is also expected to increase.

Conclusion: Rheumatologists are less likely to use biologics/small molecules for nr-axSpA, but when they do, use of brands is similar to those used in AS. While TNF inhibitors continue to account for the majority of biologic/small molecule prescriptions for these conditions, rheumatologists project increased use of agents with alternative mechanisms of action in the future.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7752

AB0707

LTBI SCREENING IN SPONDYLOARTHRITIS PATIENTS PRIOR TO ANTI-TNF TREATMENT AND FOLLOW-UP IN AN ENDEMIC AREA

Andrea Shimabuco, Ana Medeiros, Renata Miossi, Karina Bonfiglioli, Julio Moraes, Celio Gonçalves, Percival D. Sampaio-Barros, Claudia Goldenstein-Schainberg, Fernando Souza, Leandro Prado, Michelle Lopes, Eloisa Bonfa, Carla Saad.
Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Rheumatology Division, Sao Paulo, Brazil

Background: Screening for latent tuberculosis infection (LTBI) remains a concern in endemic regions for anti-TNF eligible patients and recent researches have shown peculiarities between the inflammatory arthropathies such as different groups of spondyloarthritis (SpA).

Objectives: To evaluate the long-term efficacy of LTBI screening and treatment in patients with Ankylosing Spondylitis (AS) and Psoriatic Arthritis (PsA) receiving TNF blockers in a single center.

Methods: A total of 218 SpA patients (135 AS and 83 APS) were screened for LTBI before receiving anti-TNF (infliximab, adalimumab, etanercept, certolizumab pegol and golimumab) treatment using the tuberculin skin test (TST), chest X-ray (CXR) and history of previous exposure to tuberculosis (TB). Patients were regularly followed every 2-3 months and asked about infectious symptoms or new exposure. TST was not repeated regularly. LTBI patients were treated with isoniazid (300 mg/day) for 6 months, according local guidelines.

Results: One hundred and eight patients (49.5%) were treated with a single anti-TNF agent and the total duration of biological treatment was approximately six years (5.9 ± 4.0). INF and ADA were most often used among the 422 treatment cycles analyzed, representing 45% and 30% respectively. LTBI screening was positive in 82 patients (38%): 69 (84%) were TST-positive, 23(28%) had a history of TB exposure and 5(6%) had an abnormal CXR. As isolated variables TST positivity and previous exposure accounted for 58(71%) and 11 (13%) LTBI diagnosis. There were some distinct patterns between APS and AS patients screening: despite APS patients had more cases of previous TB than AS patients (6% vs 0.7%, $P = 0.03$), they had a lower frequency of LTBI (30% vs 42%, $P = 0.04$). Among LTBI patients, TST positivity was lower in SpA than AS patients (64% vs 93%, $P = 0.002$), even with more previous exposure (52% vs 18%, $P = 0.02$) and in patients with peripheral arthritis (27% vs 42%, $P = 0.03$). During follow-up, 11 patients developed active TB: 5 under ADA, 5 under INF and 1 under ETA treatment. Five cases (45%) were extrapulmonary: 3 pleural, 1 peritoneal and 1 spondylodiscitis. Four (36%) cases occurred in patients with a positive LTBI screening and 7 in patients without LTBI. There was no difference in drug survival

according to type or class of anti-TNF, disease subgroups, duration or use of synthetic drugs and prednisone. Four (36.3%) cases occurred in the first year, median 5.3 (1.2-8.8) months after initiating anti-TNF exposure, 2 of them (50%) in patients with positive LTBI. Seven cases were probably due to re-exposure since occurred later, median 21.9 (14.2-42.8) months (5 in patients with negative LTBI screening). Six patients (54.5%) re-initiated treatment with ETA. Only the patient who developed pulmonary TB under ETA had a second TB infection after 18 months of therapy.

Conclusion: Despite the adequate screening and treatment of LTBI, according to local guidelines, TB still occurs in spondyloarthritis patients under anti-TNF therapy, even in the first year of treatment. These data point to LTBI screening/treatment failure, maybe due to anergy, mainly in PsA patients, with peripheral disease, low adherence or re-exposition in an endemic environment. The high frequency of extrapulmonary disease is also a diagnostic challenge.

REFERENCES

- [1] Cantini F, Nannini C, Niccoli L, et al; SAFEPIO (Italian multidisciplinary task force for screening of tuberculosis before and during biologic therapy). Guidance for the management of patients with latent tuberculosis infection requiring biologic therapy in rheumatology and dermatology clinical practice. *Autoimmun Rev*. 2015 Jun;14(6):503-9.

Disclosure of Interests: Andrea Shimabuco: None declared, Ana Medeiros: None declared, Renata Miossi: None declared, Karina Bonfiglioli Speakers bureau: Has received speaking fees from Roche, Pfizer, Bristol-Myers Squibb, Abbvie and Janssen, Julio Moraes: None declared, Celio Gonçalves: None declared, Percival D. Sampaio-Barros: None declared, Claudia Goldenstein-Schainberg: None declared, Fernando Souza: None declared, Leandro Prado: None declared, Michelle Lopes: None declared, Eloisa Bonfa: None declared, Carla Saad: None declared

DOI: 10.1136/annrheumdis-2019-eular.7062

AB0708

INFLUENCE OF IMMUNOGENICITY ON LONG-TERM MAINTENANCE OF ADALIMUMAB IN SPONDYLOARTHRITIS

Marine Samain^{1,2}, Emilie Ducourau^{1,2,3}, Theo Rispens⁴, Emmanuelle Derris⁵, Fabienne Le Guilhard⁶, Lucia Andras⁶, Aleth Perdriger⁷, Eric Lespessailles³, Antoine Martin⁸, Grégoire Cormier⁹, Thomas Armingeat¹⁰, Valerie Devauchelle-Pensec^{11,12}, Elisabeth Gervais¹³, Benoit Le Goff¹⁴, Annick de Vries¹⁵, Eric Piver^{16,17}, Gilles Paintaud^{2,18}, Céline Desvignes^{2,18}, David Ternant^{2,18}, Hervé Watier^{2,19}, ^{1,2}Philippe Goupille²⁰, Denis Mulleman^{1,2}. This work was a collaborative venture by the VICTOR HUGO network (Hôpitaux Universitaires du Grand Ouest—Western France University Hospitals; <http://www.srouest.fr>), dedicated to innovative research on rheumatic diseases. ¹CHRU de Tours, Service de Rhumatologie, Tours, France; ²University of Tours, EA 7501 GICC, Tours, France; ³CHU d'Orléans, Service de Rhumatologie, Orléans, France; ⁴Landsteiner laboratories, Sanquin Research, Amsterdam, Netherlands; ⁵CHU de Mans, Service de Rhumatologie, Le Mans, France; ⁶CHU de Blois, Service de Rhumatologie, Blois, France; ⁷CHRU de Rennes, Service de Rhumatologie, Rennes, France; ⁸CHU de Saint-Brieuc, Service de Rhumatologie, Saint-Brieuc, France; ⁹CHU Vendée, Service de Rhumatologie, La Roche-sur-Yon, France; ¹⁰CHU de Saint-Nazaire, Service de Rhumatologie, Saint-Nazaire, France; ¹¹Université de Brest, Inserm UMR1227 LBAI, Brest, France; ¹²CHRU de Brest, Service de Rhumatologie, Brest, France; ¹³CHRU de Poitiers, Service de Rhumatologie, Poitiers, France; ¹⁴CHRU de Nantes, Service de Rhumatologie, Nantes, France; ¹⁵Biologicals Lab, Sanquin Diagnostic Services, Amsterdam, Netherlands; ¹⁶University of Tours, Inserm U 1259, Tours, France; ¹⁷CHRU de Tours, Laboratoire de Biochimie, Tours, France; ¹⁸CHRU de Tours, Laboratoire de Pharmacologie-Toxicologie, Tours, France; ¹⁹CHRU de Tours, Laboratoire d'Immunologie, Tours, France; ²⁰CHRU de Tours, Inserm CIC1415, Tours, France

Background: Immunogenicity of anti TNF monoclonal antibodies leads to poor or secondary loss of response. Methotrexate reduces anti-drug antibodies (ADA) to adalimumab at week 26 in spondyloarthritis (SpA).¹

Objectives: Herein we sought to examine adalimumab long term persistence in ADA positive versus ADA negative SpA patients.

Methods: The CoMARIS study (Combination of Methotrexate and Adalimumab to Reduce Immunization in patients with axial SpA) is a 26-week prospective, randomised, open-labelled, multicentre study in which patients received adalimumab 40 mg subcutaneously (s.c.) every other week either in combination with MTX 10 mg s.c. for 26 weeks or without MTX. In a post hoc analysis, we reviewed the charts of patients to

assess adalimumab persistence. A Cox model analysis was performed to test the following covariates: MTX combination or not, sex, presence of ADA at week 26.

Results:

Data from 104 patients (54 without MTX and 50 with MTX) were reviewed, and time of adalimumab discontinuation was collected. The median time of follow-up was 210.57 weeks. ADA positivity at week 26 was the only covariate associated with adalimumab persistence. The median retention rate of adalimumab in ADA positive patients was 56.9 weeks, as compared with 98.6 weeks in those without ADA (log rank: $p=0.015$). In the Cox model analysis, the presence of ADA at week 26 increased the risk of adalimumab discontinuation by 1.78 [IC 95%=1.11-2.85], $p=0.016$.

Adalimumab retention rates in function of ADA status at 26 weeks

Figure. adalimumab retention rate in ADA positive patients (n=38) at week 26 versus ADA negative patients (n=66), log rank: $p=0.015$

Conclusion: Immunogenicity is a key factor that contributes to adalimumab discontinuation in SpA. MTX at initiation may therefore be considered in combination to adalimumab in SpA patients.

REFERENCES

- [1] Ducourau E, et al. Methotrexate reduces adalimumab immunogenicity in patients with spondyloarthritis: a randomized clinical trial. *EULAR17-1527*.

Acknowledgement: This work was promoted by the Regional University Hospital Center of Tours and supported by grants from the French Ministry for Health and Sport within the framework of the Programme Hospitalier de Recherche Clinique2011.

Disclosure of Interests: Marine Samain: None declared, Emilie Ducourau Speakers bureau: BMS and Abbvie, Theo Rispens Grant/research support from: Genmab, Speakers bureau: Pfizer, Abbvie, Regeneron, Emmanuelle Derris: None declared, Fabienne Le Guilhard: None declared, Lucia Andras: None declared, Aleth Perdriger: None declared, Eric Lespessailles Grant/research support from: Grants/research support from Amgen, Eli Lilly, MSD, UCB., Consultant for: Consultant for Amgen, Expanscience, Eli Lilly, MSD, UCB., Antoine Martin: None declared, Grégoire Cormier: None declared, Thomas Armingeat: None declared, Valerie Devauchelle-Pensec Grant/research support from: Roche-Chugai, Speakers bureau: MSD, BMS, UCB, Roche, Elisabeth Gervais Speakers bureau: Abbvie, BMS, MSD, Pfizer, Roche, UCB, Novartis, Benoit Le Goff Speakers bureau: Abbvie, BMS, Janssen, MSD, Pfizer, Sanofi-Genzyme, UCB, Novartis, Annick de Vries: None declared, Eric Piver: None declared, Gilles Paintaud Grant/research support from: Novartis, Roche Pharma, Sanofi-Genzyme, Chugai, Pfizer and Shire, Céline Desvignes: None declared, David Ternant Speakers bureau: Sanofi, Amgen, Hervé Watier: None declared, Philippe Goupille Grant/research support from: Financial compensation received from MSD on a pro-rotas basis for participation in Scientific Committee meetings and functions for this study, Speakers bureau: Abbvie, Biogaran, BMS, Hospira, Janssen, MSD, Pfizer, Sanofi-Genzyme, UCB, Denis Mulleman Speakers bureau: Pfizer, Novartis, Grifols

DOI: 10.1136/annrheumdis-2019-eular.4698