0.0001 respectively). Patients with falls had lower BBS ($p=0.03$) and loss of balance correlated with impairment of the quality of life ($\rho=-0.56$; $p<0.001$).

Conclusion: Balance is impaired in 1/3 of SpA patients and the BBS is associated mainly with functional and metrology indexes, showing that patients with severe cumulative damage are more affected.

REFERENCES

- [1] - Pompeu JE, Romano RS, Pompeu SM, Lima SM. Static and dynamic balance in subjects with ankylosing spondylitis: literature review. *Rev Bras Reumatol.* 2012; 52:409-16.
- [2] - Sawacha Z, Carraro E, Del Din S, Guiotto A, Bonaldo L, Punzi L, et al. Biomechanical assessment of balance and posture in subjects with ankylosing spondylitis. *J. Neuroeng Rehabil.* 2012; 9:63.

TABLE 1. - Study of BBS (balance berg scale) values according to epidemiological, clinical and treatment variables

	Median BBS with the variable (IQR)	Median BBS without the variable (IQR)	p
Male gender	51(44.0-55.0)	51.0(44.2-53.7)	0.80
White ethnic background	49.0(42.5-53.0)	54.0(30.2-55.0)	0.01
Exposed to tobacco (ex and current)	46.0(41.5-54.0)	52.0(46.5-55.0)	0.03
Bilateral sacroiliitis	51.0(45.0-55.0)	53.5(46.2-55.0)	0.63
Enthesitis	50.0(38.5-55.2)	51.0(44.5-55.0)	0.45
Dactylitis	50.5(42.0-54.2)	51.0(44.0-55.0)	0.73
Uveitis	51.0(46.5-55.0)	50.5(41.7-55.0)	0.26
Peripheral arthritis	52.0(40.7-55.7)	51.0(33.0-54.0)	0.92
HLA B27 presence	52.0(44.5-55.0)	48.5(39.5-51.1)	0.03
Anti TNF- α users	50.5(45.0-54.7)	52.0(39.0-55.0)	0.67
History of falls in the last year	46.5(40.0-50.2)	52.0(44.5-55.0)	0.03

TABLE 2. - Correlation studies of BBS (berg balance scale) values with age, disease duration, disease activity indexes, functional and metrology indexes and quality of life

	Rho	95% confidence interval	p
BASDAI	-0.28	-0.51 to -0.008	0.03
ASDAS VHS	-0.32	-0.54 to -0.05	0.01
ASDAS CRP	-0.33	-0.55 to -0.06	0.01
BASFI	-0.71	-0.82 to -0.55	<0.0001
BASMI	-0.80	-0.88 to -0.67	<0.0001
ASQoL	-0.56	-0.72 to -0.35	<0.0001
Age (years)	-0.50	-0.68 to -0.25	<0.0001
Disease duration (years)	-0.10	-0.36 to 0.17	0.44

BASDAI=Bath Ankylosing Spondylitis Disease Activity Index; ASDAS=Ankylosing Spondylitis Disease Activity Score, ESR=erythrocyte sedimentation rate; CRP=C reactive protein; BASMI=Bath Ankylosing Spondylitis Metrology Index; BASFI=Bath Ankylosing Spondylitis Functional Index; ASQoL=Ankylosing Spondylitis Quality of Life questionnaire.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5970

AB0715 ASSESSMENT OF BODY COMPOSITION IN LEAN MASS AND FAT MASS IN SPONDYLOARTHRITIS

Dhia Kaffel, selma.bouden,Kaouther Maatallah, Nahaline Abaza, Hend Riahi, Wafa Hamdi, Mohamed Montacer Kchir. *kassab institute, tunis, Tunisia*

Background: Mohamed Montacer Kchir bsorptiometry Absorptiometer (DEXA) is the gold standard for the study of body composition. It allows to sepMohamed Montacer Kchirarate the 2 compartments lean mass (LM) and fat mass (FM), and to study their distribution at the android and gynoid level

Objectives:

- Evaluate the change in total body composition measured the DEXA in patients with spondyloarthritis (SpA) compared to a control group.
- Evaluate the distribution of LM/FM at the android and gynoid levels.
- Study the influence of disease activity and bone mass on body composition.

Methods: This was a case-control study involving 50 patients with SpA compared to 50 controls. Each control was comparable in age and sexe to the matched patient.

Results: The mean age was 36 years in the SpA group and 35,5 years in the control group. Eighty per cent of the patients with SpA were male. The mean BASDAI was $4,64 \pm 2,5$. The mean ESR rate was

$32,37 \pm 20,95$, the mean CRP rate was $18,05 \pm 27,22$ [0-141]. The mean body mass index (BMI) was $25,4 \text{ Kg/m}^2$ ($25,74 \text{ Kg/m}^2$ in the control group). The mean bone mineral density (BMD) at the lumbar spine was 1.033 g/cm^2 (with a mean T score of -1.5 DS and Z score -1.3 DS) while it was 0.956 g/cm^2 at upper end of the femur level (with a mean T score of -0.8 and a mean Z score of -1.2 DS). In controls, the mean BMD was 1.055 g/cm^2 and 1.013 g/cm^2 at the lumbar spine and upper end of the femur respectively (with a mean T score of -1.4 DS and -0.5 SD and a mean Z score of -1.4 and -0.3 respectively at the lumbar and superior femoral sites). In the SpA group, the mean value of whole body FM was 31.6%, whereas this average was around 36% in controls. FM was distributed as follows: 39.2% at the android level versus 37.7% at the gynoid level. For LM, the whole body average was 68.4%, while the average was 60.8% at the android level and 62.3% at the gynoid level. The 2 groups (patients and controls) were comparable in terms of BMD, T score and Z score measurements at different lumbar and superior femoral sites. The SpA group and their healthy controls were comparable in terms of FM and LM measurements and their distributions. We found a significant correlation between the rate of CRP and the measurements of FM at the android level ($r = -0.358$, $p = 0.023$) (and conversely the LM at the android level). However, we found no association between disease activity (BASDAI) and body composition.

Conclusion: Changes in body composition (MM/MG) related to chronic inflammatory status may occur during chronic inflammatory diseases. However, few studies in the world have focused on changes in the bodily compartments and their distribution in SpA.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5450

AB0716

ESTABLISHMENT OF BASDAI CUT-OFFS FOR THE DISEASE ACTIVITY STATES BASED ON ASDAS CUT-OFFS IN TAIWANESE ANKYLOSING SPONDYLITIS PATIENTS

Hsin-Hua Chen¹, Ting-I Pan², Yi-Ming Chen¹, Yi-Hsing Chen¹, Wen-Nan Huang¹, Tsu-Yi Hsieh¹, Kuo-Lung Lai¹, Wei-Ting Hung¹, Yin-Yi Chou¹, Chih-Wei Tseng¹, Yi-Da Wu¹, Chia-Wei Hsieh¹, Ching-Tsai Lin¹. ¹Taichung Veterans General Hospital, Division of Allergy, Immunology and Rheumatology, Department of Internal Medicine, Taichung, Taiwan, Republic of China; ²Tunghai University, Department of Industrial Engineering and Enterprise Information, Taichung, Taiwan, Republic of China

Background: The Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) has been widely utilized to evaluate disease activity in patients with ankylosing spondylitis (AS). However, the cut-off of BASDAI used to indicate high disease activity (i.e., ≥ 4) was determined arbitrarily and was suggested as a criterion to initiate biological therapy for AS patients. The Ankylosing Spondylitis Disease Activity Score (ASDAS) has been developed as a new composite index to assess AS disease activity. The cut-off values for disease activity states has been defined and validated. ASDAS ≥ 2.1 was selected as a criterion of starting biological therapy. However, the BASDAI cut-off values corresponding to the ASDAS cut-off values for disease activity states were unknown.

Objectives: The purpose of this study was to estimate the corresponding BASDAI and ASDAS cut-off in a Taiwanese AS cohort.

Methods: Since November 2016, we assessed the Ankylosing Spondylitis Disease Activity Score (ASDAS) and the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) regularly and recorded demographic data, comorbidity, family history, medication use for AS patients in Taichung Veterans General hospital (TCVGH) using an electronic patient reported data system linked to an electronic medical record system. We identified 489 AS patients with complete baseline demographic and assessment data from the TCVGH electronic data system during 2016/11–2018/10. We used receiver operating characteristic (ROC) curves with Youden's J statistic to determine the cut-off values of BASDAI that correspond to ASDAS disease activity cut-offs (i.e., 1.3, 2.1 and 3.5).

Results: We included a total of 489 AS patients [114 (23.3%) females, mean age 44.1 years (S.D. 13.9), mean symptom duration 18.0 years (S.D. 11.9), 152 (31.1%) current biologic users]. Mean BASDAI, ASDAS-ESR and ASDAS-CRP scores were 2.1 (S.D. 1.5), 1.6 (S.D. 0.8) and 1.5 (S.D. 0.9) respectively. Mean levels of CRP and ESR were 0.6 (S.D. 1.5) mg/dl and 12.2 (S.D. 14.0) mm/hr respectively. Based on ASDAS-CRP, the numbers (%) of AS patients with inactive disease (<1.3), low disease activity (1.3-2.1), high disease activity (2.1-3.5) and very high disease activity (≥ 3.5) were 210 (42.9), 171 (35.0), 88 (18.0) and 20 (4.1) respectively. Based on ASDAS-ESR, the numbers (%) of AS patients with inactive disease, low disease activity, high disease activity and very high disease activity were 202 (41.3), 174 (35.6), 96 (19.6) and 17 (3.5)