Table 1.

	CLINICAL TRIAL Gladman D, et al. (N=131)	CLINICAL PRACTICE (N=41)
Baseline demographic parameters		
Age, years (mean±SD)	49.5±12.3	50.2±10.7
Sex, n (%)	67M/64F (51/49)	18M/23F (46.6/53.4)
Disease Characteristics		
Duration of psoriatic arthritis, year (mean±SD)	9.6±7.6	12.6±9.1
HAQ-DI	1.3±0.7	1.9±0.4 (n=22)
Swollen joint count, mean±SD	12.1±10.6	5.7±5.3
Painful joint count, mean±SD	20.5±13.0	8.7±6.1
Elevated CRP, n (%)	85 (65)	33 (80.5)
PASI score, median [IQR]	7.6 [0.6–32.2]	6 [0–15]
Day 1 oral glucocorticoid use, n (%)	37 (28)	25 (60.9)
Concomitant synthetic DMARD, n (%)		
- Methotrexate	98 (75)	25 (60.9)
- Leflunomide	12 (9)	8 (19.5)
- Sulfasalazine	21 (16)	0 (0)
- Others	2 (2)	0 (0)
IL of previous Anti-TNF inhibitors, mean±SD	1.7±1.0	2.5±1.4
IL of previous other biological no anti-TNF, n (%)	11 (8)	52 (126.8)
Adverse event, n (%)	72 (55)	10 (24.4)
Discontinuation due to adverse event, n (%)	2 (2)	1 (2.4)

Table 2.

	Baseline (N=41)	1st month (N=41)	6th month (N=13)
Prednisone, n (%)	25 (60.9%)	8 (43.9%)	6 (46%)
Dose (mean±SD)	4.9±6.1	7.6±4.7	5.6±2.9
Methotrexate, n (%)	9 (21.9%)	6 (14.6%)	3 (23%)
Dose (mean±SE)	18.8±5.8	19.5±3.3	21.6±2.8
N. Patients: Skin involvement, n (%)	22 (53.6%)	8 (19.5%)	2 (15.4%)
N. Patients: Nail involvement, n (%)	12 (29.2%)	7 (17.1%)	6 (46.2%)
N. Patients: arthritis, n (%)	38 (92.6%)	8 (19.5%)	2 (15.4%)
- Swollen joint count, median [IQR]	4 [2–7]	2 [1–4.3]	0 [0–1.5]
- Painful joint count, median [IQR]	7 [4.3–11.5]	2 [1–5]	1 [0–2]
N. Patients: enthesitis, n (%)	11 (26.8)	8 (19.5)	1 (7.7%)
N. Patients: dactylitis, n (%)	7 (17.1)	4 (9.7)	ND
CRP mg/dl, median [IQR]	2.36 [0.95–7.5]	1.7 [0.5–4.5]	0.98 [0.4–5]
DAS28, median [IQR]	5.1 [4.7–5.7]	3.8 [3.2–4.6]	2.9 [2.4–3.3]
DAPSA, median [IQR]	28.3 [23.2–32.3]	14 [7.2–31.1]	7.6 [4.8–19.2]
PASI, median [IQR]	6 [0–15]	5 [0.7–9.5]	2 [0–8]

Prior TOFA, most patients had received oral prednisone or equivalent (max. 17.9±13.6 mg/d), synthetic immunosuppressants (mean 1.9±0.7) and biological therapy (TB) (3.4±1.9). TB were as follows: etanercept (29), adalimumab (29), Infliximab (16), golimumab (13), certolizumab (14), secukinumab (30), ustekinumab (22). Apremilast was used in 10.

After a mean follow-up of 12.6±9.1 years after the PsA diagnosis, TOFA was started (5 mg/12 h). 60.9% associated prednisone (9.4±6.1 mg/d). In 17 (41.5%) TOFA was started in combined therapy: methotrexate (9) and leflunomide (8); in the remaining 24, monotherapy was prescribed. In addition to active arthritis patients presented skin involvement (53.6%), enthesitis (26.8%), nail involvement (29.2%) and dactylitis (17.1%).

Patients of clinical practice compared with clinical trial have a longer duration of PsA, functional disability (HAQ) and received a higher proportion of corticosteroids and TB (anti-TNF and non-anti-TNF) (table 1).

After a median follow-up of 4 [3–10.5] months, patients showed prompt improvement in the main variables, both in activity indexes (PASI, DAS28, DAPSA) and laboratory test (table 2). Minor side effects were reported in 10 patients (gastrointestinal symptoms), and TOFA was discontinued in 1 due to persistent symptoms.

Conclusion: In this preliminary study, first patients of clinical practice in Spain with TOFA in PsA had a longer evolution and received a greater number of TB than those of clinical trial. As in the TOFA clinical trial, it seems effective, rapid and relatively safe in daily clinical practice for refractory PsA.

REFERENCES

- https://www.ema.europa.eu/documents/overview/xeljanz-epar-medicine-overview_es.pdf
- Gladman D. N Engl J Med 2017; 377: 1525-36.

Disclosure of Interests: José Luis Martín-Varillas: None declared, Eva Galíndez: None declared, Esteban Rubio Romero: None declared, Agustí Sellas-Fernández: None declared, Roberto Daniel Gonzalez Benítez: None declared, Beatriz Joven-Ibáñez Speakers bureau: Celgene, Novartis, MSD, Pfizer, AbbVie, and Janssen, José Campos Esteban: None declared, Olga Rusinovich: None declared, Vanesa Calvo-Río: None declared, Francisco Ortiz-Sanjuán: None declared, Clara Ventín-Rodríguez: None declared, Luis Fernández-Dominguez Speakers bureau: Celgene, Novartis, Janssen, Antia Crespo Golmar: None declared, Ximena Elizabeth Larco Rojas: None declared, Manuel Moreno: None declared, Alejandro Escudero Contreras: None declared, Emma Beltrán: None declared, Rafael Melero: None declared, Maria Jose Moreno: None declared, Enrique Raya: None declared, Beatriz Arca: None declared, María-Luisa Peral: None declared,

Olga Maiz: None declared, Raul Veroz Gonzalez: None declared, Alfonso Corrales: None declared, Natalia Palmou-Fontana: None declared, Belén Atienza-Mateo: None declared, J. Loricera: None declared, Miguel A González-Gay Grant/research support from: Prof. MA Gonzalez-Gay received grants/research supports from Abbvie, MSD, Jansen and Roche., Speakers bureau: Consultation fees/participation in company sponsored speaker's bureau from Pfizer, Lilly, Sobri, Celgene, Novartis, Roche and Sanofi., Ricardo Blanco Grant/research support from: Abbvie, MSD, and Roche, Consultant for: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen, Speakers bureau: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen

DOI: 10.1136/annrheumdis-2019-eular.7803

AB0763

SAFETY OF ABATACEPT TREATMENT OVER 2 YEARS IN A PHASE III ACTIVE PSORIATIC ARTHRITIS RANDOMIZED TRIAL (ASTRAEA)

Philip J. Mease¹, Alice B. Gottlieb², Désirée van der Heijde³, Oliver Fitzgerald⁴, Alyssa Johnsen⁵, Marleen Nys⁶, Subhashis Banerjee⁵, Harris A. Ahmad⁵, Dafna D. Gladman⁷. ¹Swedish Medical Center and University of Washington, Seattle, United States of America; ²Icahn School of Medicine at Mount Sinai, New York, United States of America; ³Leiden University Medical Centre, Leiden, Netherlands; ⁴University College Dublin, Dublin, Ireland; ⁵Bristol Myers Squibb, Princeton, United States of America; ⁶Bristol Myers Squibb, Brainerd Alleud, Belgium; ⁷University of Toronto, Toronto, Canada

Background: In the placebo (PBO)-controlled Phase III ASTRAEA study (ClinicalTrials.gov, NCT01860976) in psoriatic arthritis (PsA), abatacept (ABA) 125 mg SC weekly significantly increased the ACR20 response rate at Week 24 versus PBO (39.4% vs 22.3%, respectively; p<0.001; primary endpoint).¹ Numerical benefits in physical function, enthesitis and dactylitis and a modest impact on psoriasis lesions were also seen at Week 24. Efficacy was maintained or improved up to Week 52.¹

Objectives: To assess ABA safety up to 2 years in patients (pts) enrolled in ASTRAEA.

Methods: Pts with active PsA were randomised 1:1 to ABA or PBO for 24 weeks followed by 28 weeks' open-label (OL) ABA (total time on study: 52 weeks); pts without ≥20% improvement in joint counts at Week 16 (early escape) were switched to OL ABA (total time on study: 44 weeks).¹ Pts could subsequently receive OL ABA in a 52-week long-term extension (LTE; Year 2) for the collection of safety data. The proportions of pts in the cumulative ABA population (all pts who received ≥1 dose of ABA at any time in the study) with AEs (all AEs, serious AEs [SAEs], deaths, AEs leading to discontinuation), laboratory marked abnormalities and anti-ABA antibodies over the cumulative ABA period (date of first treatment dose to end of LTE) were recorded in 1-year and 6-month intervals.

Table 1. Summary of safety up to Year 1 and Year 2 (cumulative ABA period; cumulative ABA population)

	From baseline up to Year 1 (N=398)		From baseline up to Year 2 (N=398)	
	n (%)	Incidence rate per 100 patient-years	n (%)	Incidence rate per 100 patient-years
Deaths	0	0	0	0
SAEs	34 (8.5)	12.1	50 (12.6)	8.9
Treatment-related ¹ SAEs	5 (1.3)	NA	10 (2.5)	NA
Discontinued due to SAEs	8 (2.0)	NA	8 (2.0)	NA
AEs	273 (68.6)	184.9	322 (80.9)	151.2
Treatment-related AEs	81 (20.4)	NA	104 (26.1)	NA
Discontinued due to AEs	13 (3.3)	NA	18 (4.5)	NA
AEs reported in ≥5% of patients				
Nasopharyngitis	25 (6.3)	9.0	44 (11.1)	7.7
Upper respiratory tract infection	28 (7.0)	10.2	40 (10.1)	7.0
Bronchitis	26 (6.5)	9.4	38 (9.5)	6.6
Urinary tract infection	18 (4.5)	6.4	29 (7.3)	5.0
Influenza	—	—	29 (7.3)	4.9
Back pain	—	—	23 (5.8)	3.9
Diarrhoea	—	—	22 (5.5)	3.7
Sinusitis	—	—	21 (5.3)	3.5
AEs of special interest				
Infections ²	162 (40.7)	75.9	226 (56.8)	62.7
Malignancies ³	4 (1.0)	1.4	4 (1.0)	0.7
Autoimmune events ⁴	1 (0.3)	0.4	4 (1.0)	0.7
Local injection-site reactions ⁵	5 (1.3)	1.8	6 (1.5)	1.0
AEs within 24 hours ⁶	94 (23.6)	38.7	141 (35.4)	34.0

¹Causal relationship between study drug and an AE was determined by an investigator physician

²SAEs of infection reported during the LTE included osteomyelitis, intervertebral discitis and cellulitis. Three pts discontinued during the LTE due to the following infections: hepatitis A, Epstein-Barr virus infection and intervertebral discitis

³Bowen's disease, carcinoma in situ of skin, prostate cancer and transitional cell carcinoma. Only transitional cell carcinoma was treatment related. No malignancies were reported during the LTE

⁴Uveitis, hyperthyroidism, Sjögren's syndrome and celiac disease. No autoimmune events reported in the LTE were serious or led to discontinuation of ABA therapy

⁵No local injection-site reactions led to treatment discontinuation

⁶None were suggestive of systemic immune reactions

ABA=abatacept; LTE=long-term extension; NA=not available; pt=patient; SAE=serious AE

Results: Overall, 398 pts were included in the analysis of the cumulative ABA period (213 received ABA from study Day 1; 185 completed initial randomisation to PBO and received ABA thereafter). Of these, 322 pts entered the LTE (initial randomisation: ABA, n=162; PBO, n=160) and continued to receive OL ABA beyond the 28-week OL period. Of 162 and 160 pts initially randomised to ABA and PBO, respectively, 19 (11.7%) and 21 (13.1%) pts discontinued ABA in the LTE. Median (range) number of ABA injections in the cumulative ABA period to end of LTE was 80 (1–133). Safety during the cumulative ABA period over 2 years is shown (Table 1); 6-month interval safety data will be available at time of presentation. In the LTE, there were 3 SAEs of infection (osteomyelitis, intervertebral discitis and cellulitis) and 3 pts discontinued due to infection (hepatitis A, Epstein-Barr virus, intervertebral discitis). No malignancies or deaths were reported, and no autoimmune events were reported as serious or led to ABA discontinuation during the LTE. AEs, SAEs and AEs of special interest occurred with a similar frequency regardless of concomitant MTX or prior TNF inhibitor (TNFi) use. The proportion of pts with anti-ABA antibodies during the cumulative ABA period was higher during the post-treatment period (≥ 22 days after last dose; 37.1% [91/245]) than in the on-treatment period (first treatment dose date to ≤ 21 days after last dose; 7.1% [28/392]). No apparent relationship between anti-ABA antibodies and efficacy in the on-treatment period or AEs suggestive of systemic immune reactions was seen.

Conclusion: Abatacept was well tolerated up to 2 years with no new safety signals during the LTE in this Phase III study in PsA. There was no impact of concomitant MTX use or prior TNFi exposure on the safety profile of abatacept. The occurrence of anti-abatacept antibodies had no impact on abatacept efficacy or safety.

REFERENCES

- [1] Mease PJ, et al. *Ann Rheum Dis* 2017;76:1550–8.

Acknowledgement: Professional medical writing: Andrea Plant, PhD, Caudex; funding: Bristol-Myers Squibb.

Disclosure of Interests: Philip J Mease Grant/research support from: AbbVie, Amgen, BMS, Celgene, Janssen, Lilly, Novartis, Pfizer, SUN and UCB, Consultant for: AbbVie, Amgen, BMS, Galapagos, Gilead Sciences, Inc., Janssen, Lilly, Novartis, Pfizer, SUN and UCB, Speakers bureau: AbbVie, Amgen, BMS, Celgene, Genentech, Janssen, Lilly, Novartis, Pfizer and UCB, Alice B Gottlieb Grant/research support from: Pl: Incyte Corporation, Janssen-Ortho Inc., Lilly ICOS LLC, Novartis, UCB, XBiotech, Consultant for: AbbVie, Dermira, Incyte Corporation, Lilly ICOS LLC, Novartis, Sun Pharmaceutical Industries Ltd., Avotres (unpaid), XBiotech (unpaid), Speakers bureau: AbbVie, Eli Lilly and Company, Janssen Biotech; advisory board: Bristol-Myers Squibb, Celgene Corporation, Janssen Biotech, Janssen-Ortho Inc., LEO Pharma, Novartis, UCB, Désirée van der Heijde Consultant for: AbbVie, Amgen, Astellas, AstraZeneca, Bristol-Myers Squibb, Boehringer Ingelheim, Celgene, Daiichi, Eli-Lilly, Galapagos, Gilead, GlaxoSmithKline, Janssen, Merck, Novartis, Pfizer, Regeneron, Roche, Sanofi, Takeda, Union Chimique Belge, Oliver FitzGerald: None declared, Alyssa Johnsen Shareholder of: Bristol-Myers Squibb, Employee of: Bristol-Myers Squibb, Marleen Nys Shareholder of: Bristol-Myers Squibb, Employee of: Bristol-Myers Squibb, Subhashis Banerjee Shareholder of: Bristol-Myers Squibb, Employee of: Bristol-Myers Squibb, Harris A Ahmad Shareholder of: Bristol-Myers Squibb, Employee of: Bristol-Myers Squibb, Dafna D Gladman Grant/research support from: AbbVie, Amgen, Celgene, Lilly, Novartis, Pfizer, and UCB, Consultant for: AbbVie, Amgen, BMS, Celgene, Galapagos, Gilead, Janssen, Lilly, Novartis, Pfizer, and UCB

DOI: 10.1136/annrheumdis-2019-eular.344

AB0764

USE OF IXEKIZUMAB IN A THIRD LEVEL HOSPITAL

Lara Mendez¹, Isabel Madroñal García¹, Esteban Rubio-Romero¹, Lourdes Rodríguez Freire², Clara Aguilera Cros¹, María Dolores Arcila Duran¹, Marina Gomez Vargas¹. ¹hospital universitario virgen del rocío, rheumatology, seville, Spain; ²HOSPITAL UNIVERTARIO VIRGEN DEL ROCIO, dermatology, seville, Spain

Background: Psoriatic arthritis (PsA) is a chronic inflammatory disease mediated by the immune system, which affects the musculoskeletal system and the skin. It is a disease with increased expression of TNF-alpha

and IL23/IL17 cytokines, which are therapeutic targets whose neutralization has a great impact on the control of the disease.

Ixekizumab is a monoclonal antibody inhibitor of IL-17 indicated for the treatment of moderate-severe psoriasis and PsA. Ixekizumab is administered subcutaneously every 4 weeks, and according to the latest studies, it is a well-tolerated drug and an effective treatment option with significant improvement in the quality of life of patients with cutaneous psoriasis and PsA.

Objectives: -Describe the clinical characteristics, tolerance, safety and survival of a cohort of patients diagnosed with psoriasis or PsA, who have started treatment with Ixekizumab.

-Correlate skin involvement and joint with the quality of life of the patient through the life questionnaire (DLQI) in patients treated with Ixekizumab.

Methods: An observational, cross-sectional and retrospective study was conducted on a cohort of 29 patients who started treatment with Ixekizumab in our hospital in the last year (2018). The data were obtained through the review of patients' medical records and through telephone contact to carry out quality of life questionnaires (DLQI).

Results: Twenty-nine patients were included, of which 22 were males (79.5%). The mean age of the patients was 47 years. Of the cases included, 22 of the patients presented cutaneous Psoriasis (75.8%), 6 of the patients with PsA (20.7%) and 1 patient presented with Dermatomyositis antiMDA 5 (3.5%). Of the 29 patients included, 24 (82.2%) did not present adverse effects. Of the 5 patients who presented adverse effects, in 100% of the cases they were mild. All patients had received prior therapy with disease modifying drugs (DMARDs), and 31% were naive to biological therapy. A total of 4 patients required concomitant therapy with corticosteroid treatment, 3 of them with PAs and 1 with dermatomyositis anti MDA 5. The average of the DLQI of the patients analyzed was 4 and an average PASI of 1 after the start of the treatment, obtaining an Overall improvement of the skin.

A statistically significant Pearson correlation of 0.895 was found between DLQI and PASI (dependent variable), with a strong degree of association. In which patients with a lower score in the DLQI had a lower PASI.

Regarding the pain produced by the autoinjector, measured according to the EVA scale, it was 2 (mild pain), and none would abandon the treatment for this reason. Of all the patients, 9.1% abandoned the treatment due to inefficiency.

Conclusion: The treatment with Ixekizumab presents some promising results in the treatment of cutaneous psoriasis and PsA.

Treatment with Ixekizumab presents a significant improvement in the quality of life of patients.

Ixekizumab is a well-tolerated drug with good efficacy in the short-medium term. It has also been shown to be a safe drug among the patients included in the study.

However, long-term studies with a larger number of patients are required to evaluate the efficacy and safety of Ixekizumab.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5766

AB0765

THE IMPACT OF PSORIASIS SEVERITY ON OUTCOMES AMONG PSORIATIC ARTHRITIS PATIENTS RECEIVING ADALIMUMAB

¹Brigham and Women's Hospital and Harvard Medical School, Boston, United States of America, ²University of Oxford, Oxford, United Kingdom, ³AbbVie, Inc., North Chicago, United States of America, ⁴Swedish Medical Center/Providence St. Joseph Health, Seattle, United States of America

Background: Psoriatic arthritis (PsA) is a heterogeneous, chronic, immune mediated disease associated with psoriasis (PsO). The American College of Rheumatology (ACR) and National Psoriasis Foundation (NPF) recently released a guideline for treatment of PsA, in which TNF inhibitors are recommended over IL-17 or IL-12/23 inhibitors in pts with active PsA.¹ The ACR/NPF conditionally recommend that IL-17 or IL-12/23 inhibitors may be used instead of TNF inhibitors in pts with severe PsO (defined as a psoriasis activity and severity index [PASI] ≥ 12 or a body surface area [BSA] of $\geq 10\%$).

Objectives: Evaluate the impact of PsO severity on clinical, functional, and structural outcomes among PsA pts treated with adalimumab (ADA) or placebo (PBO).