AB0808

COMBINATION OF ORAL PARACETAMOL AND TOPICAL NSAIDS FOR OSTEOARTHRITIS PAIN: A SYSTEMATIC SCOPING REVIEW OF THE LITERATURE

John Bell¹, Vidhu Sethi², Kamran Siddiqui², Philip G. Conaghan³. ¹Graduate School of Health, University of Technology Sydney, Sydney, Australia; ²GlaxoSmithKline Consumer Healthcare, Singapore, Singapore; ³Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, United Kingdom

Background: Topical NSAIDs and paracetamol (APAP) are frequently used for osteoarthritis (OA) pain.^{1,2} In real-world settings, combination treatment is common,^{3,4} with more than one-quarter of patients using topical NSAIDs with oral non-opioid analgesics such as APAP.⁴

Objectives: We conducted a systematic scoping review to: understand current recommendations on concomitant use of topical NSAIDs and APAP for OA pain; evaluate the extent of supporting evidence for this practice; and identify literature gaps.

Methods: We searched PubMed and the Cochrane Library for clinical studies (database inception to January 2019) of topical NSAIDs in OA in which APAP was given concomitantly, either regularly (combination treatment) or as needed (rescue treatment). Grey literature searches were conducted to identify clinical practice guidelines for management of OA from leading organizations.

Results: After removing duplicates, the literature searches returned 375 articles, of which 66 were clinical studies of topical NSAIDs in OA and were reviewed further. Of these, we identified 1 randomized controlled trial (RCT) evaluating efficacy and safety of combination treatment with APAP and a topical NSAID, in which 43 patients with knee OA currently treated with APAP were randomly assigned to double-blind treatment with ketoprofen plaster or placebo for 4 weeks.⁵ Results showed that the combination of APAP/topical ketoprofen was statistically significantly more effective in pain reduction ($P=0.03$) and physician's global assessments ($P=0.01$) compared with APAP/placebo. Of the remaining 65 topical NSAID studies, another 37 (57%) allowed rescue treatment with APAP during the treatment period (including 34 RCTs; 1 open-label extension of 2 RCTs; 1 patient preference study; and 1 single-arm open clinical trial). Because none of these studies randomly assigned patients to the combination, it was not possible to evaluate efficacy or safety of the combination based on these data. Review of OA guidelines identified 3 major organizations—EULAR, NICE, and ESCO—that recommend or allow concomitant use of topical NSAIDs and APAP for specific joints.⁶⁻⁸

Conclusion: In clinical practice, drug combinations are frequently used to manage OA pain, and some OA treatment guidelines allow the use of topical NSAIDs concomitantly with APAP. Very limited supporting data are available, with only a single small RCT demonstrating the efficacy and safety of the combination. However, it may be inferred from the fact that more than half of the topical NSAID studies identified allowed APAP rescue therapy that this combination is generally perceived to be efficacious and well tolerated. Additional, larger RCTs are needed to confirm the benefit of this commonly used analgesic combination in managing OA pain.

REFERENCES

- [1] Conaghan PG, et al. *Clin Rheumatol*. 2015;34:1581-8.
- [2] Jackson H, et al. *BMJ Open*. 2017;7:e019694.
- [3] Gupta R, et al. *Int J Res Med Sci*. 2018;6:985-9.
- [4] Wilson N, et al. *Rheumatology*. 2015;54:860-7.
- [5] Yoo B, et al. *J Korean Rheum Assoc*. 1996;3:70-5.
- [6] Bruyere O, et al. *Semin Arthritis Rheum*. 2014;44:253-63.
- [7] Geenen R, et al. *Ann Rheum Dis*. 2018;77:797-807.
- [8] NICE. Guideline CG177. 2014.

Disclosure of Interests: John Bell Consultant for: John Bell has acted as consultant for Bayer, GlaxoSmithKline, Mylan, Novartis, Pfizer and Reckitt Benckiser., Speakers bureau: John Bell has received payment for speaking from GlaxoSmithKline, Pfizer and Reckitt Benckiser., Vidhu Sethi Employee of: Vidhu Sethi is an employee of GlaxoSmithKline Consumer Healthcare, Singapore., Kamran Siddiqui Employee of: Kamran Siddiqui is an employee of GlaxoSmithKline Consumer Healthcare, Singapore., Philip G Conaghan Consultant for: Flexion Therapeutics, AbbVie, Medivir, Merck Serono, Novartis, GlaxoSmithKline

DOI: 10.1136/annrheumdis-2019-eular.7257

AB0809

A STUDY TO COMPARE ARTICULAR CARTILAGE VOLUME IN HEALTHY AND OSTEOARTHRITIC KNEE AND ITS CORRELATION WITH CLINICO – RADIOLOGICAL SEVERITY

Sudeepti Srivastava, Rajeswar Nath Srivastava, Amar Chandra Sharma. *King George's Medical University, Orthopaedic Surgery, Lucknow, India*

Background: Knee Osteoarthritis (KOA) is a persistent debilitating disease characterized by loss of articular cartilage. Known to be a disease of elderly, it may not be necessarily so, as the loss in articular cartilage volume (ACV) begins early but detected very late on X-rays and by that time no therapy works and joint replacement is the only answer.

Objectives: To determine the role of ACV in early diagnosis of KOA, which may not only monitor disease progression from a very early stage, it may also open a gateway to slow down or stop the structural changes involved in KOA.

Methods: 60 Cases and equal number of age, sex matched controls were recruited. Each subject had MRI of the reference knee. ACV was measured manually by means of image processing on an independent workstation using semi-automated machine GE Signa. VAS for knee pain and WOMAC for pain, stiffness and disability were recorded for clinical severity and X-rays for radiological severity by KL grading.

Results: With age, a statistically significant inverse correlation of ACV was found in both healthy and osteoarthritis knees. Height, weight and BMI were independent of ACV in healthy knee. In KOA, a significant positive correlation of ACV was observed with height, a significant inverse correlation with WOMAC scores and significant difference with sub-scales of WOMAC index (pain, stiffness and physical function). The statistical difference in ACV between the two categories of WOMAC scores (<32 and >32) was significant.

Conclusion: This study concluded that ACV is significantly lower in osteoarthritis knee as compared to healthy adult knee. This study also found that KOA in females occurs at an early age and progresses slowly whereas in males occurs late and progresses rapidly. ACV may become one of the most promising tools for early diagnosis of KOA and in monitoring disease progression.

REFERENCE

- [1] Beary JF III. Joint structure modification in osteoarthritis: development of SMOAD drugs. *Curr Rheumatol Rep* 2001;3:506-512.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6708

AB0810

INTRA-ARTICULAR CNTX-4975 FOR PAINFUL KNEE OSTEOARTHRITIS: ASSESSMENT OF COOLING METHODS FOR REDUCING PROCEDURAL PAIN

Randall Stevens¹, Kimberly Guedes¹, Nilam Mistry¹, Duncan Lascelles¹, David Ball². ¹Centrexion Therapeutics Corp, Boston, United States of America; ²MAC Clinical Research, Manchester, United Kingdom

Background: CNTX-4975 is a long-acting, *trans*-capsaicin injection in phase 3 trials for treatment of moderate to severe pain associated with knee osteoarthritis (OA). Intra-articular (IA) CNTX-4975 injection produces short-lived procedural pain that can be ameliorated with joint cooling. A prior analysis (Cohort 2) demonstrated that a circumferential circulating ice water wrap (CCIWW) more effectively lowered IA knee temperature and reduced procedural pain than an ice pack on top of the knee (3.4 vs 7.3, respectively, on a numeric pain rating scale [NPRS] 10 minutes post-CNTX-4975).

Objectives: Subjects were enrolled in 2 separate cohorts (C3/C4) to: compare effects of 2 circumferential cooling methods on IA and skin knee temperature and procedural pain (C3); and assess procedural pain with an abbreviated cooling schedule and with vs without post-CNTX-4975 cooling (C4).

Methods: Eligible subjects were adults aged 45–75 yrs with ≥3 months of bilateral moderate to severe OA knee pain. All subjects received each

cooling device on opposite knees; IA CNTX-4975 injections and other procedures on the left and right knees were separated by 7 (± 2) days. See **Figure** for study design. IA and skin temperatures were recorded throughout the procedures. Subjects rated procedural pain at prespecified times using NPRS (0=no pain; 10=worst possible pain).

Results: Five subjects enrolled in and completed each cohort (mean age, yrs: C3, 57.6; C4, 59.6; each 80% male). In C3, mean IA temperatures decreased with both methods; mean temperatures at baseline and 105 minutes were 33.7°C and 26.7°C for ice-gel pack cooling and 32.4°C and 28.2°C for CCIWW. Skin temperature reductions between cohorts were similar; mean temperatures were 28.8°C and 14.3°C (ice-gel pack) and 28.7°C and 15.2°C (CCIWW). Mean IA temperatures in C4 at baseline and 55 minutes (15 minutes post-CNTX-4975 and/or end of last cooling period [CCIWW only]) were 33.6°C and 29.0°C for ice-gel pack cooling and 32.1°C and 27.7°C for CCIWW. Mean skin temperatures were 28.5°C and 22.8°C for ice-gel pack and 27.8°C and 18.8°C for CCIWW. Pain levels before CNTX-4975 were low. Procedural pain peaked 10–20 minutes after CNTX-4975 injection in both cohorts. While mean pain levels were similar across cohorts (**Table**), pain was reduced more effectively by CCIWW in C3 and ice-gel pack in C4. Procedural pain was not increased in the absence of post-injection cooling in C4.

Conclusion: The ice-gel pack and CCIWW effectively reduced IA knee joint temperatures and procedural pain in both cohorts. Shorter cooling time and low pain levels observed with IA CNTX-4975 and ice-gel pack cooling in C4 suggest that this strategy may be feasible in clinical practice.

Procedural Pain Over Time (min), Mean $\pm$ SE ^a	Cohort 3		Cohort 4	
	Ice-gel pack	CCIWW	Ice-gel pack	CCIWW
Baseline	0.8 $\pm$ 0.37	0.4 $\pm$ 0.24	0.4 $\pm$ 0.24	0.2 $\pm$ 0.20
10 ^b	4.2 $\pm$ 1.36	2.2 $\pm$ 1.46	2.4 $\pm$ 1.29	4.2 $\pm$ 1.20
20 ^b	3.6 $\pm$ 1.33	2.2 $\pm$ 1.20	3.0 $\pm$ 1.22	2.6 $\pm$ 1.25
30 ^b	2.6 $\pm$ 0.81	2.2 $\pm$ 0.97	2.0 $\pm$ 1.05	1.8 $\pm$ 1.11
Last assessment ^c	2.4 $\pm$ 0.81	1.4 $\pm$ 0.40	1.6 $\pm$ 1.03	2.4 $\pm$ 1.60

Cohort 3: Comparison of 2 circumferential cooling methods; Cohort 4: Shortened cooling and with/without post-CNTX-4975 cooling.
^aLower scores=less pain; ^bPost-CNTX-4975 injection; ^cApproximately 70 min post-injection for C3; 60 min for C4.

Figure: Study Procedures

A. Study Cohort 3

*Without epinephrine, 15 mL. Subjects rated pain throughout the study (orange circles). Unfilled orange circles represent time points presented in the table.

B. Study Cohort 4

Disclosure of Interests: Randall Stevens Shareholder of: Centrexion Therapeutics Corp, Employee of: Centrexion Therapeutics Corp, Kimberly Guedes Employee of: Centrexion Therapeutics Corp, Nilam Mistry Employee of: Centrexion Therapeutics Corp, Duncan Lascelles Consultant for: Centrexion Therapeutics Corp, David Ball: None declared

DOI: 10.1136/annrheumdis-2019-eular.1156

AB0811

MULTICENTER PROSPECTIVE STUDY TO ASSESS EFFICACY AND SAFETY OF INTRA-ARTICULAR INJECTIONS OF A SOLUTION COMBINING HYALURONIC ACID AND CHONDROITIN SULPHATE IN PATIENTS WITH KNEE OSTEOARTHRITIS

Elena Taskina¹, Ludmila Alekseeva¹, Natalia Kashevarova¹, Evgenia Sharapova¹, Ekaterina Strebkova¹, Sergey Anikin¹, Lena Zonova², Tatiana Raskina³, Elvira Otteva⁴, Aleksandr Lila¹, ¹V. A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation; ²FSBEI HE Novosibirsk State Medical University, Novosibirsk, Russian Federation; ³FSBEI HE KemSMU MOH, Kemerovo, Russian Federation; ⁴SBHI CDC MOH Khabarovsk kray, Khabarovsk, Russian Federation

Background:

Objectives: To assess efficacy and safety of a single intra-articular injection of a solution combining hyaluronic acid (HA) 60 mg and chondroitin sulphate (CS) 90 mg in 3 mL on patients with knee osteoarthritis (OA) in a multicenter prospective study.

Methods: 79 outpatients (predominantly females - 81.0%) from 5 RF constituent territories with primary tibiofemoral Kellgren-Lawrence score grade II or III knee OA, ≤ 40 mm pain intensity during walking on visual analogue scale (VAS), requiring NSAIDs intake (for at least 30 days during 3 months prior to enrollment) were included into the study after signing the informed consent form. Mean age was 60.3 $\pm$ 8.7 years, mean BMI - 29.2 $\pm$ 4.7 kg/m², disease duration - 6 (3 - 10) years. Grade II OA was documented in 68.4% of patients, Grade III - in 31.6%. The study lasted for 6 months. Efficacy and safety evaluations were made based on VAS pain assessment, Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) - (WOMAC pain (0-500), WOMAC function (0-1700), WOMAC stiffness (0-200)), VAS patients' health status, EQ-5D-based assessment of patients' quality of life, global physician's and patient's efficacy assessment, and daily NSAIDs requirements.

Results: Obtained results demonstrate statistically significant VAS pain reduction during walking already in 1 week after intra-articular injection of the combination (respectively, 62 (55-72) and 41 (32-51) mm, $p < 0.0001$). Moreover, pain continued to subside during all 3 months of follow up (in 1 month - 28 (20-42), in 3 months - 22 (14-37) mm). A significant pain reduction achieved at Mo 3 persisted until Mo 6 - 20 (14-42) mm, without documented pain increase (Fig. 1). Similar trends were observed with total WOMAC score (1125 (899-1540) - at baseline, and 552 (309-837) mm - by the end of the study, $p < 0.0001$), and all WOMAC sub-scores (268 (189-312) - baseline WOMAC pain, 91 (48-171) mm - by the end of the study $p < 0.0001$ (Fig. 2); stiffness - 101 (59-130) and 40 (20-61) mm, $p < 0.0001$; function - 802 (647-1095) and 402 (191-638) mm, $p < 0.0001$, respectively). Median time to the onset of therapeutic effect was 7 (5-18) days. Statistically significant improvement of patients' quality of life by EQ-5D and general health status was observed during all follow up period (respectively, 0.52 (-0.02-0.59) and 0.69 (0.59-0.80), $p < 0.0001$; 48 (30-60) and 72 (60-80) mm, $p < 0.0001$). One injection of the drug resulted in dose reduction or discontinuation of NSAIDs therapy: at baseline 76 patients (96.2%) were taking NSAIDs, in one week 31 (39.2%) patients discontinued NSAIDs, in 1 month - 72.2%, in 3 months - 73.4%, and by the end of the study at Mo 6 - 54.4% were not taking NSAIDs. These data were consistent with physician's and patient's global assessment of the efficacy of treatment, who stated «significant improvement» and «improvement» in the majority of cases, with only few «no effect» or «worsening» cases documented in analyzed population. Adverse events, such as worsening of pain and/or swelling of the joint, were documented in 8 patients (10.1%); they resolved spontaneously or following NSAIDs intake.

Fig.1. VAS pain, mm