Figure 3.- Average of number of reported painful knee locations after pressure of <4Kg/cm2.

Figure 4.-Knee normalized Medio-Lateral forces (a) and Abduction-Adduction torques (b)

Conclusion: Although both treatment groups present the same OA radiologic grade, ART group presents significant higher stiffness and functional disability. That may affect to the gait of these ART patients, altering the forces distribution and torques between both legs. Although no differences in knee pain in life situation (Wp) between treatment group are reported, ART group present more painful sites in the knee with pressure stimuli. Emotional component may be playing a role in the pain and illness perception, influencing the patient decision to undergo ART surgery

REFERENCES

- Grazina, R, Adv Exp Med Biol.1059 111-135, 2018
- Pujol, J et al, Pain. 158(9) 1831-1838, 2017.
- R.B. Davis, Hum. Mov. Sci. 10 575-587, 1991.

Acknowledgement: Funds from the MICINN (HOLOA-DPI2016-80283-C2-1-R and DPI2016-80283-C2-1-R (AEI/FEDER, UE), RYC-2015-18888) and DTIC-UPF are acknowledged

Disclosure of Interests: Laura Tío: None declared, Francisco Castro Speakers bureau: Lilly, Simone Tassani: None declared, Santos Martinez: None declared, Raul Torres: None declared, Raquel Aredondo: None declared, Miguel Ángel González-Ballester: None declared, Joan Carles Monllau: None declared, Jérôme Noailly: None declared, Jordi Monfort Speakers bureau: Bioibérica

Procure Health

DOI: 10.1136/annrheumdis-2019-eular.5856

AB0813

AN EXPERT CONSENSUS ON THE APPROPRIATE USE OF ORAL SYSADOAS FOR THE TREATMENT OF THE OSTEOARTHRITIC PATIENT IN PRIMARY HEALTH CARE: A DELPHI STUDY

Jordi Monfort¹, Benjamín Abarca², Xavi Carné³, Sergio Giménez⁴, Ingrid Möller⁵, Montserrat Romera⁶, Marco Bibas⁷, Marianna Vitaloni⁷, Josep Verges⁷. ¹Hospital del Mar, Barcelona, Spain; ²Centro Salud Sagrado Corazón, Lugo, Spain; ³Hospital Clínic, Barcelona, Spain; ⁴Centro De Salud Y Atención Primaria Limonar, Málaga, Spain; ⁵Institut Poal de Reumatologia, Barcelona, Spain; ⁶Hospital de Bellvitge, L'Hospitalet de Llobregat, Spain; ⁷OAFI Foundation (Osteoarthritis Foundation International), Barcelona, Spain

Background: Clinical studies have demonstrated that osteoarthritic pain is linked to disability and quality of life (QoL). The therapeutic modalities in the treatment of osteoarthritis (OA) are numerous and despite the availability of evidence-based guidelines for OA management, agreement on

treatments is lacking¹. Symptomatic Slow-Acting Drugs for OA (SYSADOA), are natural compounds, used in OA treatment. Thus, there is disagreement about SYSADOAs use in clinical practice.

Objectives: Our objective was to prepare a consensus document on the appropriate use of oral SYSADOAs: chondroitin sulphate (CS), glucosamine(G), diacerein(D) and the combination of CS plus G for OA management in primary health care.

Methods: A two-round Delphi study was carried out to assess expert consensus on the appropriate use of SYSADOAs in primary care. The questionnaire validated by the expert committee (3 rheumatologists, 2PC physicians, 1 clinical pharmacologist) included 24 questions The Delphi panel was composed of 15 experts (10 PC physicians, 1 rheumatologist, 1 traumatologist, 1 rehabilitator, 1 gynaecologist and 1 clinical pharmacologist) with extensive experience in the treatment of OA and the use of oral SYSADOAs which were identify by clinical coordinator, a methodological coordinator, and four members of the Scientific Committee. Participants were asked 24 questions on SYSADOAs use. Items that reached consensus by at least 80% across both panels were included in the guidelines. The fieldwork of the study was developed for approximately 4.5 months. This study was promoted by the International Osteoarthritis Foundation (OAFI) with the support of the Spanish Ministry of Health, Social Services and Equality.

Results: Consensus statements emerged: (1)patient phenotypes affects SYSADOAs action; (2)SYSADOAs are effective in primary and secondary OA, in the three first grade of Knee OA, hand and hip; there is no evidence for erosive hands, shoulder, spine, and ankle OA; (3)CS, G and association can reduce pain, inflammation, improve QoL and functional capacity and have a chondroprotective effect; (4)CS and D can reduce synovial membrane inflammation, all oral SYSADOAs, except D, can decrease cell death and the enzymes responsible for cartilage destruction; (5) The maximum therapeutic efficacy is reached after 3 to 6 months; (6) SYSADOAs can be prescript to patients having comorbidities: cardiovascular risk or disease, digestive disease, hypertension, dyslipidemia, peripheral vascular disease, type 2 diabetes, and oesophageal reflux. There is disagreement in the prescription of oral SYSADOA in patients with liver and kidney disease.

Conclusion: This study sheds light on the appropriate use of oral SYSADOAs in primary health care by providing added value to published evidence. Results based on literature evidence on efficacy and safety, the clinical experience of the panellist experts in OA treatment and the fact that OA patient is a chronic, elderly, with multiple diseases and polymedicated person. The diffusion of our results among primary health practitioners will contribute to improving OA patient management protocols to ensure a personalized treatment to OA patients and to ameliorate their QoL.

REFERENCES

- Ethgen O, Reginster JY. Degenerative musculoskeletal disease. Ann Rheum Dis 2004;63:1-3.

Disclosure of Interests: Jordi Monfort Speakers bureau: Bioibérica

Procure Health, Benjamín Abarca: None declared, Xavi Carné: None declared, Sergio Giménez: None declared, Ingrid Möller: None declared, Montserrat Romera: None declared, Marco Bibas: None declared, Marianna Vitaloni: None declared, Josep Verges: None declared

DOI: 10.1136/annrheumdis-2019-eular.4193

AB0814

INDIVIDUAL AND SOCIAL FACTOR CAN INFLUENCE THE QUALITY OF LIFE OF KNEE OA PATIENTS: A SYSTEMATIC REVIEW

Josep Verges¹, Marianna Vitaloni¹, Marco Bibas¹, Rosa Sciortino², Maritza Quintero², Jordi Monfort³, Francisco de Abajo⁴, Patrick du Souich⁵, Ingrid Möller⁶, Elizabeth Oswald⁷, Marco Matucci-Cerinic⁷, Guy Eakin⁸, Angie Botto-van Bemden⁸. ¹OAFI Foundation (Osteoarthritis Foundation International), Barcelona, Spain; ²Universidad de los Andes, Merida, Venezuela (Bolivarian Republic); ³Hospital del Mar, Barcelona, Spain; ⁴Alcalá University, Alcalá de Henares, Spain; ⁵University of Montreal, Montreal, Canada; ⁶Institut Poal de Reumatologia, Barcelona, Spain; ⁷University of Florence, Firenze, Italy; ⁸ARTHRITIS FOUNDATION, Atlanta, United States of America

Background: Knee OA (KOA) is the most common form of chronic joint disease and bears more responsibility than any other disease for disability¹. It associates with remarkable functional restrictions due to pain. The limitations in activity caused by KOA seriously affect social relationships, emotional well-being, reducing the quality of life (QoL) of patients.