such as Osteopenia or even Osteoporosis. It has been reported in several studies, an incidence of fracture between 30-40% in this population.

Objectives: To evaluate the evolution of bone turnover biomarkers before and after treatment (according to 2015 NOF guidelines) and to describe the incidence of fracture after a SCI.

Methods: A prospective study was conducted, including 48 SCI patients followed during an observation period of 24 months, with visits every six months. Thirty two patients concluded the study. In each visit patients were enquired about fractures. A blood test was also performed. In addition a dual-energy x-ray absorptiometry (DEXA) and simple x-ray of hip and spine was requested annually. Patients were also educated in life-style measures, such as an adequate calcium and vitamin D dietary intake. Oral supplementation was initiated in cases of low intake or deficit, respectively. In patients with bone mineral density (BMD) measured by DEXA in range of Osteoporosis, pharmacological therapy was initiated (bisphosphonate or Denosumab). Statistics had been performed with mixed linear regression models.

Results: Our sample was 56.25% men with an average age of 56.08 ± 13.82 years old. In 32 patients (68.09%) low Vitamin D levels were detected. At baseline in 12 cases (25%) DEXA showed a BMD in osteoporosis range; in 12 (25%) osteopenia, plus 6 fractures (2.6 fractures per 100 years/patient). Afterwards, until the end of the study only a fracture of the tibia was observed (however, it was after a high impact accident), so the fracture rate was 1.2 fractures per 100 year/patient. The evolution of the variables can be seen on the next table.

	Month-0 N=48 Mean (SD)	Month-6 N=32 Mean (SD)	Month-12 N=36 Mean (SD)	Month-18 N=37 Mean (SD)	Month 24 N=32 Mean (SD)
CALCIUM (mg/dl)	9.1 (0.59)	9.53 (0.4)	9.37 (0.54)	9.46 (0.41)	9.55 (0.44)
PHOSPHOR (mg/dl)	3.63 (0.7)	3.42 (0.44)	3.32 (0.52)	3.33 (0.55)	3.25 (0.45)
VIT D (UI)	16.42 (9.95)	42.28 (18.93)	35.88 (13.53)	35.77 (18.08)	37.55 (15.06)
PTH (ng/mL)	38.15 (25.55)	36.09 (17.89)	38.33 (21.2)	40.1 (17.27)	43.35 (25.08)
PINP (ng/mL)	79.25 (80.9)	68.15 (35.77)	53.05 (45.76)	42.82 (24.26)	40.67 (25.17)
bCTX (ng/mL)	0.82 (0.49)	1.23 (4.22)	0.34 (0.35)	0.28 (0.21)	0.51 (1.36)
T-SCORE SPINE	-0.49 (1.9)		-0.61 (1.26)		-0.24 (1.34)
T-SCORE NECK	-0.95 (1.53)		-1.29 (1.41)		-1.53 (1.16)
T-SCORE HIP	-1.02 (1.58)		-1.38 (1.58)		-1.42 (1.33)
TOTAL					
CALCIUM SUPP	29 (60.42%)	20 (62.5%)	26 (72.22%)	27 (72.97%)	22 (68.75%)
VIT D SUPP	46 (95.83%)	2 (6.25%)	35 (97.22%)	36 (97.3%)	32 (100%)
BIPHOSPHONATE	4 (8.33%)	5 (15.62%)	3 (8.33%)	5 (13.51%)	6 (18.75%)
DENOSUMAB	7 (14.58%)	7 (21.87%)	7 (19.44%)	7 (18.91%)	7 (21.87%)

On multivariate analysis an inverse relationship between vitamin D and CTX levels was detected ($p \leq 0.05$). DEXA hip measures tend to become stabilized after Bisphosphonate treatment.

Conclusion: On our cohort, bone resorption biomarkers decrease after Vitamin D restoring. Antiresorptive therapy in high risk patients stopped bone loss, especially in hip. After two years follow up, with the treatment algorithm applied, fracture incidence was lower than other series published.

REFERENCE

- [1] *J Musculoskelet Neuronal Interact 2011; 11(3):257-265

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5953

AB0842 POTENTIAL RISKS FOR OSTEOPOROSIS AND SARCOPENIA IN ALCOHOLIC WOMEN

Kayo Masuko^{1,2}, Chie Iwahara³, Seiji Sakate⁴, Yumiko Tanaka⁵, Shigemi Kamiya³, Yuki Mizukami⁴. ¹Sanno Medical Center, Tokyo, Japan; ²International University of Health and Welfare, Tokyo, Japan; ³National Hospital Organization Kurihama Medical and Addiction Center, Kanagawa, Japan; ⁴Sagami Women's University, Kanagawa, Japan; ⁵National Hospital Organization Disaster Medical Center, Tokyo, Japan

Background: Recent statistics indicated an accumulating number of patients with alcohol dependency in Japan. In particular, there had been a considerable increase in the number of alcoholic women, reaching up to 130,000 patients. Because alcoholic women often have insufficient

dietary intake and physical inactivity, they may be at risk for low bone mineral density and decreased muscle power. However, these points may not be widely recognized, and the current program for the rehabilitation of alcoholism may not be sufficient to address this issue.

Objectives: To assess the risk for osteoporosis and sarcopenia in alcoholic women from a nutritional viewpoint

Methods: Thirty-two women (mean age: 42.9 ± 7.49 years) who were admitted to the Alcoholics Rehabilitation Program (ARP) for alcoholism, which was diagnosed according to International Classification of Diseases (ICD)-10, were enrolled in the study. The patients were subjected to self-answered questionnaire, nutritional assessment by expert dietitians, and measurement of body composition and grip strength.

Results: Of the 32 patients, 14 (44.0%) had a history of eating disorder, mostly anorexia. All 32 patients had elevated levels of transaminases at the time of admission. History of bone fracture was present in eight of the 32 (25.0%) patients; among them, at least three (9.4%) cases were considered to be due to a fragility fracture. The mean body mass index (BMI) was 20.4 kg/m²; 10 (31.3%) patients had BMI < 18. The average energy intake was low at 1,474.5 ± 324.3 kcal. Twenty-one (65.6%) patients were taking dietary calcium of <650 mg/day, which is the level recommended for adult women. The mean grip strength was 23.8 ± 6.5 kg, and 8 (25%) patients had grip strength of only <18 kg, which is the definition of sarcopenia in the Asian elderly.

Conclusion: Young to middle-aged alcoholic women should be treated as having high risks for osteoporosis and sarcopenia. ARP should include multidisciplinary assessment and interventions to prevent future locomotive disabilities.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5262

AB0843 HYPOVITAMINOSIS D IN A MARFAN POPULATION

Daniela Melchiorre¹, Elisa Pratelli², Renato Colombai³, Marco Matucci-Cerinic¹, Guglielmina Pepe¹. ¹University of Florence, Florence, Italy; ²AOU Careggi Hospital, Florence, Italy; ³Department of Public Health, Pontedera, Italy

Background: Marfan syndrome is an autosomal dominant disorder of connective tissue caused by mutations in FBN1 gene on the long arm of chromosome 15. Information is often lacking regarding an increased risk of fractures and its management and prevention in patients with Marfan syndrome. In these patients reduced bone mineral density was rarely reported. A pivotal role in bone mineralization is played by Vitamin D.

Objectives: This study evaluates whether the reduction of the values of Vitamin D is present in Marfan patients.

Methods: 60 patients (pts) (28 M/32 F; age 15-75 mean age 46,2) with Marfan syndrome were studied. The patients were followed from the clinical and research regional Marfan Center in Florence, Italy. All pts met new Ghent criteria that identified ectopia lentis, thoracic aorta dilatation with Z score = or > 2 and a large group of systemic clinical features represented by cardiovascular, musculoskeletal, central nervous system (dural ectasia) pulmonary, cutaneous and ocular clinical manifestations (systemic features) reaching a score = or > 7 as major clinical criterion. Familiarity and the detection of a gene mutation (in MFS FBN1 gene displays a mutation in about 90% of patients) are also considered criteria necessary for Marfan diagnosis.

Results: 11/60 pts (18,3%) (5M/6F age 15-75 mean age 44,9) showed reduced values of Vitamin D. 6/11 pts (45,4%) (3M/3F age 15-56, mean age 34,1) showed very poor Vitamin D values (range 8.7 ng/ml – 13,17 ng/ml).

Conclusion: Our results suggests that Vitamin D may play an important role in bone metabolism in Marfan patients. These results need confirmation since further data are missing in literature.

REFERENCES

- [1] Trifirò G, Marelli S, Viecca M, Mora S, Pini A. Areal bone mineral density in children and adolescents with Marfan syndrome: evidence of an evolving problem.
[2] Bone. 2015 Apr;73:176-80. doi: 10.1016/j.bone.2014.12.006. Epub 2014 Dec 13.

Disclosure of Interests: Daniela Melchiorre: None declared, Elisa Pratelli: None declared, Renato Colombai: None declared, Marco Matucci-Cerinic: Grant/research support from: Actelion, MSD, Pfizer, BMS, Chemomab, Sanipedia, Speakers bureau: Actelion, BMS; MSD, Janssen, Guglielmina Pepe: None declared

DOI: 10.1136/annrheumdis-2019-eular.8229