	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6
Gender	Male	Male	Male	Male	Female	Female
Age of onset of symptoms (years)	4	4	5	2	4	2
Age of diagnosis (years)	5	5	5	3	4	3
Time elapsed for diagnosis (months)	13	12	3	11	4	6
Initial manifestation	-Fever - Thrombocytopenia	-Arthritis	-Convulsive crisis	-Cutaneous vasculitis -Arthritis	-Cutaneous vasculitis -Arthritis	-Deep cutaneous lupus -Autoimmune hepatitis
Number of ACR criteria (1997)	4	6	6	5	4	4
Renal disease	LN Class IV, A:6/24 C:0/ 12)	LN Class III, A:2/24 C:2/ 12	LN Class IV A:7/24 C:3/ 12	LN Class III, A:5/24 C:1/ 12)	No	No

* LN: lupus nephritis. A: activity score. C: chronicity score.

in patients with juvenile idiopathic arthritis (JIA), are currently available¹. With regards to possible predictors of flare, the role of obesity has not been investigated yet.

Objectives: To investigate to role of obesity at start of TNFi therapy on remission duration and flare rate after TNFi withdrawal in JIA patients.

Methods: The charts of JIA patients followed at the study center and treated with TNFi as first biologic from 2009 to 2015, who reached remission and with at least one year of follow up after withdrawal, were retrospectively reviewed. Exclusion criteria were: systemic JIA and pJIA RF-positive. Demographic and clinical features were registered, including: ILAR JIA subtype, occurrence of uveitis, ANA status, number of active joints at TNFi start and at flare onset, BMI (z-score and percentile at time of biologic withdrawal for determining obesity, overweight and underweight), concomitant DMARDs. For each patient we registered the remission duration and the occurrence of flare after 12 months from TNFi withdrawal. For comparison of continuous and categorical variables we respectively used the Mann-Whitney and Fisher's exact probability test. Spearman's test was used for correlation analysis. Data were expressed as median and interquartile range (IQR 25-75) and analyzed with SPSS 25.0. P values < 0.05 were considered significant. Variables with significant association with remission duration were included in the multivariable linear regression model.

Results: 57 JIA patients were included: 34 of them (59.6%) flared within 12 months after TNFi withdrawal with a median of 5 months (IQR 3-8). Obesity was found in 6 patients (10.5%), overweight in 7 (12.2%) and underweight in 6 (10.5%). No statistically significant difference in the aforementioned factors was found in patients who flared, if compared with patients who did not flare. In obese patients we observed a shorter remission duration ($p = 0.04$). In addition, remission duration was inversely correlated with the number of active joints at TNFi start ($r = -0.40$ $p = 0.03$). A multiple linear regression analysis was performed with remission duration as dependant variable and active joints at TNFi start ($p = 0.007$; $B = -0.47$) and obesity ($p = 0.05$ $B = 0.31$) as explanatory variables.

Conclusion: In our cohort the number of active joints at TNFi start and obesity were both associated to shorter remission duration after anti-TNF withdrawal. In particular obesity, a modifiable factor involved in several pathological conditions, should be handled more carefully in order to optimize JIA management. Larger studies need to confirm our preliminary data.

REFERENCES

- [1] Shoop-Worrall SJW, Kearsley-Fleet L, Thomson W, Verstappen SMM, Hyrich KL. How common is remission in juvenile idiopathic arthritis: A systematic review. *Semin Arthritis Rheum*. 2017 Dec;47(3):331-337.

Disclosure of Interests: Fabio Basta: None declared, Hanan Jadoun: None declared, Maria Isabella Petrone: None declared, Angela Aquilani: None declared, Andrea Uva: None declared, Rebecca Nicolai: None declared, Fabrizio De Benedetti Grant/research support from: Abbvie, SOBI, Novimmune, Roche, Novartis, Sanofi, Pfizer, Silvia Magni-Manzoni Consultant for: Abbvie, Speakers bureau: Abbvie

DOI: 10.1136/annrheumdis-2019-eular.6029

AB0941

SYSTEMIC LUPUS ERYTHEMATOSUS IN CHILDREN UNDER 6 YEARS OF AGE, EXPERIENCE IN A TERTIARY CENTER

EDITH ADRIANA BENITEZ VAZQUEZ, Maria Teresa Brana Ruiz, Luis Alberto Aparicio, Sofia Osorio Sagrero, Merari Gomez Cortes, Velka Cortez, Rocio Maldonado, Enrique Faugier Fuentes. *Hospital infantil de Mexico Federico Gomez, Pediatric rheumatology, Mexico City, Mexico*

Background: Systemic lupus erythematosus (SLE) is a rare entity in patients under 5 years old, depending on the ethnic group it has a prevalence of 3.3 to 24 per 100 000 children. Before puberty no difference in prevalence associated with gender was observed. In the pediatric age, SLE has a more aggressive course.¹

Objectives: To describe the demographic and clinical characteristics of 6 patients under 6 years old diagnosed with SLE.

Methods: To describe the demographic and clinical characteristics of 6 patients under 6 years old with a diagnosis of SLE based on the 1997 ACR criteria at the Federico Gomez Children's Hospital of Mexico from June 2017 to January 2019.

Table N°1. Demographic and clinical characteristics of patients under 6 years of age diagnosed with SLE.

Results: The diagnosis of SLE in patients under 5 years old is rare, the literature describes that in these patients there is greater renal, hematological and neuropsychiatric manifestations; We present the demographic and clinical characteristics of 6 patients under 6 years of age with a diagnosis of SLE based on criteria of ACR 1997; 4 of the patients presenting are male, in three of the cases the initial manifestation was non-erosive arthritis and skin manifestations (in two purpura and in one deep cutaneous lupus), in four of the cases lupus nephritis was presented; the average time between the onset of symptoms and the diagnosis was 8 months (range between 3 and 13 months).

Conclusion: The diagnosis of SLE in patients under 6 years old is rare, the initial manifestations may be nonspecific, which causes greater complexity to establish an opportune diagnosis. It is described that in lupus cases in pediatric patients there are clinical manifestations of greater severity and early diagnosis modifies the prognosis. The initial manifestations of SLE in patients younger than 6 years old in our series are similar to those described in the literature: arthritis and seizures, as well as the presentation associated with gender, with no difference in preschool age.² It has been described that the time elapsed from the onset of symptoms to the diagnosis is shorter at younger age of patient, however in our series the time of diagnosis was an average of 8 months, probably related to the low prevalence of the disease in this age group.

REFERENCES

- [1] Malattia C., Martini A. (2013 Jun). Paediatric-onset systemic lupus erythematosus. *Best Pract Res Clin Rheumatol*;27(3):351-62.
- [2] Jia Zhu., Fengqi Wu., Xiaolan Huang. (2013 Jan). Age-related differences in the clinical characteristics of systemic lupus erythematosus in children. *Rheumatol Int* 33:111-115.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.8329