from: Prof. MA Gonzalez-Gay received grants/research supports from Abbvie, MSD, Jansen and Roche., Speakers bureau: Consultation fees/participation in company sponsored speaker's bureau from Pfizer, Lilly, Sobi, Celgene, Novartis, Roche and Sanofi., Ricardo Blanco Grant/research support from: Abbvie, MSD, and Roche, Consultant for: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen, Speakers bureau: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen

DOI: 10.1136/annrheumdis-2019-eular.3621

AB1099

CHAGAS DISEASE REACTIVATION IN THE RHEUMATOLOGIC IMMUNOSUPPRESSED PATIENT. IS IT NO LONGER AN ORPHAN DISEASE?

Ariana Ringer¹, Juan Pablo Ruffino¹, Nadia Cuadranti¹, Florencia Martinez¹, Cecilia Argento¹, Ignacio Rolla¹, Juan Manuel Vandale¹, Carla Achilli¹, Mariano Palatnik¹, Noel Cortese¹, Mariana Lagrutta², Rodolfo Leiva³, Damian Aguila⁴, Laura Cordoba⁵, Milagros Zafra⁵, T Gambande⁵, Patricia Sciaratta⁶, M Svetaz⁶, F Gonzalez⁷, F Pacini⁷, Silvina Villar⁷, Ana Rosa Perez⁷, Oscar Bottasso⁷, Marcelo Abdala¹. ¹Rheumatología, Hospital Provincial del Centenario, Universidad Nacional de Rosario, Rosario, Argentina; ²Clinica Médica, Hospital Provincial del Centenario, Rosario, Argentina; ³Cardiología, Hospital Provincial del Centenario, Rosario, Argentina; ⁴Infectología, Hospital Provincial del Centenario, Rosario, Argentina; ⁵Microbiología, Universidad Nacional de Rosario, Rosario, Argentina; ⁶Laboratorio Central, Facultad de Ciencias Bioquímicas y Farmacéuticas, UNR, Rosario, Argentina; ⁷Instituto de Inmunología Clínica y Experimental de Rosario (IDICER, CONICET, UNR), Rosario, Argentina

Background: Chagas' disease (CD) is an endemic and neglected infection in Latin America. Due to international human migration, it has become a worldwide issue. Scarce evidence is published regarding its behavior in rheumatic diseases (RD) with rheumatologic treatments (RT).

Objectives: To screen and follow up patients with CD under RT. To detect clinical and serological reactivation.

Methods: A systematic screening was conducted between January 2018 and January 2019 in a third-level Hospital in Argentina. Patients with CD and concomitant RD under RT were included. Assessments were done before and after RT. A direct (Strout) and indirect method (Polymerase chain reaction-PCR) were performed in order to detect parasitemia, levels of antibodies were evaluated by three techniques. Clinical, infectological and cardiologic features were examined. Everything was assessed in, at least, two opportunities, separated by a minimum of one month. If treatment was modified or clinical condition changed, all the evaluations were repeated as before. In case of reactivation, specific treatment was indicated and follow up controls were stricter.

Results: 38 patients were identified. RD: Rheumatoid Arthritis: 22 (57.9%), Systemic lupus erythematosus: 4 (10.5%), Systemic Sclerosis: 3 (7.9%), Vasculitis: 2 (5.3%), Psoriatic Arthritis: 1 (2.6%), others: 6 (15.8%). RT: classic, synthetic and biological disease-modifying antirheumatic drugs, cyclophosphamide and corticosteroids.

Two reactivations were detected with both direct and indirect methods, with a significant title antibody increment and consistent clinical signs and symptoms. Cardiologic abnormalities were found. A 64-years old lady with Microscopic Polyangiitis, under prednisone 60 mg/day and a 57-years old lady with Systemic Lupus Erythematosus under hydroxychloroquine and prednisone 40 mg/day, who happen to be sisters. They did not have other treatments. Both of them developed high fever, myalgias, arthralgias and asthenia. Other infections were ruled out. They received Benznidazole and Nifurtimox respectively. After one-week treatment, Strout and PCR became negative and antibodies decreased, with remarkable clinical improvement.

Conclusion: Two patients with parasitemia and concomitant symptoms were detected, after high doses of corticosteroids. Because they were sisters, a genetic background would also play a role. Parasite's screening before starting immunosuppression and its follow-up during treatment in search of reactivation should be integrated to rheumatology daily practice in endemic countries. It should also be taken into account in patients from non-endemic countries that have an epidemiological nexus. CD specific antibodies determination, Strout and PCR are economic and practical techniques, although they require qualified personnel and equipment. In areas with high prevalence of CD, the benefits of this measures would outweigh costs. A more advanced study should be performed to extend our knowledge.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3509

AB1100

PREVALENCE OF SARCOPENIA IN ECUADORIAN POPULATION BASED ON SCREENING AND DIAGNOSTIC TOOLS

Genesis Maldonado, Carlos Rios. Espiritu Santo University, Rheumatology, Guayaquil, Ecuador

Background: Sarcopenia is defined as an abnormal loss of muscle mass associated with aging. It has been validated that sarcopenia can predict functional deterioration.

Objectives: The purpose of this study is to determine the diagnosis of sarcopenia by conventional methods and gold standard.

Methods: Prospective observational study of patients attending a rheumatology center, to whom anthropometric and densitometric measurements were taken in order to determine sarcopenia.

Results: A total of 110 patients were studied, from which 91% were women and 9% men. 66% presented sarcopenia by means of the gold standard test (DXA). The mean age was 65.80±9.84 [44-89]. The mean age for women was 66.2±9.9 [44-86] and the average age of menopause was 46.08±8.53 and 63% were found to have sarcopenia. In men, the mean age was 73.3±13.8 [63-89] and 30% had sarcopenia.

Anthropometric measures were determined. The mean body mass index was 26.58±4.67, which corresponds to overweight. Three tests were performed in addition to the gold standard. The average gait speed test was 1.12±0.74 seconds and 45% were positive for sarcopenia. The mean of the muscle strength test (dynamometer) was 19.33± 9.52 and 63% had sarcopenia. The mean of the SARC-F screening questionnaire was 3.02±2.24 and 22% were positive for sarcopenia. According to the measurement of muscle mass by DXA, the mean appendicular skeletal muscle mass (ASMM) was 16.08±3.52, ASMM index 6.77±1.12 and the total skeletal muscle was 21.39±4.68. Men presented a higher muscle mass index compared to women [7.82±1.93 vs. 6.66±0.95]. When grouping patients by BMI, sarcopenia was present in all of the patients with low weight, 88% in the ideal weight group, 67% in the overweight group and 25% in the obesity group. A statistically significant relationship was found between sarcopenia and sex [0.01], weight and height [<0.001], gait test [0.04], muscle strength test [<0.001] and DXA [<0.001]. In addition, the sensitivity and specificity of the tests were determined. SARC-F had a 15% sensitivity and a 9% specificity, with a positive predictive value of 74% and a negative predictive value of 37%. The gait speed test 24% sensitivity and 38% specificity, with a positive predictive value of 57% and negative predictive value of 27%. The muscle strength test had a 78% sensitivity and 65% specificity, with a positive predictive value of 84% and negative predictive value of 96% (Figure 1).

Abstract AB1100 Figure 1. Statistical measures of sarcopenia's diagnostic tools.

Conclusion: This is the first study of determination of Sarcopenia in a Rheumatology Center in the city of Guayaquil. 66% of the studied population presented sarcopenia. It is evident that conventional methods may determine a diagnosis, however, a definitive diagnosis can be established by DXA. Because Sarcopenia is a multidisciplinary entity, more studies are needed to determine the usefulness of the conventional methods.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7374