the concentration of mast cells is increased in the synovium of affected joints. Mast cells significantly influence angiogenesis through the production of proangiogenic cytokines, including ANGPTL4.

Conclusion: Changes of the level of ANGPTL4 in the serum of patients with RA may be a potential biomarker of disease activity, severity of neovascularization, inflammation and development of bone erosion.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7868

AB1119

THE PRESENCE OF SYNOVITIS IS THE MAIN FACTOR INFLUENCING THE DEVELOPMENT OF PAIN SYNDROME IN ARTHRITIS OF THE KNEE JOINT

Ninel Aleksandrova¹, Natalia Nikitina¹, Vladislav Aleksandrov^{1,2}, Ludmila Shilova², Andrey Aleksandrov^{1,2}. ¹Research Institute of Clinical and Experimental

Rheumatology named after A.B. Zborovsky, Volgograd, Russian Federation;

²Volgograd State Medical University, Hospital Therapy, Volgograd, Russian Federation

Background: Dysfunctions and pain syndrome in lesions of the knee joint can significantly discomfort a sick person and lead to a persistent decrease in physical activity and disability. Often pain syndrome precedes radiographic appearance of the structural changes in the joint and is accompanied by an increase in number of different changes in the synovium according to ultrasound investigation.

Objectives: To investigate the clinical significance of ultrasound criteria of changes in the synovial membrane of the knee joint cavity and its role in the assessment of pain in gonarthrosis.

Methods: 30 people aged 30 to 50 years with osteoarthritis of the knee joint were under observation; assessment of the severity of pain in the knee when walking was at least 40 mm on a visual analogue scale (VAS). Ultrasound examination of the knee joint was carried out according to standard procedure using a linear sensor with a frequency of 05.12 MHz in an ultrasonic diagnosis system Accuvix V10 (Samsung Medison, Korea).

Results: The evaluation of ultrasound changes was performed in the upper inversion of the knee joint according to the following criteria: the severity of intraarticular effusion (1), synovial proliferation (2), local vascularization of the synovial membrane by power doppler (3). All patients were divided into three groups, according to the severity of pain in the knee joint: group I (12 people) - 41–59 mm, group II (10 people) - 60–79 mm, group III (8 people) - 80–100 mm on the VAS scale. By comparing changes in the knee joint by ultrasound data in patients of different groups, the following results were obtained: group I: severity of intraarticular effusion - 10 people (minimal changes in 60%, moderate in 20%, expressed in 20%), synovial proliferation - 4 people (moderate changes in 50%, expressed in 50%), local vascularization of synovium - 6 people (minimal changes in 66.7%, moderate in 16.7%, expressed in 16.6%); Group II: severity of intraarticular effusion - 9 people (55.6%, 22.2% and 22.2%), synovial proliferation - 3 people (0%, 33.3% and 66.7%), local vascularization of the synovial membrane - 4 people (25%, 25% and 50%, respectively); group III: severity of intraarticular effusion - 8 people (62.5%, 12.5% and 25%), synovial proliferation - 5 people (20%, 40% and 40%), local vascularization of the synovial membrane - 3 people (per 33.3%, respectively).

Conclusion: The use of ultrasound in the diagnosis of diseases of the knee joints allows to reliably determine the structural and functional changes in all components of the knee joint. The severity of pain in gonarthrosis is most associated with the presence of synovitis in the joint.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7065

AB1120

DIAGNOSTIC ROLE OF NEUROMUSCULAR ULTRASOUND IN CUBITAL TUNNEL SYNDROME

Fatma Ali¹, Mona Nasr¹, Ahmed Hafez², Adham Khalil³. ¹Minia university, Rheumatology and Rehabilitation, Minia, Egypt; ²Minia university, Rheumatology and Rehabilitation, Minia, Egypt; ³New Kasr El Eini Teaching Hospital, Cairo University, Cairo, Egypt

Background: Cubital tunnel syndrome (CuTS) is the second most common compressive neuropathy of the upper limb following carpal tunnel syndrome and is the most common site for entrapment for the ulnar nerve

Objectives: Our aim is to evaluate the role of ultrasonography (US) as a diagnostic tool for Cubital tunnel syndrome (CuTS) in comparison with nerve conduction study (NCS).

Methods: twenty elbows with CuTS and twenty asymptomatic controls were assessed by NCS and underwent ultrasonography of elbows. Data from patients and controls were compared to determine the diagnostic relations in patients with CuTS and the grade of severity

Results: There was a high degree of correlation between NCS of the ulnar nerve, clinical parameters and variable US measurements. The CSA of the ulnar nerve was the most sensitive parameter and a cut-off point of 9.5 mm² behind medial epicondyle was found to be 100% sensitive and 80% specific. The ulnar nerve ratios (UNR) had a diagnostic accuracy of 95% with 85% specificity.

Conclusion: Ultrasonographic measurements of the ulnar nerve CSA and UNR have a comparable diagnostic value as a non-invasive and an alternative modality for the evaluation of CuTS

REFERENCES

- [1] Simon N, Ralph J, Poncelet A, Engstrom J, Chin C, Kliot M. A comparison of ultrasonographic and electrophysiologic 'inching' in ulnar neuropathy at the elbow. *Clin Neurophysiol.* 2015;126(2):391-398.
- [2] Omejec G, Podnar S. Normative values for short segment nerve conduction studies and ultrasonography of the ulnar nerve at the elbow. *Muscle nerve.* 2015;51(3):370-377.
- [3] Yoon J, Walker F, Cartwright M. Ultrasonographic swelling ratio in the diagnosis of ulnar neuropathy at the elbow. *Muscle nerve.* 2008;38(4):1231-1235.
- [4] Bayrak A, Bayrak I, Turker H, Elmali M, Nural M. Ultrasonography in patients with ulnar neuropathy at the elbow: Comparison of cross sectional area and swelling ratio with electrophysiological severity. *Muscle nerve.* 2010;41(5):661-666.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4629

AB1121

EFFICACY AND SAFETY OF ULTRASOUND GUIDED ASPIRATION AND INTRA LESIONAL CORTICOSTEROIDS INJECTION OF RUPTURED BAKER'S CYST

Mohammed A. Mortada¹, Youmna A. Amer². ¹Faculty of medicine, Zagazig University, Rheumatology and Rehabilitation, Zagazig, Egypt; ²Faculty of medicine zagazig university, Rheumatology and rehabilitation, Zagazig, Egypt

Background: Baker's cyst is the most common mass in the popliteal fossa and results from fluid distension of the gastrocnemio-semimembranosus bursa. The most common complication of Baker's cyst is the rupture or dissection of fluid into the adjacent proximal gastrocnemius muscle belly, which results in a pseudothrombophlebitis syndrome mimicking symptoms of DVT.

Treatment of ruptured Baker cysts ranged from conservative management to surgical resection.

Ultrasonographic guided aspiration and corticosteroids injection may be an effective and easy method of management of these cases.

Up to the best of our knowledge, this is the first study to detect the efficacy and safety of ultrasonographic guided aspiration and injections of ruptured Baker cysts.

Objectives: To evaluate the efficacy and safety of ultrasonographic guided aspiration and local corticosteroids of ruptured Baker cysts based on follow-up clinical and sonographic results.

Methods: A retrospective study was conducted on 42 patients (12 males and 30 females, mean age 36 +/- 10 SD years) affected by a ruptured Baker cysts associated to knee joint disorders in the period between January 2013 to January 2019. The diagnosis was done by clinical presentation of acute calf pain, swelling, tenderness at the calf muscles and ultrasonographic evidences of ruptured baker cysts in the form of free fluid collection in the calf connected to a well defined cyst at the back of knee.

All cases were treated by ultrasonographic guided aspiration and intralesional injection of corticosteroids once or twice one week apart. Follow up were done on a weekly basis until complete resolution of symptoms then 3 months later. Visual analogue scale (VAS) for calf pain and Rauschnig-Lindgren and Lysholm Knee Scoring Scales were used to assess pre/post-injection knee functions.

Results: The primary diagnoses to patients presented with ruptured Baker cyst in this study were as follow: 18 (42.8%) cases with rheumatoid

arthritis, 15 (37.5%) cases with osteoarthritis and 9 (21.4%) cases with psoriatic arthritis.

Clinical parameters (VAS for calf pain and Rauschnig-Lindgren score) improved significantly in all patients at both post injection evaluation visits.

Rauschnig-Lindgren score was significantly lower after US guided injections (mean, 0; range, 0-1) than at baseline (mean, 2; range, 1-2); $p < 0.001$ (table1).

VAS for calf pain also significantly lower after US guided injections (mean, 0.5; range, 0-1) than at baseline (mean, 9.5; range, 0-10); $p < 0.001$.

Ultrasonographic features improved significantly with complete disappearance of free fluid at the calf in 35 (83.3%) cases one week after the injection.

As regards Baker cyst only 5 (11%) cases showed complete disappearance of the backer cyst and in the majority of cases 37 (88%) there were persistent Baker cysts.

No side effects were reported in all cases.

Abstract AB1121 Table 1. Clinical results of Baker cyst excision (Rauschnig and Lindgren scale and VAS for calf pain)

	Pre injection	1 week later	3 months later
grade 0	0	30	39
Grade 1	0	8	3
Grade 2	6	4	0
Grade3	36	0	0
VAS for calf pain	9.5	1.5	0.5

VAS; visual analogue scale

Conclusion: Ultrasonographic guided aspiration and intra_lesional corticosteroids injection is an effective and safe method in management of ruptured Baker cysts.

REFERENCES

- [1] Nigerian Journal of Orthopaedics and Trauma June 2009; 8 (1): 31-32

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3649

AB1122 PREDICTION OF THE UTILITY OF ULTRASOUND EVALUATION IN RHEUMATOID ARTHRITIS BY THE CLINICAL FACTORS

Takahiro Asano, Masei Suda, Ryo Rokutanda, Haruyuki Yanaoka, Sho Fukui, Haruki Sawada, Yukihiko Ikeda, Ayako Koido, Rui Imai, Hisanori Shimizu, Hiromichi Tamaki, Tokutarou Tsuda, Mitsumasa Kishimoto, Kenichi Yamaguchi, Masato Okada. *St. Luke's International Hospital, Immuno-Rheumatology Center, Chūō-ku, Tokyo, Japan*

Background: It is known that ultrasound detects synovitis with higher sensitivity than physical examination in rheumatoid arthritis (RA), and the residual inflammation detected by ultrasound even in the patient achieving clinically remission well correlates with progression of erosion.¹ However, it is impossible to perform ultrasound evaluation on all the joints in all the patients in terms of time and human resource.

Therefore, ultrasound use in RA should be more focused on situations when the physical findings and ultrasound findings likely to dissociate. Hirata et al showed the concordance between swelling or tenderness joint and ultrasound-detected synovitis are low in the PIP, MCP, and wrist joint.²

Objectives: To investigate the factors that can predict dissociation between clinical and ultrasound-detected synovitis in the patient with rheumatoid arthritis.

Methods: Patients aged 18 years or older who satisfied ACR/EULAR 2010 criteria who visited St. Luke's International Hospital from June 1, 2018 to November 9, 2018 and were performed ultrasound on PIP, MCP, and wrist joints were eligible for this study. We retrospectively reviewed the electronic medical chart and ultrasound data, and analyzed the dissociation of clinical synovitis defined as existence of swelling or tenderness and ultrasound-detected synovitis defined as both of power doppler grade and gray scale grade 1 or gray scale grade 2. Clinical evaluation was done by the attending physician who ordered the ultrasound evaluation. Ultrasound evaluation was done using standard scan technic³ by the sonographers who are blinded to the clinical evaluation.

The correlation between patient's characteristics and dissociation was analyzed using χ^2 test, Mann-Whitney U test and logistic regression analysis.

Results: 51 joints in 32 patients (4 males, 28 females) were included in the study. Among the 32 patients, 18 was classified in the dissociation group (DG) and 14 in the non-dissociated group (NDG). No significant difference was observed in the patient's characteristics of DG and NDG. In the univariate analysis, the absence of swelling in the target joint had tendency to be the risk of dissociation. The same tendency was preserved in the multivariate analysis with the presence of tenderness in the target joint assigned as a covariate (OR 3.68, 95% CI = 0.85 - 15.90, $p = .08$).

Conclusion: There was a numerical tendency that no swelling on the joint is a risk factor for the dissociation between clinical synovitis and ultrasound-detected synovitis. Future study with increased number of participants may identify more sensitive risk factors for dissociation, and enable to make an effective indication of ultrasound evaluation in the care of RA.

REFERENCES

- [1] Kawashiri S, et al. *Rheumatology* 53.3, 2013: 562-569.
[2] Hirata A, et al. *Arthritis Care & Research* Vol. 69, No. 6, June 2017: 801-806
[3] Naredo E, et al. *Annals of the rheumatic diseases* 64.3, 2005: 375-381.

Disclosure of Interests: Takahiro Asano: None declared, Masei Suda: None declared, Ryo Rokutanda: None declared, Haruyuki Yanaoka: None declared, Sho Fukui: None declared, Haruki Sawada: None declared, Yukihiko Ikeda: None declared, Ayako Koido: None declared, Rui Imai: None declared, Hisanori Shimizu: None declared, Hiromichi Tamaki: None declared, Tokutarou Tsuda: None declared, Mitsumasa Kishimoto Consultant for: Kyowa Hakko Kirin Co., Ltd., Kenichi Yamaguchi: None declared, Masato Okada: None declared

DOI: 10.1136/annrheumdis-2019-eular.5472

AB1123 SHOULDER PAIN: ARE THERE PREDICTIVE FACTORS OF RESPONSE TO TREATMENT AND OF ULTRASOUND FINDINGS?

Soraia Azevedo¹, Joana Rodrigues¹, Daniela Faria¹, Joana Neves², Joana Silva¹, Daniela Peixoto¹, Sérgio Alcino¹, José Tavares-Costa¹, Carmo Afonso¹, Filipa Teixeira¹. ¹*Unidade Local de Saúde do Alto Minho, Rheumatology department, Ponte de Lima, Portugal*; ²*Hospital de Braga, Rheumatology department, Braga, Portugal*

Background: Shoulder pain is a common cause of consultation in Primary Health Care, and may correspond to up to 30% of the reasons for consultation. Pathology of the rotator cuff is the most common cause of pain. Ultrasound is a valuable diagnostic tool in assessing shoulder disorders; it can be as effective as magnetic resonance imaging.

Objectives: To determine the predictive factors of response to treatment and ultrasound findings in shoulder pain.

Methods: We performed an analysis of the patients' cases sent to the rheumatology consultation with shoulder pain, every patients had an echography shoulder evaluation, and the rheumatologist decided treatment based on the guidelines for the treatment of shoulder tendinopathies. The use of nonsteroidal anti-inflammatory drug (NSAIDs) and muscle relaxant medications as well as the following techniques: corticosteroids local injection, barbotage, capsular distension and physiotherapy programs were some of the variables assessed. Posteriorly, the patients were clinically assessed in a follow-up visit.

Results: A total of 119 patients were evaluated. Men presented a higher risk of distension of the subscapular bursa. There was a statistically significant relationship between the time from the beginning of the symptomatology and treatment response. Diabetes mellitus, arterial hypertension and dyslipidaemia were statistically significantly associated with some rotator cuff lesions and distention of the subscapular bursa.

Conclusion: In patients with shoulder pain, early intervention positively influences the response to treatment. Thus, it is important that these patients are evaluated more promptly.

Some comorbidities seem to be associated with a higher risk of specific rotator cuff lesions.

No relationship was found between response to treatment and age, sex, occupation, previous treatments or type of therapy selected. The associations found in this study seem to have clinical implications. Prevention of rotator cuff disease is a matter of major relevance as well as early institution of treatment.