Carestream, UCB, Abbvie, Pfizer, Astra Zeneka, Roche, Siemens, Image Analysis Group, Jonathan Vela: None declared, Soeren Skou: None declared, Mikkel Eld: None declared, Nasir Musa Al-Mashkur: None declared, Robert Riis Consultant for: Image analysis Group, Ole Simon-sen: None declared

DOI: 10.1136/annrheumdis-2019-eular.5695

AB1126

ULTRASOUND CHANGES IN PATIENTS WITH CALCIFICATIONS IN THE SUPERSPINATUS TENDON BEFORE AND AFTER THE DEVELOPMENT OF A SUBACROMIAL PAINFUL SYNDROME

Carlos Guillén-Astete, Andrea Briones-Figueroa, Ramon y Cajal University Hospital, Madrid, Spain

Background: The relationship between subacromial syndrome (SS) and the presence of calcifications in the thickness of the superspinatus tendon is complex. It is undeniable that the presence of a solid body hinders the movement of the rotator cuff through the subacromial space and to some extent contributes to the trapping that conditions pain and functional limitation, but, on the other hand, calcifications in the thickness of the superspinatus tendon are incidental findings that have no clinical significance or whose presence precedes the painful syndrome in months or years. We understand that in a SS related to calcification, it contributes as an irreducible agent that conditions an inflammatory response of the tendon when it attempts to cross the subacromial space, however, there are no studies available that explain this relationship from a sonoanatomic perspective.

Objectives: To determine the extent to which the disposition and characteristics of a calcification of the supraspinatus tendon, or of the tendon itself, are modified when an SS is triggered.

Methods: A quasi-experimental, "before-after" study was conducted with individuals who presented a SS and in whom incidental calcification in the TSE had been previously diagnosed in an ultrasound study. Demographic data and static images from the ultrasound studies of each patient were compared. The explanatory variable was time (before-after) and the response variables in image were: (1) Increased tendon thickness measured at 30mm from the deep edge of the acromion (GROSORB), (2) Increased tendon thickness measured at the height of the middle of the longitudinal axis of the calcification (GROSORC), (3) Longitudinal axis of the calcification (LONGC), (4) Thickness or height of the calcification, when this was measurable (ALTC), and (5) Distance between the calcification and the deep end of the acromion (DISTC). In addition, two indices were determined: (a) Increased tendon thickness in the calcification zone (GROSORC/GROSORB) and (b) proportion of tendon thickness occupied by calcification (ALTC/GROSORC). The measurements were carried out using the software of the equipment in which the original and subsequent study was made (GE Logiq e; Toshiba Nemio XG).

Variables measured by ultrasound assessment	Before measurements	After measurements	T Student (paired data)
ALTC (N=20) mm	5.075 DE 1.055	5.160 DE 0.994	0.0073
LONGC mm	6.184 DE 2.414	6.236 DE 2.446	0.1066
GROSORB mm	6.348 DE 0.691	6.436 DE 0.674	0.1285
GROSORC mm	6.845 DE 0.718	7.263 DE 0.872	0.0003
GROSORC/GROSORB	1.082 DE 0.085	1.129 DE 0.069	0.0072
ALTC/GROSORC	0.748 DE 0.164	0.716 DE 0.130	0.0611

Results: The "before" ultrasounds were performed between 2012 and 2016 and the "after" ultrasounds between 2015 and 2018. The time fashion between the two ultrasounds for each patient was 2 years. Thirty-three subjects were included, with an average age of 47.5 SD 7.6 years. Proportion of women 66.6%. Proportion of dominant shoulders (51.1%). As shown in table 1, the most significant changes were the thickness of the tendon measured at the height of the middle of the longitudinal axis of calcification and the increase of this measure with respect to the thickness measured at 30mm from the deepest region of acromion (GROSORC/GROSORB). Although the calcification height also showed a statistically significant increase, it could only be measured in 20 of the 33 subjects.

In the correlation tests, the GROSORC/GROSORB index increase correlated strongly with the DISTC (Coef. Pearson 0.686, $P < 0.01$) and with the longitudinal diameter of the calcification (LONGC) (Coef. Pearson 0.797, $P < 0.01$).

Conclusion: The most important factor that explains the development of an SS associated with calcification is the increase in tendon thickness at the height of the calcification itself and, at the same time, this measure increases when the calcification is more lateralized and the more elongated its longitudinal diameter. These modifications could have implications when deciding the therapeutic management of these processes.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.5918

AB1127

ULTRASOUND INTER-READER RELIABILITY OF INFLAMMATORY FINDINGS IN PATIENTS WITH POLYARTHRITIS

Luisa Brites¹, Mariana Santiago^{1,2}, João Rovisco^{1,2}, João Dinis de Freitas¹, José Antonio P. Da Silva^{1,2}, Inês Silva³, Filipa Teixeira⁴, Daniela Faria⁴, Cândida G. Silva^{5,6}, Sandra Falcão^{3,7}. ¹Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal; ²Faculdade de Medicina de Coimbra, Coimbra, Portugal; ³Hospital Egas Moniz, Centro Hospitalar de Lisboa Ocidental, Lisbon, Portugal; ⁴Unidade Local de Saúde do Alto Minho, Serviço de Reumatologia, Ponte de Lima, Portugal; ⁵Higher School of Health Sciences, Polytechnic Institute of Leiria, Leiria, Portugal; ⁶Centro de Química da Universidade de Coimbra, Coimbra, Portugal; ⁷Faculdade de Ciências Médicas da Universidade Nova de Lisboa, Lisbon, Portugal

Background: Ultrasonography is an imaging technique that allows rheumatologists to visualise structural and inflammatory changes within a joint.

Objectives: The objective of this study was to assess the inter-reader reliability of interpretation of inflammatory and destructive changes in a wide range of joints in patients with polyarthritis.

Methods: This study was divided in two parts: 1) consensus process and 2) reliability exercise. For the first part, a written questionnaire was sent by email to 6 sonographers from 3 portuguese hospitals with the highest level of competence (EULAR competency assessment level 2). The questionnaire included 17 questions divided in two groups: 1) elementary components in B-mode and Doppler assessment (effusion, synovial hypertrophy (SH), power Doppler (PD), erosions and synovitis definition) and 2) approach at the joint level (the definition of which plan and recess will be assessed in each joint). The participants were asked to rate their level of agreement/disagreement for each statement using a 1-5 Likert scale (1=strongly disagree to 5=strongly agree). For the reliability exercise, video clips of US examinations of 40 joints (wrist, metacarpophalangeal (MCP) from 1 to 5, proximal interphalangeal (PIP) from 1 to 5, knee, tibiotarsal (TT) and metatarsophalangeal (MTP) joints from 1 to 5, elbow and shoulder) from each of 15 patients were collected (showing a multiplanar bilateral ultrasound approach). Each joint in each video was scored by individual ultrasonographers for the presence/absence of elementary components: effusion (Yes/No), SH (No/Grade 1 to 3), PD (No/Grade 1 to 3) and erosions (Yes/No). Inter-reader agreement analysis was assessed through Fleiss' kappa coefficient and classified according to Landis and Koch[8]: κ values 0 were considered poor, 0-0.20 slight, 0.21-0.40 fair, 0.41-0.60 moderate, 0.61-0.80 good and 0.81-1.00 excellent. Statistical significance was defined as $p < 0.05$. Statistical analysis was performed using STATA V.14.

Results: Thirty seven joints of the 600 joints were excluded due to dislocation of the joint or presence of objects (rings/catheters) and the videos of a total of 563 joints were analysed by the 6 ultrasound experts. Inter-reader agreement was superior for TT joints and inferior for wrist; the identification of the erosions had the better agreement in the elementary components (Table 1).

Table 1. Inter-observer agreement for each elementary component and for anatomical region

Elementary component	κ	Joint	κ	Joint	κ
Effusion	0.6044	Wrist	0.6767	Shoulder	0.7271
Synovial	0.6291	MCP	0.6866	Knee	0.7192
hypertrophy					
Power doppler	0.7195	PIP	0.7107	TT	0.8043
Erosions	0.7314	Elbow	0.7291	MTP	0.7040

MCP - Metacarpophalangeal joints, MTP - Metatarsophalangeal joints, PIP - Proximal interphalangeal joints, TT-Tibiotarsal joint. All $p < 0.001$

Conclusion: The reliability of interpretation of inflammatory and destructive changes using video clips was in general good to excellent and it was better for erosions and tibiotarsal joint (regarding elementary component and anatomical region, respectively).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.2181

AB1128

SENSITIVITY AND SPECIFICITY OF THE AUTOMATED SQUEEZE TEST (GAENSLER'S MANEUVER) FOR IDENTIFYING METACARPOPHALANGEAL SYNOVITIS BY MAGNETIC RESONANCE IMAGING

Diana Daniela Castañeda Martínez, David Vega Morales, Jorge Antonio Esquivel Valerio, Mario Alberto Garza Elizondo, Brenda Roxana Vázquez Fuentes, Dionicio Ángel Galarza-Delgado. Hospital Universitario "Dr. José Eleuterio González" Universidad Autónoma de Nuevo León, Servicio de Reumatología, Monterrey, Mexico

Background: It is well known that an early diagnosis and treatment for Rheumatoid Arthritis (RA) prevents its complications, therefore there are many efforts to identify individuals at risk to develop RA. The Squeeze test has been used to detect synovitis in metacarpophalangeal joints, even though it is used in daily practice, there is a great variability in their performance among rheumatologists. [2]

Objectives: The aim of this study is to determine the diagnostic performance of the automated squeeze test (AST) on the metacarpophalangeal (MCP) joints to detect the presence of synovitis, edema, or erosions by magnetic resonance imaging (MRI) using the rheumatoid arthritis magnetic resonance imaging score (RAMRIS) in first-degree relatives (FDR) of RA patients, of whom clinically suspect arthralgia (CSA) in hands was suspected, as well as in RA patients.

Methods: It is an observational and cross-sectional study for a diagnostic test. A total of 60 patients, older than 18 years, were included and divided in three groups: CSA group: 22 with less than 1 year with arthralgia and required to be FDR of RA patients; early RA group: 22 patients who met ACR/EULAR 2010 Classification Criteria with less than 1 year with the disease; and late RA group: 16 patients who met ACR/EULAR 2010 Classification Criteria, with more than 1 year with the disease. The AST was performed in the 60 participants' dominant hand. The device was evaluated by MRI, which examined the same hand in the 60 patients.

Results: A total of 240 MCP joints were evaluated. The AUC for the total RAMRIS score >10 was [0.480 (95% CI 0.301-0.617) $P=0.597$], for synovitis RAMRIS score > 7 was [0.459 (95% CI 0.331-0.669) $P=0.791$], and for the presence of any synovitis by RAMRIS was [0.575 (95% CI 0.428-0.723) $P=0.331$]. For the RAMRIS synovitis score, the most sensitive and specific cut-off of the force by AST was 4.645 kg with a 66.7% sensitivity and 50% specificity.

Abstract AB1128 Table 1. Baseline characteristics

	TOTAL	CSA n=22	ERA n=22	LRA n=16
Age mean (SD)	44.7 (13.7)	37.6 (10.9)	49.05 (11.9)	47.5 (16.2)
Female n (%)	50 (83.3)	18 (81.8)	18 (81.8)	14 (87.5)
TJC median (IQR)	4 (12)	0.5 (4)	5 (13)	10 (11)
SJC median (IQR)	3 (8)	0 (3)	5 (12)	7.5 (9)
DAS28-ESR median (IQR)			4.8 (1.5)	5.4 (1.4)
HAQ median (IQR)			0.64 (0.75)	1.47 (0.94)
CDAI median (IQR)			18.07 (16.8)	23.8 (15.5)
Married n (%)	41 (68.3)	12 (54.5)	15 (71.4)	14 (93.3)
Smoker n (%)	19 (31.7)	2 (22.7)	10 (45.5)	4 (25)
Morning stiffness n (%)	31 (51.7)	4 (18.2)	16 (72.7)	11 (68.8)
RF IgM positivity n (%)	46 (76.7)	9 (40.9)	12 (70.6)	4 (25)
ACPA positivity n (%)	25 (41.7)	19 (86.4)	20 (90.9)	7 (43.8)

Clinically suspect arthralgia, **CSA**; Early Rheumatoid Arthritis, **ERA**; Late Rheumatoid Arthritis, **LRA**; Tender Joint Count, **TJC**; Swollen Joint Count, **SJC**; Disease Activity Score – Erythrocyte Sedimentation Rate, **DAS28-ESR**; Health Activity Questionnaire, **HAQ**; Clinical Disease Activity Index, **CDAI**; Rheumatoid Factor, **RF**; Anti-citrullinated peptides antibodies, **ACPA**.

Figure 1

Conclusion: The application of AST with a force of 4.645 kg, to exert pain in the dominant hand, is a sensitive and specific technique to establish MCP joint inflammation, compared with other automated methods that had been recently tested.

REFERENCES

- [1] van Steenberg HW, Aletaha D, Beart-van de Voorde LJ, et al. EULAR definition of arthralgia suspicious for progression to rheumatoid arthritis. Ann Rheum Dis 2017;76:491-6.

- [2] Vega-Morales D, Esquivel-Valerio JA, Arana-Guajardo AC. Automated squeeze test (Gaensler's manoeuvre) to identify patients with arthralgia suspicious for progression to RA: improving time delay to rheumatology consultation. Ann Rheum Dis 2017;76:e40.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4182

AB1129

ULTRASOUND MEASUREMENT OF MUSCLE THICKNESS AT THE PROXIMAL FOREARM: VALIDITY ISSUES

Sonia Castell¹, Gianluca Smerilli², Edoardo Cipolletta², Fausto Salaffi², Emilio Filippucci², Walter Grassi². ¹Hospital del Mar, Rheumatology, Barcelona, Spain; ²Università Politecnica delle Marche, Rheumatology Unit, Ancona, Italy

Background: The term *sarcopenia* defines the muscle strength loss and muscle mass loss due to aging, and it is one out of three criteria for the diagnosis of frailty in the elderly. A number of studies have related the handgrip strength with muscle thickness (MT) in the forearm measured by ultrasound (US)¹, but a standardized scanning protocol has not been described and its interobserver reliability has not been investigated yet.

Objectives: The aims of this study were to provide detailed description of the scanning protocol to measure the MT in the forearm and to test its feasibility and interobserver reliability.

Methods: A total of 27 consecutive subjects were enrolled at Ospedale "Carlo Urbani", in Jesi (Ancona, Italy): 2 healthy volunteers recruited from among our staff, and 25 patients referred to the Rheumatology Department affected by rheumatoid arthritis (5), psoriatic arthritis (4), spondyloarthritis (3), polymyalgia rheumatica (2), systemic lupus erythematosus (1), systemic sclerosis (1), overlap SLE/SSc (1), Sjögren syndrome (1), anti-synthetase syndrome (1), undifferentiated connective tissue disease (1), osteoporosis (1), small vessels vasculitides (2) and fibromyalgia (2). The female to male ratio was 19/27, the mean age was 52.1 years (SD ± 13.7), and the mean Body Mass Index was 27.4 Kg/m² (SD ± 4.2). Four rheumatologists (SC, EC, GS, EF) trained in musculoskeletal US, with a different degree of experience, performed the examinations using a MyLab ClassC (Esaote SpA) equipped with a broadband linear probe (frequency range 4-13 MHz). All subjects sat in front of the sonographer with their hands supinated and the forearm resting on the examining table. First, the coronoid process was imaged according to the "longitudinal scan of the coronoid recess" as indicated by the 2017 EULAR US guidelines. Then the probe was moved distally following bony cortex until the ulnar tuberosity was identified. Immediately distally to the ulnar tuberosity the bone turns flat and hyperechoic and this was taken as the anatomical reference for the measurement. Afterwards the probe orientation was changed to obtain a transverse view. During the rotation, the proximal third of the diaphysis of the radius was imaged. Two MT were measured, the ulnar MT (UMT) and the radial MT (RMT), between the subcutaneous tissue-muscle interface and the muscle-bone interface of each bone respectively. The measurement of UMT and RMT of both arms were registered, as well as the scanning time of all the examiners.

Results: We found an excellent interobserver reliability of this scanning protocol, with an interclass correlation coefficient (ICC) among the four sonographers of 0.975 (CI 0.955 - 0.987) for the right UMT, an ICC of 0.968 (CI 0.942 - 0.984) for the left UMT, an ICC of 0.932 (CI 0.878 - 0.966) for the right RMT and an ICC of 0.949 (CI 0.908 - 0.974) for the left RMT. The mean time required to acquire all measurements in each subject was less than five minutes (SC 4.4 min; EC 4 min; GS 4.2 min; EF 4.5 min).

Conclusion: The results of this study provide evidence in favour of both feasibility and interobserver reliability of US measurement of the forearm MT.

REFERENCES

- [1] Abe T, Thiebaux R S, Loenneke J P, Ogawa M and Mitsukawa N. Association between forearm muscle thickness and age-related loss of skeletal muscle mass, handgrip and knee extension strength and walking performance in old men and women: a pilot study. Ultrasound in Med. & Biol., Vol. 40, No. 9, pp 2069-2075, 2014.

Disclosure of Interests: Sonia Castell: None declared, Gianluca Smerilli: None declared, Edoardo Cipolletta: None declared, Fausto Salaffi: Grant/research support from: Abbvie, Roche, Novartis, BMS, Pfizer, Sanofi, Speakers bureau: Abbvie, Roche, Novartis, Pfizer, Sanofi, BMS, Emilio Filippucci: None declared, Walter Grassi: None declared