AB1234

OUTCOME OF EDUCATION WITH REGARD TO INFLUENZA AND PNEUMOCOCCAL VACCINATIONS IN INFLAMMATORY ARTHRITIS PATIENTS ON DMARDS

Michela Frendo, *Ntoriana Tziatzou*, Daniel Debattista, Thomas Degiorgio. *Mater Dei Hospital, Rheumatology, Tal-Qroqq, Malta*

Background: Inflammatory arthritis (IA) patients on immunosuppressant disease modifying drugs (DMARDs) are at an increased risk of infections. Influenza and pneumococcal vaccines are recommended as part of the BSR and EULAR guidelines for the clinical management of these patients. Prior to commencing DMARDs, the patients are reviewed by the nurse specialist, who discusses the benefits versus risks of DMARDs, necessary monitoring and recommends the pneumococcal and influenza vaccines.

Objectives: The aim of this audit is to assess the uptake of the pneumococcal and the influenza vaccine in IA patients prior to starting biologic or synthetic DMARDs as advised by the nurse specialist during the education visit with the patient.

Methods: The study sample included 139 patients with various types of IA, including rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis, who attended the rheumatology nurse education sessions prior to starting DMARDs in a secondary care hospital in 2017. Verbal advice supported by a vaccination leaflet developed in 2016 was given by the rheumatology nurse. Data was compiled by means of a telephone questionnaire.

Results: One hundred and twenty six (90.6%) participants recalled being given advice on vaccinations. Seventy eight (62%) of these patients received the influenza vaccine. The rest (28%) did not receive the vaccine for various reasons including fear of side effects, fear of developing a worse infection, belief of inefficacy and fear of injections.

A significant improvement ($p=0.0084$) in the influenza vaccination rates was noted since a previous audit in 2016, where following verbal education by a rheumatologist, only 41.4% received the influenza vaccine.

A significant improvement in uptake was also noted in the pneumococcal vaccination rates since only 17.2% of the patients received the pneumococcal vaccine in 2016 compared to 62.7% in 2017 ($p<0.0001$). Various reasons including fear of side effects, belief of inefficacy, fear of injections and financial implications were given by patients who did not receive the pneumococcal vaccine.

Overall, 62% of the patients received both vaccines after education given by the rheumatology specialist nurse and receiving the vaccination leaflet.

Conclusion: This audit showed a significant progress in the uptake of the influenza and pneumococcal vaccinations in patients with inflammatory arthritis following verbal advice by the specialist rheumatology nurse and the introduction of a vaccinations' educational leaflet.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.872

AB1235

COST-SAVING FOR HEALTH SERVICES DETECTING THE MISDIAGNOSIS OF RHEUMATOID ARTHRITIS USING IMAGING IN THE PROCESS OF DIAGNOSIS: EVIDENCE FROM REAL-WORLD

Omaira Valencia¹, Pedro Santos-Moreno², Edwin Castillo³, Nelson Alvis⁴. ¹*Biomab Center for Rheumatoid Arthritis, Research, Bogota, Colombia*; ²*Biomab Center for Rheumatoid Arthritis, Scientific direction, Bogotá, Colombia*; ³*Biomab Center for Rheumatoid Arthritis, Rheumatology, Bogotá D.C., Colombia*; ⁴*Alzack Foundation, Health Economics, Bogotá, Colombia*

Background: The diagnosis of rheumatoid arthritis (RA) using EULAR criteria through conventional assessments remains controversial, especially for those with seronegative results, many of patients without other diagnostic aids as imaging, could be wrong diagnosed followed by expensive treatments (1-3).

Objectives: To evidence through real-world data how the use of imaging within a screening process of diagnosis of RA can saving future costs for unnecessary treatments in patients with misdiagnosis of RA.

Methods: A retrospective real-world data (RWD) analysis was developed from medical records of patients with presumptive RA diagnosis reportedly seronegative for both rheumatoid factor (RF) and anti-cyclic citrullinated peptide antibodies (ACPA), and who met ACR/EULAR 2010 classification criteria, in the period between July 2016 and June of 2017; patients were assessed by imagenology (X-ray, ultrasound (US) or magnetic resonance imaging (MRI) according to the screening diagnosis protocol in a center of integral attention for rheumatoid arthritis (CIA-RA) in order to confirm diagnosis of RA, or classify patients in an alternative proper diagnosis. Direct costs of diagnosis was estimated in two scenarios: the conventional diagnosis and the screening process of diagnosis in the CIA-

RA. To quantify the cost-savings for this process we also estimated the cost of treatment for the first year after diagnosis for patients with RA and patients with the most common diagnosis found after screening.

Results: 440 patients were referred to our center with presumptive diagnosis of RA in the period, who were assessed for ACPA and RF obtaining a seronegative result for both. After screening process just 115 patients were classified as RA, 99 as SRA and 16 as Nonspecific RA; 12.2% were identified by X-Ray, 67.7% were identified by US and 20% by MRI. The most frequently misdiagnosis found was Osteoarthritis in the 72.5% of patients assessed by the screening process. In that way, the conventional diagnosis cost \$54.4, while the CIA-RA screening diagnosis cost was \$247.1 per patient, however there was found a potential cost-saving from using the CIA-RA screening process of diagnosis of \$1,440,494 per year due to the pharmacological cost saving of 325 patients who requires treatment for OA and not for RA.

Conclusion: According with our findings the use of imaging within a diagnostic screening process combining conventional criteria is a useful tool to discard false positive diagnosis of RA. Despite the fact that at first sight, the cost of screening process of diagnosis is more expensive than conventional diagnosis, after one year of treatment it can be assumed potential cost-savings using the proposed approach.

REFERENCES

- [1] Hu H, Luan L, Yang K, Li S-C. Burden of rheumatoid arthritis from a societal perspective: A prevalence-based study on cost of illness for patients with rheumatoid arthritis in China. *Int J Rheum Dis* [Internet]. 2017 Feb 17 [cited 2017 Mar 1]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28211251>
- [2] Cross M, Smith E, Hoy D, Carmona L, Wolfe F, Vos T, et al. The global burden of rheumatoid arthritis: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis*. 2014;73(6).
- [3] Østergaard M, Pedersen SJ, Døhn UM. Imaging in rheumatoid arthritis - status and recent advances for magnetic resonance imaging, ultrasonography, computed tomography and conventional radiography. *Best Pract Res Clin Rheumatol* [Internet]. 2008;22(6):1019-44. Available from: <http://dx.doi.org/10.1016/j.berh.2008.09.014>

Disclosure of Interests: Omaira Valencia: None declared, Pedro Santos-Moreno Grant/research support from: Dr Santos has received research grants from Janssen, Abbvie and UCB, Speakers bureau: Dr Santos has received speaker fees from Sanofi, Lilly, Bristol, Pfizer, Abbvie, Janssen and UCB, Edwin Castillo: None declared, Nelson Alvis : None declared

DOI: 10.1136/annrheumdis-2019-eular.7623

AB1236

IMPROVING CLINICAL OUTCOME AND REDUCING COST FOR PATIENTS WITH RHEUMATIC DISEASES VIA ONLINE INTERACTION WITH RHEUMATOLOGISTS BASED ON SMART SYSTEM OF DISEASE MANAGEMENT (SSDM) MOBILE TOOL

Yong Wang¹, Li Luo², Qin Li³, Fen Wang⁴, Anbin Huang⁵, Hong Zhang³, Peng Ji², Yanping Zhao⁶, Lingxun Shen⁵, Zhengang Wang⁷, Feng Wei⁸, Hui Xiao⁹, Yuhua Jia⁹, Fei Xiao⁹, Fengchun Zhang¹⁰, SSDM Collaboration Group, China.

¹The first Hospital Affiliated to AMU (Southernwest Hospital), Chongqing, China;

²The first affiliated hospital of xinjiang medical university, Urumchi, China; ³The

First People's Hospital of Yunnan Province, Kunming, China; ⁴First Affiliated

Hospital of Medical University Of Anhui, Hefei, China; ⁵Union Hospital Affiliated

Tongji Medical College of Huazhong University of Science and Technology,

Wuhan, China; ⁶First Affiliated Hospital of Harbin Medical University, Harbin, China;

⁷TongRen hospital, Capital University, Beijing, China; ⁸Jiangmen Central Hospital

Affiliated Jiangmen Hospital of Sun Yat-Sen University, Jiangmen, China;

⁹Shanghai Gothic Internet Technology Co., Ltd., Shanghai, China; ¹⁰Peking Union

Medical College hospital, Beijing, China

Background: Without efficient primary medical care and follow-up system in China, patients can choose any hospitals or doctors they like in seeking care. As a result, most patients rush to large hospitals. Once patients left those clinics, no follow up data is available. Surveys show that over 40% of the rheumatic patients don't need to go to a hospital, only need advice from rheumatologists. SSDM is a series of applications for chronic diseases management, which strengthens the interaction between doctors and patients based on valuable clinical data. Our previous study showed that patients can master the SSDM and perform self-management after training, including evaluations on disease activity and health assessment questionnaire (HAQ), as well as medication and lab test data entries.