AB1286

EVALUATION OF CARDIOVASCULAR RISK FACTORS AND PREVALENCE OF METABOLIC SYNDROME IN PATIENTS WITH DIAGNOSIS OF ANKYLOSING SPONDYLITIS:

Severino Correia Do Prado Neto¹, Érika Carvalho de Aquino², Jozelia Rêgo¹, Nilzio Da Silva¹, Fabiana Pina¹. ¹Hospital das Clínicas de Goiânia – Universidade Federal de Goiás, Goiânia, Brazil; ²Universidade Federal de Goiás, Goiânia, Brazil

Background: Cardiovascular events are the main causes of mortality in patients with Ankylosing Spondylitis. In addition, a higher prevalence of Metabolic Syndrome is reported in this group. With immunobiological therapy, much progress has been made in controlling the inflammatory process, but the cardiovascular risk remains high.

Objectives: To evaluate the prevalence of cardiovascular risk factors and Metabolic Syndrome in patients with Ankylosing Spondylitis at the Rheumatology Outpatient Clinic of HC-UFG and to correlate them with epidemiological, clinical, laboratory and radiographic characteristics of the disease.

Methods: Data from 59 patients were collected in medical records between July and August 2018. Clinical characteristics, cardiovascular risk factors and metabolic components were analyzed. Descriptive analyzes of the data were made, prevalences and their reasons were calculated. The associations between variables were assessed by chi-square test and Fisher's exact test.

Results: 81.4% were male, 67.8% self-denominated non-whites, average age 46.7 years. Isolated axial joint involvement was the most frequently observed (54.2%). Enthesopathic lesions were identified in 30.51% of the cases. Uveitis had a higher prevalence (22.02%), 4.2 times higher in the subgroup with HLA-B27 positive ($p=0.0002$). HLA-B27 was present in 67.8%. Advanced sacroiliitis (grades 3 and 4) and syndesmophytosis were identified in 61.1% and 49.2%, respectively, but didn't presented significant correlation with the presence of HLA-B27. 64.4% were using anti-TNF alone or in combination. The majority of the patients (72.9%) were sedentary. Clinical comorbidities were reported in 66.1% of the cases, with dyslipidemia (45.8%), systemic arterial hypertension (37.6%) and diabetes mellitus (13.6%) being the most common. Metabolic Syndrome was diagnosed in 23.7% of the samples. Normal levels of triglycerides and fasting glycemia and lower prevalence of Metabolic Syndrome had a statistically significant association with the presence of HLA-B27 ($p=0.006$; 0.026 and 0.014, respectively). Patients with HLA-B27 present had 61% lower frequency of Metabolic Syndrome. There was no association between the syndrome and its components with anti-TNF therapy; BASDAI; BASFI; metric evaluation; uveitis; degree of sacroiliitis and syndesmophytosis.

Conclusion: The prevalence of Metabolic Syndrome was below that is described in other populations. The presence of HLA-B27 was considered a protective factor for Metabolic Syndrome, triglyceride levels and fasting glycemia. This finding may be related to the presence of several HLA-B27 alleles, usually non-subtyped, and to the great ethnic miscegenation of the Brazilian population. Our observation suggests that spondylitis should be routinely evaluated for cardiovascular risk factors.

REFERENCES

- [1] Maia, DG, et al. Metabolic Syndrome in patients with ankylosing spondylitis receiving anti-TNF alfa therapy: association with predictors of cardiovascular risk. *Clin Rheumatology* 2011;36(10):2371-76.
- [2] Fiorillo, MT, et al. CD8+ T-cell autoreactivity to an HLA-B27-restricted self-epitope correlates with ankylosing spondylitis. *The Journal of Clinical Investigation* 2000;106(1):47-53.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6987

AB1287

DISEASE EVOLUTION OF PRIMARY SJOGREN'S SYNDROME – A LONGITUDINAL STUDY

Inês Rego de Figueiredo¹, Jessica Tam², Dennis Lendrem², Josephine Vila³, Elizabeth Kidd³, Julie Norris³, Ben Hargreaves³, Wan Fai Ng^{2,3}. ¹Unidade de Doenças Auto-imunes/Medicina 7.2, Hospital de Curry Cabral, Centro Hospitalar Universitário Lisboa Central (CHULC), Lisboa, Portugal; ²Musculoskeletal Research Group, Institute of Cellular Medicine, Newcastle University, Newcastle-Upon-Tyne, United Kingdom; ³Newcastle-upon-Tyne Hospitals NHS Foundation Trust, Newcastle-Upon-Tyne, United Kingdom

Background: The natural history of primary Sjogren's syndrome (pSS) remains poorly understood. It has been suggested that the initial presentations may predict the course of disease (1), with gradual reduction of biological activity as measured

by serum Immunoglobulins levels (2), while symptoms such as fatigue remaining stable over time(3).

Objectives: To evaluate how pSS progresses over time using ESSDAI (Eular Sjögren's Syndrome Disease Activity Index) and the ESSPRI (Eular Sjögren's Syndrome Patient Index), EQ5D-Time trade off (TTO) and Visual Analogue Scale (VAS) of health states.

Methods: Routine clinical data from a large single centre cohort in the UK were analysed on patients with ≥ 3 clinical visits, including up to the 10th visit. Outcomes (ESSDAI/ESSPRI and their component scores, EQ5D-3L and VAS) were analyzed by a random effects linear regression using STATA 14.

Results: 346 patients out of 858 included, with a female preponderance of 89%, and median age of 63 years. The median follow up time was 4.9 years, and median disease duration of 12 years. Anti-Ro was positive in 61% and anti-La in 42% of the patients.

ESSDAI score decreased 0.1 point per visit ($p=0.006$). Anti-Ro and anti-La positive patients exhibiting a lower score ($p=0.012$ and 0.031), and patients with MALT at presentation had a score up to 6 points higher ($p=0.002$). Regarding the ESSDAI domains, the constitutional and haematological domains showed increased activity ($p\text{-value}<0.0001$ and 0.018), with Anti-Ro and/or Anti-La patients having higher scores ($p<0.0001$ and 0.001). In contrast, the glandular, articular and peripheral nervous system domains showed decreases over time ($p\text{-values } 0.003, 0.006$ and <0.0001).

ESSPRI score increased 0.05 point/visit ($p<0.0001$). While Dryness scores remained relatively constant, Pain and Fatigue components of the ESSPRI increased over time (both $p<0.0001$). Dryness scores were higher in female patients ($p=0.04$).

EQ5D-TTO worsened by 0.03 point/visit ($p<0.0001$), but the VAS health states remained stable.

Conclusion: Our data suggest that symptoms of fatigue and pain as well as health utility worsen over time, whereas different ESSDAI domains showed different trends over time. Longer term follow-up to further understand the natural history of pSS is warranted.

REFERENCES

- [1] Skopouli FN, Dafni U, Ioannidis JPA, Moutsopoulos HM. Clinical evolution, and morbidity and mortality of primary Sjogren's syndrome. *Semin Arthritis Rheum*. 2000;29(5):296–304.
- [2] Pertovaara M, Pukkala E, Laippala P, Miettinen A, Pasternack A. A longitudinal cohort study of Finnish patients with primary Sjögren's syndrome: Clinical, immunological, and epidemiological aspects. *Ann Rheum Dis*. 2001;60(5):467–72.
- [3] Haldorsen K, Bjelland I, Bolstad AI, Jonsson R, Gorgas Brun J. A five-year prospective study of fatigue in primary Sjögren's syndrome. *Arthritis Res Ther*. 2011;13.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6189

AB1288

PREVENTION OF VENOUS THROMBOEMBOLISM AND THE RISK OF POSTOPERATIVE COMPLICATIONS IN PATIENTS WITH RHEUMATOID ARTHRITIS AFTER TOTAL HIP ARTHROPLASTY:

Alexey Rybnikov¹, Evgenij Bialik², Tatiana Reshetnyak², Sergey Makarov², Alexander Khramov², Sergey Maglevaniy². ¹Nasonova Research Institute of Rheumatology, Moscow, Russian Federation; ²Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: According to the administrative data, confirmed by several meta-analyses, patients with rheumatoid arthritis (RA) in comparison with the General population shows an increased risk of venous thromboembolic complications more than twice. Drug prevention can reduce the risk of venous thromboembolism (VTE) in patients with rheumatic diseases.

Objectives: The aim of this study was to analyze frequency of VTE, risk of bleeding and complications of the postoperative wound in patients with RA and osteoarthritis (OA) after total hip arthroplasty (THA).

Methods: Study included 486 patients (212 - with RA and 274 - with OA) who underwent primary THA. Each group of patients was divided into 3 subgroups by type of drug therapy (1-nadroparin calcium; 2-dabigatran etexilate; 3-nadroparin calcium with transfer to dabigatran etexilate). Intra- and postoperative blood loss and the wound healing process were assessed during the first 7 days after surgery.

Results: Postoperative VTE were reported in 36 (7.4%) of 486 patients. VTE in patients with RA were detected significantly less frequently than in OA (1.2% and 6.1%, $p = 0.0013$). Bleeding that required transfusion