Results: Among 64 patients included a musculoskeletal syndrome was detected in 46 patients (71.8%), whereas diffuse muscle pain was observed in 42 (65.6%) patients, and 83.3% of them had associated arthritis. Serological, imaging and clinical findings suggested more frequently a seronegative arthritis, whereas few patients developed rheumatoid arthritis and arthritis secondary to a connective tissue disease. A direct correlation between pre-existing autoimmune disease and pain symptoms before cancer diagnosis and risk of developing arthritis under AI therapy was observed. Mean vitamin D levels were significantly lower in patients with diffuse myalgia compared to patients with no diffuse myalgia (11.2 ± 5.8 versus 29.5 ± 10.8 ng/ml; t-test $p < 0.0001$), suggesting a potential involvement of vitamin D deficiency in diffuse muscle pain.

Conclusion: The etiology of AI-associated pain syndrome remain unknown, but clinical, serological, imaging data may be useful to identify the true origin of the musculoskeletal syndrome, as well as a detailed anamnestic history may be useful to identify autoimmune predisposing factors. Myalgias and generalized weakness are likely associated with hypovitaminosis D and might be misdiagnosed as fibromyalgia. The management of these patients should become a multispecialistic task.

REFERENCES

- [1] Beckwée D, Leysen L. Prevalence of aromatase inhibitor-induced arthralgia in breast cancer: a systematic review and meta-analysis. *Support Care Cancer* 2017 May;25(5):1673-1686.
- [2] Borrie AE, Kim RB. Molecular basis of aromatase inhibitor associated arthralgia: known and potential candidate genes and associated biomarkers. *Expert Opin Drug Metab Toxicol*. 2017 Feb;13(2):149-156.

Disclosure of Interests: Stefania Sciacca: None declared, Nadia Melillo: None declared, Francesco Paolo Cantatore Speakers bureau: PFIZER, ROCHE

DOI: 10.1136/annrheumdis-2019-eular.7371

AB1291 ASSOCIATION OF CCL2 GENE POLYMORPHISMS AND THEIR SERUM LEVELS WITH SUSCEPTIBILITY TO KNEE OSTEOARTHRITIS

Amar Chandra Sharma, Rajeswar Nath Srivastava, Sudeepti Srivastava, Saloni Raj. *Indian Institute Of Toxicology Research, Gheru Campus, Developmental Toxicology Division, Lucknow, India*

Background: Osteoarthritis (OA) is one of the most depleting chronic disorders and its pathogenesis is yet to be unfold. It is characterized by pain, swelling, stiffness, decreased ability to move and, sometimes, the formation of bone spurs. Knee Osteoarthritis (KOA) is multifactorial and remains largely understudied at molecular level. The available literature suggested that the development, onset and progression of OA is significantly affected by genetic factors. Chemokines are small, secreted proteins that released from chondrocytes and/or synovial fibroblasts of KOA patients. In this case control study, we investigated the possible association between promoter polymorphisms rs1024611 and the polymorphism rs4586 in exon 2 of the chemokine (C–C motif) ligand 2 (CCL2) gene and knee OA.

Objectives: To investigate the possible association between promoter polymorphisms rs1024611 and the polymorphism rs4586 in exon 2 of the CCL2 gene and KOA.

Methods: DNA was obtained from 300 primary knee OA patients and 300 healthy controls. CCL2 genomic variants (rs1024611 and rs4586 polymorphisms) were detected by polymerase chain reaction restriction fragment length polymorphism (PCR-RFLP). In addition, the effect of serum levels of rs1024611 and rs4586 on cases and controls were examined by enzyme-linked immunosorbent assay (ELISA).

Results: The rs1024611 A/G promoter polymorphism was associated with KOA [genotype frequency, $p = 0.007$; allele frequency, $p = 0.01$]. Significant association was observed between the G carrier of the rs1024611 A/G promoter polymorphism and primary knee OA patients ($p = 0.01$). However, no significant difference was found in the rs4586 polymorphism. Haplotype frequency analysis revealed a significant difference ($\chi^2 = 8.01$, $p = 0.02$). The CCL2 serum level of subjects with the G carrier (285.0 ± 86.5 pg/mL) of the rs1024611 A/G promoter polymorphism was statistically higher than that of subjects with the non-G carrier (160.5 ± 47.8 pg/mL). Further in relation with clinical severity of KOA, we observed significant association of the G carrier of the rs1024611 A/G promoter polymorphism with both Visual analogue scale (VAS) ($p < 0.01$) and Western Ontario and McMaster Universities (WOMAC) score ($p < 0.05$).

Conclusion: The G carrier of the rs1024611 A/G promoter polymorphism was found to be associated with primary knee OA, and could be a susceptibility factor in the development of primary knee OA in the North Indian population.

REFERENCES

- [1] Hulin-Curtis SL, Bidwell JL & Perry MJ. Association between CCL2 haplotypes and knee osteoarthritis. *International journal of Immunogenetics*. 2015
- [2] Park HJ, Yoon SH, Zheng LT, Lee KH, Kim JW, Chung JH, Lee YA, Hong SJ: Association of the 22510A/G chemokine (C–C motif) ligand 2 polymorphism with knee osteoarthritis in a Korean population. *Scand J Rheumatol* 2007;36:299–306

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6642

AB1292 PREVALENCE OF VITAMIN D DEFICIENCY AND ITS ASSOCIATED FACTORS AMONG RHEUMATOID ARTHRITIS PATIENTS MANAGED IN A RHEUMATOLOGY UNIT OF A TERTIARY CARE HOSPITAL IN SRI LANKA

sethugesilva¹, Gunendrika Kasthuriratna², Buddhika Mahesh³, Janalini Sashikaran^{1,2}. ¹College of Specialists in Rheumatology and Rehabilitation – Sri Lanka, Colombo, Sri Lanka; ²College of Specialists in Rheumatology and Rehabilitation – Sri Lanka, Colombo, Sri Lanka; ³University of Melbourne, Melbourne, Australia

Background: Prevalence of unrecognized vitamin D deficiency can be high among Rheumatoid Arthritis (RA) patients. Lack of mobility among of these patients prevent them receiving adequate sun exposure. Low levels of vitamin D could potentially cause a higher disease burden and disease activity.

Objectives: To determine the prevalence of vitamin D deficiency and its associated factors among RA patients managed in a rheumatology unit of National Hospital of Sri Lanka (NHSL).

Methods: A descriptive cross-sectional study was done among patients with RA with a calculated sample size of 137. Being diagnosed according to ACR – EULAR criteria and availability of serum vitamin D level were among the inclusion criteria. Patients with disability due to causes other than RA were excluded. All patients satisfying the eligibility criteria were invited to be recruited. A data extraction sheet was utilized. Data was collected by investigators. The associations were evaluated with Chi square test and Spearman correlation coefficient at a significance level of 5%. Ethics approval was obtained from ethics committee of NHSL.

Results: The response rate was 92%. The median (IQR) age of participants' were 56.5 (49.0 to 64.25). Among participants, majority (n=117, 92.9%) were females. Only 11.1% (n=14) had normal vitamin D levels. The insufficient and deficient categories comprised of 38.1% (n=48) and 50.8% (n=64). The commonest symptoms included; joint pain (n=101, 80.2%), fatigue (n=84, 66.7%) and muscle pain (n=78, 61.9%). Deficiency or insufficiency was lowest in the occupation category of "agricultural and labourer" (37.5%) while 100% in many indoor-occupied categories and among Muslims. "Deficiency or insufficiency" was significantly associated with muscle pain ($p=0.001$) but not with CDAI ($p=0.896$), fatigue ($p=0.549$), joint pain ($p=0.735$).

Conclusion: Vitamin D "deficiency or insufficiency" is common among patients with RA and commoner in the sub-categories with muscle pain and with restricted sun-exposure. More research must be promoted in this regard.

REFERENCES

- [1] Imboden J.B., Hellmann D.B., Stone J.H. 2013. Current treatments and diagnosis rheumatology 3rd edition. New York : McGraw-Hill Medical
- [2] Baker JF, Baker DG, Toedter G, Shults J, Von Feldt JM, Leonard MB (2012) Associations between vitamin D, disease activity, and clinical response to therapy in rheumatoid arthritis. *Clin Exp Rheumatol* 30:658–664 2.
- [3] Braun-Moscovici Y, Toledano K, Markovits D, Rozin A, Nahir AM, Balbir-Gurman A (2011) Vitamin D level: is it related to disease activity in inflammatory joint disease? *Rheumatol Int* 31:493–499
- [4] Craig SM, Yu F, Curtis JR, Alarcon GS, Conn DL, Jonas B, Callahan LF, Smith EA, Moreland LW, Bridges SL Jr, Mikuls TR (2010) Vitamin D status

and its associations with disease activity and severity in African Americans with recent-onset rheumatoid arthritis. *J Rheumatol* 37:275–281(2)

- [5] Haque UJ, Bartlett SJ (2010) Relationships among vitamin D, disease activity, pain and disability in rheumatoid arthritis. *Clin Exp Rheumatol* 28:745–747 5.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3570

AB1293 SMAD3 GENE POLYMORPHISMS AND EXPRESSION IN SERUM AND CARTILAGE INFLUENCE THE RISK OF KNEE OSTEOARTHRITIS

Rajeswar Nath Srivastava, Amar Chandra Sharma, Sudeepti Srivastava, Saloni Raj. *King George's Medical University, orthopaedic surgery, Lucknow, India*

Background: Knee Osteoarthritis (KOA) is the most common degenerative arthritis, a type of arthritis that is caused by breakdown of articular cartilage with eventual loss of the cartilage of the joints. Smad3 is a key intracellular messenger in the transforming growth factor β signaling pathway. Previous study suggested Smad3 gene mutation is a possible predisposing factor for human OA and found gene mutation in OA, providing insight into the function of SMAD3 mediated TGF- β signals in the development of OA and also suggested that Smad3 gene mutation may be a risk factor for genetic susceptibilities to OA. In this case control study, we investigated the possible correlation between the SNPs Smal (C/T; rs6494629), FokI (A/C; rs2289263) in Smad3 gene and susceptibility to knee OA and validated in serum and cartilage.

Objectives: To investigate the possible association between SNPs (rs6494629 and rs2289263) of the Smad3 gene and KOA.

Methods: In this study cases consisted of men and women ≥ 40 years that fulfilled American College of Rheumatology (ACR) clinical and radiographic criteria for knee OA. Venous blood samples were obtained from all cases as well as controls for genetic analysis. Polymerase chain reactions were performed for SNP analysis using specific primer. Total protein was measured in serum by an enzyme linked immunosorbent assay according to the manufacturer's Protocol (ELISA) and in cartilage tissue done by western blot.

Results: A total of 200 cases that confirmed radiographic knee OA and equal number of age and sex matched healthy controls were enrolled. There was no significant difference in demographic characteristics between the cases and controls. A SNP (rs6494629 and rs2289263) mapping to intron 1 of SMAD3 was associated with knee OA ($P < 0.013$ and $P < 0.044$, respectively). Within the SNPs (rs6494629) of Smad3 gene, genotype CC and CT was found to be significantly ($p < 0.013$) associated with knee OA as compared with the CC genotype and SNP rs2289263, genotype CC and CA was found to be significantly ($p < 0.044$) associated with knee OA as compared with the AA genotype. In addition when alleles were compared, C allele of both the studied SNP was observed to be significantly associated with knee OA. Serum levels of Smad3 in KOA patients with rs6494629 TT, CT and CC genotypes were significantly higher than healthy subjects with the same rs6494629C/T genotypes (all $P < 0.0001$, $P < 0.0006$, $P < 0.017$ respectively). Increased serum levels of Smad3 were also observed in KOA patients with rs2289263 AA, CA and CC genotypes compared to controls (all $P < 0.006$, $P < 0.0006$, $P < 0.004$). We performed Immunoblot analysis on cartilage tissue from 15 cases and 10 controls. The smad3 level in cases was significantly higher than controls ($P < 0.006$).

Conclusion: Our data indicate that genetic variation in the SMAD3 gene is involved in the risk of knee OA in North Indian populations, confirming the results from previous studies on the potential importance of this gene in the pathogenesis of OA. Further we also validated these genetic variations at protein level in both blood and tissue and found significant association.

REFERENCES

- [1] Valdes AM, Spector TD, Tamm A, Kisand K, Doherty SA, Dennison EM. Genetic variation in the SMAD3 gene is associated with hip and knee osteoarthritis. *Arthritis and rheumatism*. 2010;10.1002/art.27530
- [2] Su SL, Yang HY, Lee HS, Huang GS, Lee CH, Liu WS. Gene-gene interactions between TGF- β /Smad3 signalling pathway polymorphisms affect susceptibility to knee osteoarthritis. *BMJ Open*. 2015;5:e007931.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6897

AB1294

IMPACT OF INFERTILITY, PREGNANCY LOSS AND CHILDBEARING DECISION ON FAMILY SIZE IN WOMEN WITH SYSTEMIC LUPUS ERYTHEMATOSUS AND RHEUMATOID ARTHRITIS

Dayang Masyrinartie Suahilai¹, Nurulraziqin Mohd Jamid¹, Noraini Mat Husin², Hairul Hadi Ariff¹, Rashidah Bahari³, Nor Shuhaila Shahril¹. ¹Hospital Putrajaya, Medicine, Putrajaya, Malaysia; ²Hospital Raja Permaisuri Bainun, Ipoh, Medicine, Ipoh, Malaysia; ³Hospital Putrajaya, Clinical Research Centre, Putrajaya, Malaysia

Background: Systemic lupus erythematosus (SLE) and Rheumatoid Arthritis (RA) often affects women in their reproductive years. These women are faced with a life-long illness that may have considerable impact not only on their physical health, but also on their reproductive potential. Fertility of these women may also be affected by the disease, treatment and/or organ damage.

Objectives: To determine the role of infertility, pregnancy loss and childbearing decision and patients concerns on family size in women with SLE and RA

Methods: A cross sectional study using a self-administered reproductive history questionnaire completed by woman with SLE/RA attending Rheumatology clinic follow up in Hospital Putrajaya, Hospital Tengku Ampuan Rahimah, and Hospital Raja Permaisuri Bainun, Malaysia from 1 January 2017 to 30 June 2017.

Within each disease cohort, women were identified into 3 groups, those with fewer children than planned (group A), those with same number of planned children (group B) and those with completed family or not interested in having any children (group C). Data on number of children, pregnancies, miscarriages and self reported infertility were recorded. For group A, data on patient concerns and the factors that could impact family building were also obtained.

Results: Total of 110 women with SLE and 91 women with RA were surveyed. The mean age of women with SLE and RA were 37.6 years ($\pm$ SD 7.4) and 45.37 years ($\pm$ SD 11.7) respectively. Majority of women (48.8%) with SLE and RA were in group A with 59% ($n=65$) of women with SLE and 33% ($n=33$) of women with RA had fewer children than originally planned.

The average numbers of pregnancies were similar in both cohorts, but women with SLE had 1 less child and were more likely to report infertility and had higher rate of miscarriages (Table 1). SLE group A had a similar number of pregnancies, but 1 less child compared to SLE group B and C (Table 2). Similarly, among women with RA, group A had 1 less child with similar number pregnancies and miscarriage rate (Table 2). In both groups of women, concerns about inability to care for a child, damage from medications, and genetic transmission of their disease were associated with a lower pregnancy rate.

Table 1. Number of pregnancies, live births, miscarriages and rate of infertility among women with SLE and RA*

	SLE	RA	P
No. of pregnancies	2.28 ± 1.626	2.60 ± 1.949	0.128
No. of children	1.85 ± 1.235	2.19 ± 1.731	0.019
No. of miscarriages	0.46 ± 0.809	0.36 ± 0.659	0.040
No reporting infertility (%)	5 (4.5)	2 (2.2)	0.366

*values are the mean $\pm$ SD unless otherwise indicated

Table 2. Pregnancy outcomes for women with SLE ($n=110$) and RA ($n=91$) *

	Fewer children than desired (group A)		Expected number of children (group B)		completed family (group C)	
	SLE	RA	SLE	RA	SLE	RA
No. of woman, n (%)	65 (59.1)	33 (36.3)	20 (18.2)	31 (34.1)	25 (22.7)	27 (29.7)
No. of pregnancies	2.11 $\pm$ 1.592	2.03 $\pm$ 1.723	2.3 $\pm$ 0.865	2.81 $\pm$ 2.12	2.72 $\pm$ 2.092	3.07 $\pm$ 1.9
No. of children	1.57 $\pm$ 1.212	1.55 $\pm$ 1.416	2.1 $\pm$ 0.553	2.39 $\pm$ 4.856	2.36 $\pm$ 1.497	2.74 $\pm$ 1.745
No. of miscarriages	0.51 $\pm$ 0.831	0.36 $\pm$ 0.603	0.25 $\pm$ 0.716	0.32 $\pm$ 0.653	0.52 $\pm$ 0.823	0.41 $\pm$ 0.747

*values are the mean $\pm$ SD unless otherwise indicated

Conclusion: In this population, more than half of women with RA or SLE had fewer children than desired. Other than patient choice, infertility and miscarriage also play an important role on family size.