Figure c

REFERENCES

- Strand V, Singh JA. Patient Burden of Axial Spondyloarthritis. *J Clin Rheumatol*. 2017 Oct;23(7):383-391.
- Scott DL, Wolfe F, Huizinga TW. Rheumatoid arthritis. *Lancet*. 2010 Sep 25;376(9746):1094-108.
- Anghel LA, Farcaș AM, Oprean RN. Medication adherence and persistence in patients with autoimmune rheumatic diseases: a narrative review. *Patient Prefer Adherence*. 2018 Jul 3;12:1151-116

Disclosure of Interests: Servet Akar Grant/research support from: MSD, Abbvie, Roche, UCB, Novartis, Pfizer, Amgen, Consultant for: MSD, Abbvie, Roche, UCB, Novartis, Pfizer, Amgen, Speakers bureau: Pfizer, Umut Kalyoncu Grant/research support from: MSD, Roche, UCB, Novartis and Pfizer, Consultant for: MSD, Abbvie, Roche, UCB, Novartis, Pfizer and Abdi Ibrahim, Speakers bureau: MSD, Abbvie, Roche, UCB, Novartis, Pfizer and Abdi Ibrahim, Ediz Dalkılı Grant/research support from: MSD and Abbvie, Consultant for: MSD, Abbvie, Roche, UCB, Pfizer and Novartis, Speakers bureau: MSD, Abbvie, Roche, UCB, Pfizer and Novartis, Hakan Emmungil Grant/research support from: MSD, Roche, Pfizer, Abbvie, Consultant for: Novartis, Roche, Speakers bureau: MSD, Roche, Pfizer, Abbvie, Celltrion, Novartis

DOI: 10.1136/annrheumdis-2019-eular.5674

AB1393-HPR COMPARISON OF THE QUALITY OF LIFE, FUNCTIONAL AND EMOTIONAL STATUS OF INPATIENTS AND OUTPATIENTS WITH RHEUMATIC DISEASES

Gamze Arın¹, Aybüke Seven¹, Yasemin Özel Aslıyüce¹, Aykut Özçadırcı¹, Fatma Birgül Kumbaroğlu¹, Şule Apraş Bilgen², Edibe Ünal¹. ¹Hacettepe University Faculty of Physical Therapy and Rehabilitation, Ankara, Turkey; ²Hacettepe University Faculty of Medicine, Department of Rheumatology, Ankara, Turkey

Background: Quality of life, function and emotional status of inpatients with rheumatic diseases may be estimated worse than outpatients. Previous studies had shown that there was a worsening in the emotional state of inpatients (1).

Objectives: The aim of this study is to compare the quality of life, functional and emotional status in inpatients and outpatients with rheumatic diseases.

Methods: The study included 78 patients (inpatient, n = 31; outpatient, n = 47) with rheumatic disease. The Health Assessment Questionnaire (HAQ) (2) and SF-36 (3) were used to evaluate the functionality and quality of life, and the Hospital Anxiety and Depression Scale (HADS) (4) was used to determine their emotional status. The variables were investigated using visual and analytical methods to determine whether or not they are normally distributed. Since physical function, mental health and general health perception values of SF-36 and HADS-Anxiety were normally distributed, the Students T-test was used to compare these parameters between two groups. Physical role limitation, pain, social status, emotional difficulty, energy viability of SF-36, HAQ and HADS-Depression were not normally distributed. Thus, Mann-Whitney U test was used to compare these scores between two groups.

Results: The mean age of the subjects (n = 78) included in the study was 46.09 13.89 years and the mean BMI was 27.59 15.08. There was a significant difference in depression, anxiety and pain and social functioning sub-parameters of SF-36 (p <0.005) but there was no significant difference in other parameters (p > 0.005).

Abstract AB1393HPR Table 1. Comparison of the scores of inpatients and outpatients

	INPATIENTS (XSD)	OUTPATIENTS (XSD)	p
HAQ	0,930,82	0,890,68	0,152
HADS-Anxiety	9,873,8	8,215	0,022
HADS-Depression	9,935,02	7,474,34	0,006
SF-36 Physical Function	47,9428,42	48,3623,81	0,669
SF-36 Physical Role Limitation	27,533,7	27,6336,2	0,869
SF-36 Pain	33,6632,28	54,0721,42	0,011
SF-36 Social Status	39,5828,63	55,0627,09	0,032
SF-36 Mental Health	6121,8	62,7316,08	0,761
SF-36 Emotional Difficulty	31,1038,08	33,3243,83	0,717
SF-36 Energy Vitality	30,8320,63	43,2820,86	0,317
SF-36 General Health Perception	36,1621,68	47,4716,15	0,393

Conclusion: It was thought that during the period of admission to the hospital, inpatients should be supported in terms of pain management, social functioning and anxiety and, depression as well as taking medication. Besides, caregivers in hospitals should encourage inpatients with regard to maintaining physical activity.

REFERENCES

- Lambert, C. M., Hurst, N. P., Forbes, J. F., Lochhead, A., Macleod, M., & Nuki, G. (1998). Is day care equivalent to inpatient care for active rheumatoid arthritis? Randomised controlled clinical and economic evaluation. *Bmj*, 316(7136), 965-969.
- Fries JF, Spitz PW, Young DY: The dimensions of health outcomes: the Health Assessment Questionnaire, Disability and Pain Scales. *J Rheumatol* 9:789-793, 1982
- Brazier, J. E., Harper, R., Jones, N. M., O'cathain, A., Thomas, K. J., Usherwood, T., & Westlake, L. (1992). Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *Bmj*, 305(6846), 160-164.
- Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta psychiatrica scandinavica*, 67(6), 361-370.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7315

AB1394-HPR CAN SUPPORT FROM SIGNIFICANT OTHERS REDUCE SICKNESS ABSENCE IN EARLY RHEUMATOID ARTHRITIS?

Maria Bergström¹, Örfan Dahlström², Ingrid Thyberg^{3,4}, Mathilda Björk^{1,4}. ¹Linköping university, Department of social and welfare studies, Division of occupational therapy, Norrköping, Sweden; ²Linköping university, Department of behavioural sciences and learning, Division of psychology, Linköping, Sweden; ³Linköping university, Department of clinical and experimental medicine, Division of neuro and inflammation science, Linköping, Sweden; ⁴Region stergtland, Department of rheumatology, Heart and medicine center, Linköping, Sweden

Background: Persons with rheumatoid arthritis (RA) are at higher risk of sickness absence, and the probability of returning to work is lower compared to the general population [1]. In order for persons with RA to continue working, support from the social environment is claimed to be of importance [2]. However, this relation needs to be further investigated.

Objectives: To analyze how support from significant others affects the associations between disease related variables (medication, disease activity and activity limitations) at time for RA diagnosis and sickness absence one year after diagnosis.

Methods: Data were collected from 326 (71% women) patients in working age (18-63 years) included in the Swedish early RA cohort TIRA-2 [3] during 2006-2009. At time of inclusion, mean age was 50 years (SD=11), 89% were prescribed disease modifying anti-rheumatic drugs (DMARDs), mean disease activity score 28 joint count (DAS28) was 4.73 (SD=1.34), and mean score for activity limitation reported by Health Assessment Questionnaire (HAQ) was 0.91 (SD=0.60). The number of days with sickness absence during the first year after diagnosis and inclusion was retrieved from the Swedish Social Insurance Agency. Perceived support from significant others, family and friends separately, were self-reported