muscle cells undergo continuous cycles of death and regeneration as a consequence of the pressure of the immune response targeting antigens preferentially expressed within regenerating fibers (1, 2).

Objectives: I will review recent evidence on the relative role of unconventional cellular and molecular pathways activated in the tissues of subjects with IMM and in relevant experimental models of skeletal muscle regeneration and autoimmunity, focusing on the action of antigen-specific and regulatory T cells and immune cells populations of innate cells.

Methods: Review of available published data and of data recently generated on the topic.

Results: Muscle necrosis and the ensuing inflammation was consistently associated in mice with up-regulated expression of bona fide autoantigens targeted in patients with IIM, in particular of histidyl-transfer RNA synthetase (HisRS). Phagocytosis of HisRS-expressing apoptotic myoblasts and appropriate inflammatory costimuli are necessary to elicit the production of anti-HisRS autoantibodies and to jeopardize muscle regeneration. Soluble pattern recognition receptors, and in particular the prototypic long pentraxin PTX3 play a critical role in preventing the accumulation of cell debris in the inflamed skeletal muscle, restricting the immunogenic potential of dying and regenerating myofibers.

Conversely the local expression of type I IFN is necessary for the prolonged autoantibody response and for the maintenance and spreading of the immune-mediated muscle damage, as demonstrated using IFN α / β R-null mice (3-5). A specialized population of regulatory T (Treg) cells, has been recently characterized in the inflamed and regenerating muscle, which influence both the immune response, by promoting the M1-to-M2 switch, and the activation of precursor/stem muscle cells. Treg cells control the efficacy of muscle repair and might be involved in preventing the systemic effects of the autoimmune response generated as a consequence of the disruption of the immunologically protected environment of the skeletal muscle (6, 7).

Conclusion: Identification of the pathways that are physiologically involved in restricting the onset and accelerating the termination of autoimmune responses targeting the skeletal muscle might be valuable for the identification of novel targets for molecular intervention in patients with IIM.

REFERENCES:

- [1] Mammen AL, Casciola-Rosen LA, Hall JC, Christopher-Stine L, Corse AM, Rosen A. Expression of the dermatomyositis autoantigen Mi-2 in regenerating muscle. *Arthritis Rheum.* 2009 Dec;60(12):3784-93.
- [2] Sciorati C, Rigamonti E, Manfredi AA, Rovere-Querini P. Cell death, clearance and immunity in the skeletal muscle. *Cell Death Differ.* 2016 Jun;23(6):927-37.
- [3] Sciorati C, Monno A, Ascherman DP, Seletti E, Manfredi AA, Rovere-Querini P. Required role of apoptotic myogenic precursors and toll-like receptor stimulation for the establishment of autoimmune myositis in experimental murine models. *Arthritis Rheumatol.* 2015 Mar;67(3):809-22.
- [4] Vezzoli M, Sciorati C, Campana L, Monno A, Doglio MG, Rigamonti E, Corra G, Touvier T, Castiglioni A, Capobianco A, Mantovani A, Manfredi AA, Garlanda C, Rovere-Querini P. The clearance of cell remnants and the regeneration of the injured muscle depend on soluble pattern recognition receptor PTX3. *Mol Med.* 2017 Jan;22:809-820.
- [5] Sciorati C, Monno A, Doglio MG, Rigamonti E, Ascherman DP, Manfredi AA, Rovere-Querini P. Exacerbation of Murine Experimental Autoimmune Myositis by Toll-Like Receptor 7/8. *Arthritis Rheumatol.* 2018 Aug;70(8):1276-1287.
- [6] Schiaffino S, Pereira MG, Ciciliot S, Rovere-Querini P. Regulatory T cells and skeletal muscle regeneration. *FEBS J.* 2017 Feb;284(4):517-524. doi: 10.1111/febs.13827
- [7] DiSpirito JR, Zemmour D, Ramanan D, Cho J, Zilionis R, Klein AM, Benoist C, Mathis D. Molecular diversification of regulatory T cells in nonlymphoid tissues. *Sci Immunol.* 2018 Sep 14;3(27).
- [8] Panduro M, Benoist C, Mathis D. Treg cells limit IFN- γ production to control macrophage accrual and phenotype during skeletal muscle regeneration. *Proc Natl Acad Sci U S A.* 2018 Mar 13;115(11):E2585-E2593.

Disclosure of Interests: Patrizia Rovere-Querini Grant/research support from: Adienne Pharma & Biotech

DOI: 10.1136/annrheumdis-2019-eular.8577

SP0104 THE JANUS-FACED GLADIATOR: NEUTROPHILS IN STERILE INFLAMMATION AND AUTOIMMUNITY

Markus Hoffmann. University of Maryland, School of Medicine, United States of America

Background: Neutrophils, and particularly neutrophil extracellular traps (NETs), have been connected with inflammatory and autoimmune diseases, such as RA, SLE, gout and many others. However, the outcome of a deficiency of NET-formation can be observed in chronic granulomatous disease and Papillon-Lefèvre

syndrome and is characterized by non-resolving inflammation and frequent autoimmunity.

Objectives: I will give an overview over pro- and anti-inflammatory effects of neutrophils and NETs, respectively, focusing on our results in gout and SLE.

Methods: We investigated the impact of NET-deficiency and/or neutrophil-depletion on animal models of SLE (pristane-induced lupus) and gout (MSU crystal-induced arthritis). Mechanistic studies were performed with isolated neutrophils from mice and humans.

Results: In experimental lupus and gouty arthritis neutrophil depletion or neutropenia worsens disease. Also mice strains with normal neutrophil numbers but defects in NET formation (NOX2 or PAD4-deficient mice) have exacerbated and unresolving inflammation.

Release of inflammatory mediators by stimulated neutrophils was highest at intermediate cell densities ($20-40 \times 10^6$ cells/cm³). Above such densities, mediator release by normal neutrophils was outweighed by proteolytic degradation via proteases expressed on aggregated NETs. Additionally, binding to aggregated NETs conferred protection of the serine protease neutrophil elastase (NE) against inactivation by α 1-antitrypsin (A1AT).

Conclusion: Although neutrophils are often regarded as archetypical pro-inflammatory cells, evidence is increasing that they can also exert anti-inflammatory and immunoregulatory function. Formation and aggregation of NETs and degradation of inflammatory mediators by serine proteases are important neutrophil tools to resolve gouty arthritis and other forms of localized inflammatory conditions. On the downside, NETs come at the price of collateral tissue damage and may cause occlusion of ductal structures.

REFERENCES:

- [1] Schauer, et al. *Nature Med* 2014, 20(5):511-7.
- [2] Kienhöfer et al. *JCI Insight*, 2(10).
- [3] Hahn et al. *FASEB J* 2019, 33(1):1401-1414.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.8501

SP0105 REGULATION OF AUTOANTIBODY ACTIVITY BY T CELL SUBSETS

Gerhard Krönke. University Hospital Erlangen, Internal Medicine 3, Erlangen, Germany

Our recent data show that the IL-23–T_H17 axis controls the intrinsic inflammatory activity of autoantibodies and thereby triggers the clinical onset of autoimmune arthritis in mice and humans suffering from rheumatoid arthritis. TH17 cells regulate expression of sialyltransferases in newly differentiating antibody-producing cells and determine the glycosylation profile and activity of immunoglobulin G (IgG) that is produced by the consecutively emerging plasma cells.

Disclosure of Interests: Gerhard Krönke Grant/research support from: Lilly, Pfizer, Speakers bureau: Novartis

DOI: 10.1136/annrheumdis-2019-eular.8521

FRIDAY, 14 JUNE 2019

08:15:00 – 09:45:00

Seeing is believing: Nanotechnologies in tissue imaging

SP0106 LOOKING THROUGH MY BACK BONE

Matthias Gunzer. University Hospital, University Duisburg-Essen, Institute for Experimental Immunology and Imaging, Essen, Germany

Background: Because bone marrow is the source of all immune cells, knowing the blood supply of bones is essential to understand diverse mechanisms such as arthritic bone inflammation, anti-infectious immune defense or sterile inflammation.

Objectives: We wanted to clarify, how immune cells are able to function inside of the bone and also leave bones for peripheral functions.

Methods: Using an array of different imaging approaches and innovative animal models as well as insights from clinical studies in humans we have made the bone open to ways of studying, that have not previously possible.

Results: Using these methods we have defined an entirely new concept of how a closed circulatory loop is established in long bones. Thereby we show the mechanistic role of osteoclasts and define, how T cells and neutrophils use the system to fight against viral infection or invade ischemic tissue. This allows an integrated

view how the blood circulation inside and outside of bones are interconnected to mediate functional or dysfunctional immunity.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.8490

SP0107 THE 4-D+ NANOSCOPE: THE RESOLUTION FOR BIOLOGICAL SAMPLES

L. Kling^{1,3}, L. Mill^{3,5}, A. Grüneboom⁵, G. Sarau^{1,3}, M. Herrmann⁴, G. Schett⁴, A. Maier⁵, S.H. Christiansen^{1,2,3}, ¹Helmholtz-Zentrum Berlin für Materialien und Energie, Forschergruppe Christiansen, Berlin, Germany; ²Christiansen Research Group, Helmholtz Zentrum Berlin für Materialien und Energie, Hahn-Meitner-Platz 1, 14109 Berlin, Germany; ³Physics Department, Free University, Berlin, Arnimallee 14, 14195 Berlin, Germany; ⁴Max Planck Institute for the Science of Light, Günther-Scharowsky-Str. 1, 91058 Erlangen, Germany; ⁵Universitätsklinikum der Friedrich-Alexander-Universität Erlangen-Nürnberg, Med. 3, Ulmenweg 18, 91054 Erlangen, Germany; ⁶Friedrich-Alexander-Universität Erlangen-Nürnberg, Pattern Recognition Lab, Martensstr. 3, 91058 Erlangen, Germany

Background: In ageing societies all over the globe, the number of people suffering from bone disease, e.g. osteoporosis (OP) in the first place, has increased dramatically. OP considerably impairs patients' life quality, and results in high societal costs. However, current understanding of OP is still insufficient due to the lack of appropriate high resolution tools that permit a thorough analysis of scale bridging bone architectures from macro to nano with statistical significance to support the development of better treatments by drugs or surgical intervention.

Objectives: To revolutionize our knowledge of bone diseases based on an increase in understanding of the underlying bone anatomy, cutting-edge correlative high-resolution microscopy and spectroscopy together with advanced data analysis including machine learning approaches permit reaching the next, so far unprecedented level of understanding.

Methods: Correlative workflows starting from X-ray microscopy (XRM) volume analysis, with voxel sizes of <1µm, over light-sheet fluorescence microscopy and large scale scanning electron microscopy data acquisition, to dual beam microscope analysis (focused electron- and ion beams) permit the scale bridging investigation of bone architectures and thus merging the "big picture" and the underlying ultrastructure with statistical significance. In combination with additional analytical add-ons to these microscopes, physical properties such as optical, mechanical, compositional, structural etc. deliver a highly detailed correlative dataset of bone (cf. Figure 2).

The present paper demonstrates novel findings related to various partly age related bone diseases using advanced correlative data acquisition (cf. Figure 2).

Results: Figure 1 shows the different bone fine structures, composed of trabecular and vascular networks as well as a three-dimensional arrangement of osteolacunae that host osteocytes, as obtained by volume analysis in a new generation of lab-based x-ray microscope (Zeiss XRM Versa 520). These fine structures can be assessed quantitatively with statistical significance and can be correlated with additional modalities as shown in Figure 2. Elucidating examples demonstrating the power of such an approach will be given [ref 1, ref 2, ref 3, ref 4].

Figure 2 shows the an example correlative workflow from sample collection over sample preparation and the acquisition of various image modalities utilizing correlated data from electron-, ion-beam imaging and analytics, probes and focused laser light to study scale bridging bone architectures. We will demonstrate how these correlative workflows will permit to advance the current understanding of bone architectures and function substantially and will show some first systematic correlative microscopy and spectroscopy studies on mouse tibia and human bone [ref1, ref2, ref3, ref 4].

Abstract SP0107 Figure 1. Bone anatomy with scale – bridging architecture: mass oriented assessment with conventional methods such as DEXA and µCT and function-oriented assessment using correlative microscopy including volume analysis of trans-cortical vessels, osteo-lacunae and canaliculi by XRM and FIB/SEM.

Abstract SP0107 Figure 2. Correlative microscopy/spectroscopy workflow in bone research including (i) the challenges of sample preparation, (ii) the interplay of various image and data modalities, (iii) data evaluation, resolution enhancement and data analysis based on deep learning strategies; finally, new insights in open questions related to bone remodelling and bone structure at various length scales in relation to vasculature will be extracted from statistical correlative microscopy data.

Conclusion: Based on our correlative analytics and the data availability, the advanced data interpretation using machine learning approaches will be discussed. Using this approach, all details of bone micro-/nano-architecture can be used to provide a novel clinical tool-set for future for early detection of bone diseases.

REFERENCES:

- [1] P. Milovanovic, et al., Bone tissue aging affects mineralization of cement lines, *bone* 110, 187 (2018).
- [2] A. Grüneboom, et al., A network of trans-cortical capillaries forms the mainstay for blood circulation in long bones, *Nature Metab.* Published online (02/2019).
- [3] P. Milovanovic, et al., The Formation of Calcified Nanospherites during Micropetrosis Represents a Unique Mineralization Mechanism in Aged Human Bone, *small* 13(3), 1602215 (2017).

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.8516

SP0108 LSFM AND CATCHING UP IN JOINT IMAGING

Anika Grueneboom¹.

Background: Bone tissue differs from other organs by its hardness and high optical density. These physical characteristics are challenging in the context of fluorescence based imaging studies addressing bone anatomy and physiology. The development of novel imaging technologies, such as light-sheet fluorescence microscopy and X-ray microscopy, and tissue-specific preparation procedures like optical clearing allowed us to reveal a so far overseen network of blood vessels in the cortical bone. These trans-cortical vessels (TCVs) directly connect the bone marrow vascularization with the periosteum and make the major contribution to total blood flow in long bones. Furthermore, we could show that immune cells are recruited to the peripheral circulation via TCVs and that TCVs are highly remodeled under inflammatory conditions. In case of chronic arthritis we observed a massive increase in TCV numbers, especially at the metaphysis of long bones. This increase in bone vascularization confirms observations of previous studies describing increasing numbers of microcanals in the bare area of metacarpal heads in the hands of rheumatoid arthritis patients. These alterations in bone anatomy might hint to an alternative mechanism by which immune cells are recruited into inflamed joint tissue. While current concepts describe the recruitment of immune cells from the systemic circulation, immune cells can potentially infiltrate the arthritic joints directly from the underlying bone marrow.

Objectives: Clarifying the anatomical routes of cell recruitment as well as exploring the regulatory mechanism guiding immune cells from the bone marrow to the site of inflammation might bear novel therapeutic strategies to resolve inflammation and tissue injury in rheumatoid arthritis.

Methods: 3D optical imaging approaches like Light-sheet fluorescence microscopy and 2-Photon laser-scanning microscopy enable the visualization of entire bones and joints. In combination with tissue-specific preparation procedures and labeling strategies inflammatory mediated changes in vascularization and cell recruitment can be analyzed.

Results: The joint tissue architecture is massively affected by inflammatory arthritic processes leading to changes in vascularization, permeability and cellular tissue composition. These processes are directly connected to the onset as well as the resolution of inflammatory arthritis.

Conclusion: Newly available imaging techniques as light-sheet fluorescence microscopy facilitate new insights into bone and joint tissue architecture, function and inflammatory processes. These findings might guide the identification of novel