THU0092

DEFINING A LEARNING CURVE FOR CLINICIANS PERFORMING ULTRASOUND-GUIDED DEFINING A LEARNING CURVE FOR CLINICIANS PERFORMING ULTRASOUND-GUIDED NEEDLE SYNOVIAL BIOPSIES: A RETROSPECTIVE ANALYSIS OF THE EXPERIENCE AT THE EXPERIMENTAL MEDICINE AND RHEUMATOLOGY, QUEEN MARY UNIVERSITY OF LONDON

Alessandra Nerviani¹, Frances Humby¹, Gloria Lliso Ribera¹, Felice Rivellese¹, Stephen Kelly², Rebecca Hands¹, Costantino Pitzalis¹. ¹William Harvey Research Institute, Queen Mary University of London, United Kingdom, Centre for Experimental Medicine and Rheumatology, London, United Kingdom; ²Barts Health NHS Trust, Rheumatology, London, United Kingdom

Background: Despite the significantly better outcome reached by patients with Rheumatoid Arthritis following the introduction of the biologics, the absence of predictors of individual response to the available agents constitutes a huge unmet clinical need. The histological analysis of the diseased synovial tissue (ST) may represent a valuable prognostic tool. The ultrasound-guided needle synovial biopsy (US-nSB) is a safe, effective and increasingly used method to retrieve ST from both large and small joints. Therefore, assessing the 'learning curve' for clinicians training in this technique is critically important.

Objectives: To retrospectively evaluate the learning curve for clinicians performing US-nSB at the Centre for Experimental Medicine and Rheumatology, Queen Mary University of London.

Methods: The performance of 5 clinicians has been evaluated from their first to >20 procedures based on the training phase (Table). The standard teaching method included: US/US-guided injections; observation of US-nSB; initial procedures occurring under the strict active supervision of an expert.

Training Phase	1 st - 10 th	11 th - 20 th	21 st - 30 th	31 st - 40 th	41 st - 50 th	>50 th
Total n° of procedures included	50	50	43	35	30	90

Results: The total number of ST fragments retrieved during the procedure did not significantly change during the learning phase (17.3 ± 3.5 from 1st to 10th biopsy vs 17.2 ± 3.4 if expert performer). Independently of the training phase, a significantly higher number of samples were retrieved from knees/wrists in comparison with metacarpo-, metatarso-, and proximal inter-phalangeal joints. The average weight of a ST fragment at the beginning of the learning curve was comparable with the mean weight of ST samples retrieved by expert performers (3.7 ± 2.5 mg from 1st to 10th biopsy vs 2.9 ± 2.4 mg if experts). ST from knees had a significantly higher mean weight, which could be explained by the more frequent use of a 14G (rather than 16G) needle. At the beginning of the learning curve, 48.2%±29.5% of the retrieved ST-samples were considered histologically 'gradable' (visible lining and/or clear sub-lining based on H&E/CD68 IHC staining); the rate of gradable ST progressively increased and remained ~60% when >20 procedures have been performed. The duration of the US-nSB significantly decreased during the learning time (49 ± 18 min from 1st to 10th biopsy vs 30 ± 9 min if expert performer). No serious adverse events (AEs) such as haemarthrosis or joint infection have occurred during the learning time. The overall rate of minor AEs (e.g., minor bleeding) was not the highest at the beginning of the training. During the 1st-10th procedures, 18.2% of the patients reported severe pain compared to the average 8.4%.

Conclusion: In this single-centre retrospective study we showed that, with an appropriate training and the presence of the initial strict supervision, performing US-nSB is safe and effective since the beginning of the learning curve. Further prospective and multi-centre studies are needed to confirm the optimal learning time and training method.

REFERENCES:

- [1] Ultrasound-guided synovial biopsy: a safe, well-tolerated and reliable technique for obtaining high-quality synovial tissue from both large and small joints in early arthritis patients. S Kelly et al. Ann Rheum Dis. 2015 Mar;74(3):611-7.
- [2] Patient-reported outcomes and safety in patients undergoing synovial biopsy: comparison of ultrasound-guided needle biopsy, ultrasound-guided portal and forceps and arthroscopic-guided synovial biopsy techniques in five centres across Europe. SA Just, et al. RMD Open. 2018 Oct 26;4(2):e000799.

Disclosure of Interests: Alessandra Nerviani: None declared, Frances Humby: None declared, Gloria Lliso Ribera: None declared, Felice Rivellese: None declared, Stephen Kelly: None declared, Rebecca Hands: None declared, Costantino Pitzalis Grant/research support from: Celgene
DOI: 10.1136/annrheumdis-2019-eular.7302

THU0093

BASAL METABOLIC RATE AND CHARLSON COMORBIDITY INDEX ARE INDEPENDENT PREDICTORS OF METABOLIC SYNDROME IN PATIENTS WITH RHEUMATOID ARTHRITIS

Dong-Jin Park, Haimuzi Xu, Sung-Eun Choi, Ji-Hyoun Kang, Shin-Seok Lee. Cheonnam National University Hospital, Gwangju, Korea, Rep. of (South Korea)

Background: Metabolic syndrome (MetS) is associated with 2-fold increase in cardiovascular disease (CVD) and 1.5-fold increase in all-cause mortality in the general population. The overall pooled prevalence of MetS in the meta-analysis of rheumatoid arthritis (RA) patients is 30.65%. Along with high prevalence of MetS in RA, several studies showed the contribution of MetS to the increased risk of CVD or cardiovascular mortality in RA patients. Although controversial, the inflammatory activity of RA was significantly associated with MetS in several cross-sectional studies. However, no prospective studies have confirmed an increased risk of MetS development during the course of RA.

Objectives: This 2-year prospective study investigated the predictors of metabolic syndrome (MetS) in patients with rheumatoid arthritis (RA).

Methods: We recruited 319 consecutive RA patients who did not have MetS at baseline. MetS was defined according to the modified NECP/ATP III 2005 for Asian populations. Predictors of MetS were assessed using univariate and multivariate logistic regression analyses.

Results: Of the 247 RA patients who finished 2-year follow-up, 37 (15.0%) developed MetS. At baseline, the patients who developed MetS were older and had higher BMI ($P = 0.002$), higher waist circumference ($P = 0.011$), higher waist-hip ratio ($P = 0.035$), greater skeletal muscle mass ($P = 0.005$), higher body fat mass ($P = 0.011$), higher percent body fat ($P = 0.004$), lower basal metabolic rate ($P = 0.004$) and higher Charlson comorbidity index score ($P < 0.001$). The percent of diabetes ($P = 0.007$) and hypertension ($P < 0.001$) were higher in patients with MetS than without MetS. And they also had elevated glucose ($P = 0.004$), triglyceride levels ($P < 0.001$), and a lower low-density lipoprotein cholesterol (LDL-C) ($P < 0.001$). These patients were also more likely to take less hydroxychloroquine ($P = 0.044$), more oral hypoglycemic agents ($P < 0.001$) and had a lower EQ-5D score ($P = 0.025$). In the multivariate analysis, when variables that belong to the composition of MetS were excluded, basal metabolic rate (OR = 0.205, 95% CI: 0.078–0.541, $P = 0.001$) and Charlson comorbidity index score (OR = 2.191, 95% CI: 1.280–3.751, $P = 0.004$) remained significant predictors of MetS.

Conclusion: The development of MetS in RA patients was associated with basal metabolic rate and Charlson comorbidity index at baseline. Physicians should pay more attention to RA patients who have these risk factors to avoid CVD and cardiovascular mortality.

Disclosure of Interests: None declared
DOI: 10.1136/annrheumdis-2019-eular.3419

THU0094

SERONEGATIVE RHEUMATOID ARTHRITIS: MILD NO MORE

Sofia Pazmino¹, Annelies Boonen², Veerle Stouten¹, Diederik De Cock¹, Delphine Bertrand^{1,3}, Kristien Van der Elst³, Johan Joly³, Rene Westhovens^{1,3}, Patrick Verschueren^{1,3}, CareRA study group. ¹Katholieke Universiteit Leuven, Skeletal Biology and Engineering Research Centre, Leuven, Belgium; ²Maastricht University Medical Centre, Department of Internal Medicine, Division of Rheumatology, Maastricht, Netherlands; ³University Hospitals Leuven, Rheumatology, Leuven, Belgium

Background: Seronegative rheumatoid arthritis (RA) patients have been historically considered to have a milder disease. This confers a potential risk for under treatment. The quality of their response to early intensive therapies has not been thoroughly studied.

Objectives: To compare the disease course in seronegative and seropositive patients from the Care in early RA (CareRA) trial treated with COBRA Slim.