THU0125

CARDIOVASCULAR DISEASE RISK FACTORS MAY NEGATIVELY IMPACT RHEUMATOID ARTHRITIS DISEASE OUTCOMES: FINDINGS FROM THE ONTARIO BEST PRACTICES RESEARCH INITIATIVE

Kangping Cui^{1,2}, Mohammad Movahedi², Claire Bombardier^{2,3}, Bindee Kuriya^{2,4}.¹Western University, Internal Medicine, London, Canada; ²University Health Network, Toronto General Hospital Research Institute, Toronto, Canada;³University of Toronto, Department of Medicine and Institute for Health Policy, Management and Evaluation, Toronto, Canada; ⁴Mount Sinai Hospital, Rheumatology, Toronto, Canada

Background: Rheumatoid arthritis increases the risk of cardiovascular disease (CVD). Less is known about the direct influence of CVD and CVD risk factors (RF) on RA outcomes, but higher comorbidity burden has been suggested to adversely affect RA treatment response¹⁻².

Objectives: We tested our hypothesis that CVD risk factors (RFs) alone, in the absence of CVD, are associated with higher disease activity and disability in RA.

Methods: The Ontario Best Practices Research Initiative (OBRI) is a clinical registry of RA patients followed in routine care. RA subjects with complete data to calculate disease activity according to the Disease Activity Score-28 (DAS28), Clinical Disease Activity Index (CDAI), 28 swollen joint count (SJC28) and functional status (Health Assessment Questionnaire Disability Index [HAQ-DI]) at cohort entry were selected. Patients were divided into mutually exclusive groups by baseline CVD status as: (1) no CVD/no CVD RFs; (2) CVD including coronary artery disease, myocardial infarction, cerebral vascular accidents, and peripheral arterial disease; (3) no CVD but CVD RFs including hypertension (HTN), dyslipidemia (DLD), diabetes (DM), or smoking. We performed separate linear regression analyses for each outcome, adjusted for baseline clinical and demographic variables, to determine the independent effect of CVD status on disease outcomes at baseline and one year follow-up.

Results: Of 2033 patients examined, 49.5% had no CVD, 5.4% had CVD and 45.1% had CVD RFs alone. The most common RF was HTN (33%) followed by DLD (19.7%), current smoking (17%), and DM (8.1%). At cohort entry, having a CVD RF was associated with significantly higher DAS28 (β 0.13, 95%CI 0.002-0.26, p 0.04) and HAQ-DI (β 0.16, 95%CI 0.10-0.23, p <0.0001). At one year, CVD RF was associated with worse DAS28 (β 0.17, 95%CI 0.05-0.30, p 0.01) and CDAI (β 0.96, 95%CI 0.05-1.87, p 0.04) but not HAQ-DI (β 0.03, 95%CI -0.002-0.08, p 0.17). Having higher number of CVD RF was associated with worse disease outcomes. No association between CVD status and swollen joint count was observed.

Conclusion: Even in the absence of CVD, traditional CVD RF are associated with greater RA disease severity and disability both at baseline and one year. Self-perceived impact of comorbidity (patient global assessment of health) may be driving this relationship. Moreover, patients with CVD RF maybe more treatment-resistant, suggesting that co-management of CVD RF in RA patients may be beneficial on both fronts.

REFERENCES:

- Avina-Zubieta, J. A., Thomas, J., Sadatsafavi, M., Lehman, A. J. & Lacaille, D. Risk of incident cardiovascular events in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Ann. Rheum. Dis.* 71, 1524-1529 (2012).
- Crepaldi, G. et al. Cardiovascular Comorbidities Relate More than Others with Disease Activity in Rheumatoid Arthritis. *PLoS One* 11, (2016).

Abstract THU0125 Table 1. Multivariable linear regression of disease activity outcomes and functional status at one-year according to CVD status.

	β - coefficient (95% CI); p -value			
	DAS28-ESR	CDAI	SJC-28	HAQ-DI
CVD status	N=1965	N=1965	N=1965	N=1965
No CVD/No CVD RFs	Ref	Ref	Ref	Ref
CVD	0.12 (-0.14,0.38), 0.37	0.50 (-1.45,2.46), 0.61	0.44 (-0.21,1.10), 0.19	0.09 (-0.02,0.19), 0.10
CVD RFs/No CVD	0.17 (0.05,0.30), 0.01	0.96 (0.05,1.87), 0.04	0.29 (-0.01,0.60), 0.06	0.03 (-0.02,0.08), 0.17

Acknowledgement: OBRI investigators

Disclosure of Interests: Kangping Cui: None declared, Mohammad Movahedi Employee of: ONTARIO BEST PRACTICE RESEARCH INITIATIVE (OBRI), Claire Bombardier Grant/research support from: Abbvie, Amgen, Bristol-Myers Squibb, Celgene, Eli Lilly, Hospira, Janssen, Merck, Novartis, Pfizer Inc, Roche, Sanofi, UCB, Consultant for: AbbVie, Hospira, Janssen, Merck, Novartis, Pfizer Inc, Sanofi, Speakers bureau: Roche, Bindee

Kuriya Consultant for: Advisory Board Member: Abbvie, Sanofi Genzyme, Eli Lilly, Pfizer Canada, Roche

DOI: 10.1136/annrheumdis-2019-eular.2484

THU0126

FACTORS AFFECTING THE DISCREPANCY BETWEEN PATIENT AND PHYSICIAN GLOBAL ASSESSMENT OF DISEASE ACTIVITY IN PATIENTS WITH EARLY RHEUMATOID ARTHRITIS OVER TIME

Pooneh Akhavan¹, Mohammad Movahedi², Claire Bombardier^{1,2}. ¹University of Toronto, Mount Sinai Hospital, Rheumatology, Toronto, Canada; ²Toronto General Research Institute, University Health Network, Ontario Best Practices Initiatives, Toronto, Canada

Background: Rheumatoid Arthritis (RA) is a chronic disease that requires regular follow-ups over time. Physicians make treatment adjustments based on their evaluation of disease status and patients' response to treatment while patients' perception of disease activity may change over time.

Objectives: we aimed to assess the discrepancy between patient global assessment of disease activity and physician assessment of disease activity at baseline and after one year in patients with early RA. We also evaluated factors affecting this discrepancy at these two time points.

Methods: Patients enrolled in the Ontario Best Practices Research Initiative (OBRI), a prospective observational cohort of patients with RA who had disease duration of <12 months (early RA) and had visits both at baseline and after one year were included. The discrepancy between patient global assessment of disease activity (PTGA) and physician global assessment of disease activity (MDGA) was calculated by simple subtraction (PTGA-MDGA). The value of ≥ 30 was considered discordant which could be either positive or negative in favor of PTGA or MDGA. Linear regression analysis was used to assess factors affecting PTGA, MDGA, and PTGA-MDGA discrepancy at baseline and one year.

Results: A total of 460 patients with early RA were analyzed. Majority were female (72.4%) with mean ($\pm$ SD) age of 57 years (± 0.6). the discordance rate was 109 (24%) and 99 (21%) at baseline and one year, respectively. In the majority of the discordant cases, PTGA was higher (98 (90%) at baseline and 85 (86%) at one year). Discordant patients had significantly higher fatigue score, health assessment questionnaire disability index (HAQ-DI), pain score, PTGA, MDGA, tender joint, composite measures of disease activity and higher number of comorbidities at baseline.

Multivariable regression analysis showed that the higher PTGA is significantly associated with higher SJC, TJC and fatigue both at baseline and one year and with higher pain score only at one year. Similar associations were found for MDGA except fatigue which was not significant. Multivariable regression analysis showed that the higher discrepancy between PTGA and MDGA is associated significantly with lower SJC (β =2.75), higher TJC (β =1.17), and higher fatigue (β =0.85) at baseline. At one-year follow-up, higher pain was the strongest factor (β =4.99) for affecting the higher PTGA-MDGA discrepancy compared with SJC (β =1.65) and fatigue (β =1.24).

Conclusion: Significant discrepancy between PTGA and MDGA exists in about a quarter of patients with early RA at baseline and slightly less after one year. In the majority of these patients PTGA is higher over time. The number of active joints affects both assessments and this effect persists at one year. Clinicians should consider the above associations when making

Acknowledgement: OBRI-RA Investigators: Drs. Ahluwalia, V., Ahmad, Z., Akhavan, P., Albert, L., Alderdice, C., Aubrey, M., Aydin, S., Bajaj, S., Bensen, B., Bhavsar, S., Bobba, R., Bombardier, C., Bookman, A., Cabral, A., Carrette, S., Carmona, R., Chow, A., Ciaschini, P., Cividino, A., Cohen, D., Dixit, S., Haaland, D., Hanna, B., Haroon, N., Hochman, J., Jaroszynska, A., Johnson, S., Joshi, R., Kagal, A., Karasik, A., Karsh, J., Keystone, E., Khalidi, N., Kuriya, B., Larche, M., Lau, A., LeRiche, N., Leung, Fe., Leung, Fr., Mahendira, D., Matsos, M., McDonald-Blumer, Midzic, I., Milman, N., H., Mittoo, S., Mody, A., Montgomery, A., Mulgund, M., Ng, E., Papneja, T., Pavlova, P., Perlin, L., Pope, J., Purvis, J., Rohekar, G., Rohekar, Ruban, T., S., Samadi, N., Shaikh, S., Shickh, A., Shupak, R., Smith, D., Soucy, E., Stein, J., Thompson, A., Thorne, C., Wilkinson, S.g treatment decisions.

Disclosure of Interests: Pooneh Akhavan: None declared, Mohammad Movahedi: None declared, Claire Bombardier Grant/research support from: Abbvie, Amgen, Bristol-Myers Squibb, Celgene, Eli Lilly, Hospira, Janssen, Merck, Novartis, Pfizer Inc, Roche, Sanofi, UCB, Consultant for: AbbVie, Hospira, Janssen, Merck, Novartis, Pfizer Inc, Sanofi, Speakers bureau: Roche

DOI: 10.1136/annrheumdis-2019-eular.4888

THU0127

IMMUNOGENICITY AND SAFETY OF 23-VALENT PNEUMOCOCCAL VACCINE IN PATIENTS WITH RHEUMATOID ARTHRITIS: RESULTS FROM 5-YEAR FOLLOW UP

Daria Bukhanova, Boris Belov, Galina Tarasova, Marina Sergeeva, Yury Muravyev, Galina Lukina, Natalia Demidova, Maria Cherkasova. V.A. Nasonova Research Institute of Rheumatology, Moscow, Russian Federation

Background: In rheumatology, comorbid infections have significant impact on patients' morbidity, mortality and quality of life. Prevention of infections is an integral part of supervision of these patients.

Objectives: The aim of the study is to investigate immunogenicity and safety of 23-valent polysaccharide pneumococcal vaccine (PPV-23) in patients with rheumatoid arthritis (RA) in a five-years period.

Methods: The study included 79 RA patients with ≥ 2 recent episodes of respiratory tract infections (bronchitis, pneumonia). 52 RA pts were treated with methotrexate (MT), 14 – with leflunomide (Lef), 13– with TNF- α inhibitors + MT. One dose (0,5 ml) of PPV-23 was administered subcutaneously without discontinuing MT/Lef or 28-30 days prior to initiation of TNF- α inhibitors. Control visits were scheduled as follows: at baseline (Visit 1), and in 1, 3, and every year after immunization. 39 out of 110 pts were followed for 24 months, 23 pts – for 36 months, 23 pts – for 48 months, and 18 – for 60 months. Standard clinical examination and lab tests were performed at each visit.

Levels of serum antibodies (AB) to Pneumococcal capsular polysaccharide were measured with VaccZymeTM PCP IgG 2 panels (The Binding Site Group Ltd, Birmingham, UK). Coefficient of post-immunization response (CPR) was determined for each patient as the ratio of AB levels at Visits II, III, IV, V, VI and VII to AB level at Visit I.

Results: 53 (67%) patients did not have any reactions to the vaccine, 26 (33%) patients indicated pain, swelling and hyperemia of the skin (diameter up to 2 cm) at the injection site of the vaccine, low-grade fever. These typical reactions following vaccination are completely regressed within days without any additional treatment. They were not associated with RA therapy and did not require changes in ist schemes.

Pronounced positive immune reaction to PPV-23 was documented in all RA patients on different therapeutic regimens as significant AB incremental growth. Significantly high level of post-immunization response still persisted after 4 and 5 years of follow-up. The proportion of responders to PPV-23 after 5 years reached 73% among RA patients. (Table).

No case of clinically or radiographically confirmed pneumonia was documented during the follow up period, although one case of interstitial viral pneumonia was registered in 6 month after immunization.

Not a single vaccination-related RA exacerbation episode or new autoimmune phenomenon was documented during the follow-up.

Table. AB dynamics in RA pts during 5-year FUP, M $\pm$ σ .

	Visit II (1 month)	Visit IV (12 month)	Visit V (24 month)	Visit VI (36 month)	Visit VII (48 month)	Visit VIII (60 month)
RA	82,20	250,62*	298,70*	107,8	140,5*	194,8*
patients	[46,00; 133,50]	[187,70; 316,90]	[175,99;420,81]	[98,2;159,4]	[107,9;208,3]	[120,1;361,4]

*p<0,05

Conclusion: Thus, all given prove the sufficient immunogenicity and safety of 23-valent pneumococcal vaccine in RA patients after 5 years of follow-up, getting different therapeutic regimens.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.3173

THU0128

GLUCOCORTICOID TAPERING IN MONTHLY 1-MG DECREMENTS DOES NOT RESULT IN CLINICALLY MANIFEST ADRENAL INSUFFICIENCY IN PATIENTS WITH RHEUMATOID ARTHRITIS: LEARNINGS FROM THE PHASE 3/4 SEMIRA STUDY

Frank Buttgerit¹, J. Michael Nebesky², Gerd Rüdiger Burmester¹, Marc Hochberg³, Corrado Bernasconi², Markus R. John², Marc Y. Donath⁴.
¹Charité-Universitätsmedizin Berlin, Free University and Humboldt University of Berlin, Berlin, Germany; ²F. Hoffmann-La Roche, Basel, Switzerland; ³University of Maryland School of Medicine, Baltimore, United States of America; ⁴University Hospital Basel, Basel, Switzerland

Background: Systemic glucocorticoids (GCs) are used to treat serious inflammatory diseases but are associated with adverse events. Guidelines recommend tapering GCs to the lowest possible dose and discontinuing them as soon as possible. Physicians have concerns that reductions, even from low doses, may increase disease symptoms or cause adrenal insufficiency (AI), especially in patients receiving GCs long-term. The expectation of AI risk may be inflated by false positives in cortisol testing of patients without relevant symptoms and by a lack of robustly designed GC taper trials. The international, multicenter SEMIRA trial evaluated a taper scheme in rheumatoid arthritis (RA) patients receiving tocilizumab (TCZ) $\pm$ conventional synthetic disease-modifying antirheumatic drugs (csDMARDs).¹

Objectives: To determine whether GC tapering in the SEMIRA trial resulted in any cases of AI.

Methods: Eligible patients had low disease activity or remission for ≥ 4 weeks and were receiving a stable prednisone regimen (5 mg/day) + TCZ $\pm$ csDMARDs for ≥ 4 weeks. Patients had to have received ≥ 6 months' total TCZ + GCs (prednisone equivalent 5-15 mg/day). Patients were randomly assigned (1:1) to double-blind continued prednisone 5 mg/day (n = 128) or prednisone taper (n = 131). TCZ $\pm$ csDMARDs remained stable during the 24-week study. The prednisone taper consisted of 1-mg decrements every 4 weeks starting at randomization and ending at GC discontinuation (week 16). RA flares were treated with prednisone 5 mg/day for 2 weeks. The primary assessment was maintenance of disease control with GC discontinuation. The protocol provided evidence-based guidance for diagnosis and management of AI. Confirmatory testing was recommended for patients with suspected AI, but routine precautionary ACTH stimulation testing was not mandated.

Results: In the taper arm, 65% of patients achieved "treatment success" by meeting all key secondary end point components (maintained low disease activity, experienced no flares, and had no confirmed AI necessitating replacement therapy) at week 24 versus 77% of continued patients (risk ratio for treatment success, 0.833 [95% CI, 0.714-0.972]; p = 0.021). None of the taper patients required ACTH stimulation testing, and no cases of AI were reported.

Conclusion: SEMIRA demonstrated the usefulness of a new standardized GC dose taper scheme. Two-thirds of patients receiving TCZ underwent successful tapering and could stop GCs entirely, which is higher than the spontaneous 35% discontinuation rate observed in real-world RA patients² and underscores the potential to further reduce glucocorticoid burden. Clinical AI was not observed; thus, routine laboratory testing may be unnecessary in real-world applications of this taper scheme.

REFERENCES:

- [1] Burmester GR, et al. *Arthritis Rheumatol.* 2018;70(suppl 10):L18. 2. Yagiz B, et al. *Arthritis Rheumatol.* 2018;70(suppl 10):627.

Disclosure of Interests: Frank Buttgerit: None declared, J. Michael Nebesky Shareholder of: F. Hoffmann-La Roche, Employee of: F. Hoffmann-La Roche, Gerd Rüdiger Burmester Consultant for: Roche, Sanofi-Genzyme, Speakers bureau: Roche, Sanofi-Genzyme, Marc Hochberg Shareholder of: BriOri Biotech, Theralogix LLC., Consultant for: Bristol Myers Squibb, Eli Lilly, EMD Serono, Novartis Pharma AG, Pfizer Inc., Samumed LLC, Symic Bio Inc., Theralogix LLC, TissueGene Inc., TLC Biopharmaceuticals, Inc., Zynerba, Galapagos, IQVIA, Hoffman LaRoche., Corrado Bernasconi Consultant for: Roche, Markus R. John Shareholder of: Roche, Employee of: Roche, Marc Y. Donath Consultant for: Roche

DOI: 10.1136/annrheumdis-2019-eular.2113