Disclosure of Interests: Sarah Wood Grant/research support from: Pfizer stipend received for MPhil in epidemiology in 2013, Employee of: Employed by GSK in 2012 as a pre-registration pharmacist, Kimme Hyrich Grant/research support from: Grants to institution: BMS, Pfizer, UCB, Suzanne Verstaappen: None declared, Douglas Steinke: None declared
DOI: 10.1136/annrheumdis-2019-eular.4721

THU0201 ANALYSIS OF GUT MICROBIOTA DIVERSITY IN PATIENTS WITH RHEUMATOID ARTHRITIS AND BENEFICIAL EFFECTS OF PROBIOTICS ON INTESTINAL FLORA

Ming Yan¹, Sheng-Xiao Zhang¹, Chong Gao², Xiao-Bin Zheng¹, Lili Shang¹, Xu-Fang Yin¹, Zhang Mingxing¹, Jia Wang¹, Ting Cheng¹, Nalin Lai¹, Li Xiao-Feng¹.
¹The Second Hospital of Shanxi Medical University, Department of Rheumatology, Taiyuan, China; ²Brigham and Women's Hospital, Harvard Medical School, Department of Pathology, Boston, United States of America

Background: Rheumatoid arthritis (RA) is a progressive, irreversible disease characterized by autoimmune imbalance. Recent studies have found that the number and type of intestinal flora in RA patients were changed (called disorders), which in turn leads to immune imbalance in RA patients. The breath test can indirectly assess the growth of bacteria in the intestine. This study aimed to investigate the intestinal flora imbalance in RA patients and the effect of probiotics on the intestinal flora of RA patients using methane and hydrogen breath test.

Objectives: To study the beneficial effects of probiotics on the intestinal flora of RA patients by detecting the concentrations of methane and hydrogen and the microbial populations in patients with rheumatoid arthritis (RA) before and after probiotic treatment.

Methods: The lactulose methane and hydrogen breath technique were used to detect and compare the concentration of methane and hydrogen in 36 patients with RA before and after probiotic treatment (mainly *Bifidobacterium*, *Clostridium butyricum*, *Lactobacillus*) for 3 weeks. At the same time, the 16S rRNA V3 region in stool samples of 17 patients with RA was qualitatively analyzed by Roche/45 high-throughput sequencing platform and compared with 9 healthy adult stool samples.

Results: Compared with pre-intervention, after 3 weeks of probiotic treatment, the exhaled methane concentration was significantly increased in RA patients at 30 minutes, 60 minutes, and 90 minutes ($P < 0.05$). The sum of methane and hydrogen concentrations increased at 30 minutes ($P = 0.041$). There was no significant difference in hydrogen concentration before and after intervention ($P > 0.05$) (Table 1). Compared with healthy adults, the fecal diversity index (Shannon) of RA patients increased ($P < 0.05$) and the species uniformity index (flatness) also increases.

Figure 1: Compared with healthy subjects (n=9), RA patients (n=17) increased the fecal diversity index, $P < 0.05$.

Abstract THU0201 – Figure 1

Conclusion: Patients with RA had a significant difference in the structure of intestinal flora compared with that of healthy controls, which may represent perturbations of microbial communities and contribute to the pathogenesis and condition of RA. Probiotics can restore the composition of the gut microbiome and introduce beneficial functions to gut microbial communities through diet, probiotics or fecal transplantation, providing a promising prospect for clinical treatment of RA.

Figure 2: Compared with healthy people (n=9), RA patients (n=17) The average index of feces increased, $P < 0.003$.

Abstract THU0201 – Figure 2

REFERENCES:

- [1] Schett I B M a G, *The pathogenesis of rheumatoid arthritis*. The New England Journal of Medicine, 2011. 365(23): p. Number of 2205–2219.
- [2] Edwards C J, *Commensal gut bacteria and the etiopathogenesis of rheumatoid arthritis*. J Rheumatol, 2008. 35(8): p. Number of 1477–1479.
- [3] Horta-Baas G, Romero-Figueroa MDS, Montiel-Jarquín AJ, Pizano-Zárate ML, García-Mena J, Ramírez-Durán N. Intestinal Dysbiosis and Rheumatoid Arthritis: A Link between Gut Microbiota and the Pathogenesis of Rheumatoid Arthritis. J Immunol Res. 2017;2017:4835189

Disclosure of Interests: None declared
DOI: 10.1136/annrheumdis-2019-eular.6436

SLE, Sjögren's and APS – etiology, pathogenesis and animal models

THU0202 INTERFERON STIMULATED GENE 15 PROTECTS REGULATORY T CELL OF SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS FROM INTERFERON ALPHA-MEDIATED DEPLETION

Francesca Romana Spinelli, Ilenia Pacella, Silvia Piconese, Fulvia Ceccarelli, Francesca Miranda, Cristiano Alessandri, Vincenzo Barnaba, Guido Valesini, Fabrizio Conti. Sapienza University of Rome, Rome, Italy

Background: Data on T regulatory (T reg) cells number and functions in patients with Systemic Lupus Erythematosus (SLE) are conflicting (1). Type I interferon (IFN) decreases Treg proliferation and induces Treg apoptosis (2). IFN stimulated gene 15 (ISG15) is an IFN-induced protein with a negative effect on IFN pathway.

Objectives: Aim of the study was to evaluate the effect of IFN alpha and ISG15 on Treg number and on STAT1 phosphorylation in patients with SLE. **Methods:** We recruited patients with SLE classified according to 1997 ACR criteria. We collected demographic and clinical data including disease duration, disease activity scored according to *Systemic Lupus Erythematosus Disease Activity Index* (SLEDAI) 2K. Peripheral blood mononuclear cells (PBMC) were isolated from whole blood samples. Intracellular phosphorylated STAT1 (pSTAT) levels were evaluated in Treg (CD4+FOXP3+CD127 low cells) and conventional T cell (Tconv, CD4+FOXP3- cells) by multiparametric flow cytometry, *ex vivo* or after stimulation with recombinant IFN alpha. PBMC were cultured for 48 hours with anti-CD3 or anti-CD3 and IFN alpha; the frequency of Treg was analysed by multiparametric flow cytometry. ISG15 mRNA expression was evaluated by RT-PCR on PBMC from SLE patients. Data were expressed as mean $\pm$ standard deviation or median (interquartile range) according to the distribution; Mann-Whitney and Spearman tests were applied. P value < 0.05 were considered statistically significant.

Results: We enrolled 21 SLE patients [F:M 20:1; mean age 45.2 ± 12.9 years; mean disease duration 157.5 ± 103 months; median SLEDAI 2(4)]. Overall, median baseline ISG15 mRNA expression was 0.06 (IQR 0.16); patients were divided into ISG15-high (n=9) and ISG15-low (n=12), according to the mean mRNA expression. *Ex vivo*, after short-term treatment with IFN alpha (15 and 25 minutes), we observed a significant increase in STAT1 phosphorylation compared to baseline both in Treg and Tconv ($p < 0.00001$), in ISG15-high and ISG15-low patients (Figure 1 A and B); however, high ISG15 levels protect both Treg and Tconv from STAT1 phosphorylation (Figure 1 C). After 48 of culture with IFN alpha, we observed a decrease in Treg and Tconv number, significantly greater in ISG15-low cells (Figure 1D). Treg frequency and ISG15 expression in Treg were higher in active vs inactive SLE patients [SLEDAI > 4 vs SLEDAI < 4] ($p < 0.05$).

Conclusion: ISG15 protects T conventional cells from STAT1 phosphorylation and induces resistance to T regulatory cells depletion. The results of the study suggest that ISG15, a protein induced by IFN in the acute phase of the disease, exerts a negative feedback, allowing the recovery of T reg.

Figure 1. Effect of ISG15 on STAT phosphorylation and number of Treg and Tconv **Abstract THU0202 – Figure 1****REFERENCES:**

- [1] Zhang SX, et al. J Immunol res. 2018
- [2] Piconese S, et al. Cytokine Growth Factor rev. 2014

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7625

THU0203

PROMOTER DNA METHYLATION AND HSA-MIR-424-5P REGULATE ATF6 ALPHA EXPRESSION IN SALIVARY GLANDS OF PATIENTS WITH SJÖGREN'S SYNDROME

Patricia Carvajal¹, Sergio Aguilera², María-José Barrera³, Carolina Lagos¹, Isabel Castro⁴, Sergio González⁵, Daniela Jara¹, Claudio Molina³, María-Julieta González¹. ¹ICBM, Facultad de Medicina, Universidad de Chile, Santiago, Chile; ²Clinica INDISA, Santiago, Chile; ³Facultad de Odontología, Universidad San Sebastián, Santiago, Chile; ⁴Departamento Tecnología Médica, Universidad de Chile, Santiago, Chile; ⁵Escuela de Odontología, Universidad Mayor, Santiago, Chile

Background: Salivary glands (SG) from Sjögren's syndrome (SS)-patients show chronic inflammation and endoplasmic reticulum (ER) stress. To counteract ER stress, eukaryotic cells activate adaptive mechanisms referred to as the unfolded protein response via sensors present in the ER membrane, such as ATF6α^{1,2}. In response to ER stress, ATF6α is translocated to the Golgi apparatus where is processed releasing ATF6f, which translocates into the nucleus as a transcription factor to activate ER-associated degradation and protein folding-related gene transcription³. The overexpression of ATF6α observed in Labial SG (LSG) from SS-patients¹ could be modulated by epigenetic mechanisms.

Objectives: To evaluate the involvement of epigenetic mechanisms such as promoter-specific DNA methylation and microRNA (miRNA) regulation on ATF6α expression in LSG from SS-patients and controls. The effect of IFN-γ on both mechanisms was also evaluated in a 3D-acini model.

Methods: SG biopsies from 12 SS-patients and 10 controls were analyzed. For in vitro assays 3D-acini incubated with 1 or 10 ng/mL IFN-γ for 24 h were employed. Specific DNA methylation of ATF6α gene promoter was evaluated by methylation sensitive-high resolution melting PCR. The profile of miRNAs differentially expressed was determined by miRNA sequencing (seq), of which hsa-miR-424-5p was selected because the *in silico* analysis showed that ATF6α mRNA is a target of this miRNA. The levels of hsa-miR-424-5p were determined by Taqman miRNA assays. Functional assays using miRNA mimic and miRNA inhibitors were also performed.

Results: LSG from SS-patients showed a significant decrease of ATF6α promoter DNA methylation percentage (p=0.04), which was inversely correlated with high ATF6α mRNA levels (p= 0.02). The hsa-miR-424-5p was found downregulated by miRNA-seq in LSG from SS-patients. Taqman miRNA assays validated the significant decrease of hsa-miR-424-5p levels found in LSG from SS-patients (p=0.02), which were inversely correlated with ATF6α mRNA levels (p=0.02). Functional assays suggest that hsa-miR-424-5p regulates the mRNA levels of ATF6α. Finally, stimulation of 3D-acini with IFN-γ induced a significant increase of ATF6α mRNA levels together with a decrease in ATF6α promoter DNA methylation percentage and hsa-miR-424-5p levels.

Conclusion: The results suggest that the ATF6α overexpression is regulated by two complementary epigenetic mechanisms: the DNA hypermethylation of its promoter and the silencing of its mRNA by hsa-miR-424-5p in LSG from SS-patients. IFN-γ affects the expression of ATF6α by modulation of epigenetic mechanisms, regulating the ability of epithelial cells to handle the chronic ER stress. Interestingly, these changes carry out under a global DNA hypomethylation condition⁴.

REFERENCES:

- [1] J Autoimmun 2016;75:68-81
- [2] Rheumatology 2018;57:1021-1032
- [3] Autoimmun Rev 2018;17:796-808
- [4] Clin Immunol 2018;196:85-96

Acknowledgement: Fondecyt-1160015, Fondecyt-Postdoctorado-3170023, Fondecyt-Iniciación-11170049.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4712

THU0204

ASSOCIATION OF LILRA3 GENE WITH LYMPHOMAGENESIS RISK IN YOUNG SS PATIENTS

Evangelia Argyriou^{1,2}, Petros Roussos¹, Adrianos Nezos¹, Aliki Venetsanopoulou³, Kyriaki A. Boki², Athanasios Tzioufas³, Haralampos M. Moutsopoulos⁴, Clio Mavragani^{1,3,5}. ¹Medical School, National and Kapodistrian University of Athens, Department of Physiology, Athens, Greece; ²Rheumatology Unit, Sismanogleio General Hospital, Athens, Greece; ³Medical School, National and Kapodistrian University of Athens, Department of Pathophysiology, Athens, Greece; ⁴Academy of Athens, Athens, Greece; ⁵Joint Academic Rheumatology Program, National and Kapodistrian University of Athens, School of Medicine, Athens, Greece

Background: Primary Sjögren's Syndrome (SS) is a systemic autoimmune disease, characterized by increased risk of non-Hodgkin's lymphoma development. A functional polymorphism of the immunoreceptor LILRA3 (Leukocyte immunoglobulin-like receptor A3) has been recently identified as a potential etiopathogenic factor for both pSS susceptibility and lymphomagenesis in the general population.

Objectives: The aim of the current study was to explore the potential contribution of the LILRA3 functional polymorphism in primary SS susceptibility and SS related lymphomagenesis.

Methods: In this study, 346 primary SS patients (ACR/EULAR classification criteria, 2016) and 350 healthy controls were included. The presence of LILRA3 functional polymorphism was detected by polymerase chain reaction (PCR), whereas serum levels were measured by Enzyme-Linked Immunosorbent Assay (ELISA), respectively. Statistical analysis was performed by Graph Pad Prism.

Results: LILRA3 wild type gene variant was detected in 84% of healthy controls, 86.3% of pSS non-Lymphoma patients and 89.7% of patients complicated with lymphoma. Of interest, in pSS patients ≤40 years complicated by lymphoma, LILRA3 wild type variant was detected in 100% in contrast to pSS - non lymphoma patients (81.8%) and healthy controls (HC) (83.2%) [p=0.03]. In line with these results, LILRA3 protein serum levels were increased in pSS-lymphoma patients with pSS onset ≤40 years compared to healthy controls (p=0.009).

Conclusion: Increased LILRA3 protein levels, as a result of predominance of the LILRA3 wild type gene variant in the young pSS onset patients complicated by lymphoma, imply a potential role of the LILRA3 gene in pSS related lymphomagenesis.

Disclosure of Interests: Evangelia Argyriou: None declared, Petros Roussos: None declared, Adrianos Nezos: None declared, Aliki Venetsanopoulou: None declared, Kyriaki A. Boki: None declared, Athanasios Tzioufas: Grant/research support from: ABBVIE, PFIZER, AMGEN, NOVARTIS, GSK, Haralampos M. Moutsopoulos: None declared, Clio Mavragani: None declared

DOI: 10.1136/annrheumdis-2019-eular.6054