Speakers bureau: Abbvie, BMS, Celgene, Janssen, MSD, Novartis, Pfizer, Roche and UCB, Robert G Lambert Consultant for: Bioclinica, Parexel, Abbvie, Ulrich Weber Consultant for: Abbvie, Susanne Juhl Pedersen: None declared, Mikkel Østergaard Grant/research support from: Abbvie, Celgene, Centocor, Merck, Novartis, Consultant for: Abbvie, BMS, Boehringer-Ingelheim, Celgene, Eli Lilly, Hospira, Janssen, Merck, Novartis, Novo, Orion, Pfizer, Regeneron, Roche, and UCB, Speakers bureau: Abbvie, BMS, Boehringer-Ingelheim, Celgene, Eli Lilly, Hospira, Janssen, Merck, Novartis, Novo, Orion, Pfizer, Regeneron, Roche, and UCB, Joachim Sieper Consultant for: Abbvie, Boehringer Ingelheim, Janssen, Lilly, Merck, Mylan, Novartis, Pfizer, UCB., Speakers bureau: Abbvie, Boehringer Ingelheim, Janssen, Lilly, Merck, Mylan, Novartis, Pfizer, UCB., Martin Rudwaleit Consultant for: Abbvie, BMS, Celgene, Janssen, Eli Lilly, MSD, Novartis, Pfizer, Roche, UCB Pharma, Consultant for: Abbvie, BMS, Celgene, Janssen, Eli Lilly, MSD, Novartis, Pfizer, Roche, UCB Pharma, Xenofon Baraliakos Grant/research support from: Abbvie, Boehringer Ingelheim, Bristol-Myers Squibb, Celgene, Centocor, Chugai, Janssen, MSD, Novartis, Pfizer Inc, Roche and UCB, Grant/research support from: Abbvie, Pfizer, Merck Sharp & Dohme, UCB Pharma, Novartis, Consultant for: Abbvie, Bristol-Myers Squibb, Boehringer Ingelheim, Celgene, Chugai, Janssen Biologics, Novartis, Pfizer, UCB Pharma, Galapagos, Speakers bureau: Abbvie, Chugai, Janssen, Novartis, Pfizer, UCB Pharma, Walter P Maksymowych Grant/research support from: Abbvie, Pfizer, Janssen, Novartis, Consultant for: Abbvie, Eli Lilly, Boehringer, Galapagos, Janssen, Novartis, Pfizer and UCB Pharma; Chief Medical Officer for Canadian Research and Education Arthritis

DOI: 10.1136/annrheumdis-2019-eular.7901

THU0367

ANALYSING IMPAIRMENTS IN PHYSICAL PERFORMANCE (AS ASSESSED BY THE AS PERFORMANCE INDEX (ASPI)) IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS

Eerik Ahomaa^{1,2}, Uta Kiltz^{1,2}, Björn Bühring^{1,2}, Xenofon Baraliakos^{1,2}, Salima van Weely³, Juergen Braun^{1,2}, ¹Rheumazentrum Ruhrgebiet, Herne, Germany; ²Ruhr-University, Bochum, Germany; ³Leiden University Medical Center (LUMC), Leiden, Netherlands

Background: In patients with axial spondyloarthritis (axSpA) physical functioning is often impaired. The current gold standard to assess physical functioning is self-reported questionnaires (i.e. BASFI), which can be influenced by patients' subjective feelings. Therefore, a performance-based test-battery was designed to measure physical functioning more objectively: the ankylosing spondylitis (AS) Performance Index (ASPI) [1]. Based on domains taken from BASFI tasks were designed to imitate activities of daily living (ADL). Although the ASPI has been evaluated a thorough analysis of the deficits of physical functioning and factors which influence the performance of patients with axSpA has not been performed to date.

Objectives: The aim of the present study assesses the relation between self-reported assessments of physical functioning and actual performance of patients, and to detect influencing factors.

Methods: Consecutive axSpA patients underwent standardized assessments concentrating on the following variables: patient and disease characteristics, patient-reported outcomes (ASDAS, BASFI, BASMI, ASAS Health Index (ASAS HI), PHQ-9, IPAQ), mSASSS and ASPI (ASPI 1: Bending, 2. Putting on socks, 3. Getting up from the floor) [1]. The performance was measured in seconds as time to complete a task based on published instructions. Impairment in physical performance was defined as inability of patients to perform ≥ 1 ASPI test. Spearman Rho correlation was used to compare self-reported functioning and performed physical functioning. Logistic regression analysis was used to identify factors associated with impaired physical performance.

Results: A total of 200 patients (AS 66%, nr-axSpA 34%) was included: 69% males, 44.3 $\pm$ 12.5 years old, mean symptom duration 17.9 $\pm$ 12.6 years, BMI 27.2 $\pm$ 5.5, mean ASDAS 2.5 $\pm$ 1.1, BASFI 4.0 $\pm$ 2.7, BASMI 3.5 $\pm$ 1.8), ASAS HI 7.0 $\pm$ 4.1, PHQ-9 8.8 $\pm$ 6.2, and mSASSS (n=157) 10.2 $\pm$ 18.8). 133 patients were treated with bDMARDs (66.5%). In total 44 patients (22%) were not able to perform one or more ASPI tests. The mean time for bending was 18.6 $\pm$ 9.5 sec (n=179/90%), for putting on socks 12.8 $\pm$ 6.4 sec (n=156/78%), and for getting up from floor 6.5 $\pm$ 5.0 sec (n=187/94%). A significant correlation was found for all three ASPI-tests with BASFI (0.5-0.7), ASAS HI (0.4-0.6) and spinal mobility as

assessed by BASMI (0.4-0.7). Self-reported physical activity (IPAQ) correlated weakly with ASPI (all 0.2) and structural damage correlated only with the task putting on socks (r=0.3), whereas the other tests did not correlate. Logistic regression showed influence of obesity, spinal mobility and global functioning on actual performance but not of disease activity and self-reported physical function, (Figure 1).

Conclusion: This study confirms a good correlation of the ASPI with standard questionnaires but it showed a substantial floor effect strongly suggesting that additional information on actual performance is needed. Thus, to obtain a complete picture of function and impairments of patients with axSpA the actual performance needs also to be assessed. Moreover, obesity should be addressed as a potential modifying factor contributing to limitation in actual performance.

	Univariable		Multivariable	
	OR (CI)	p	OR (CI)	p
Age	1.08 (1.04-1.11)	<0.001		
Male sex	1.46 (0.68-3.11)	0.332		
BMI	1.17 (1.01-1.35)	<0.001	1.20 (1.09-1.32)	<0.001
ASDAS	2.35 (1.67-3.41)	<0.001		
BASFI 0-10	1.95 (1.58-2.40)	<0.001		
ASAS HI 0-17	1.41 (1.26-1.60)	<0.001	1.40 (1.06-1.86)	0.019
BASMI 0-10	2.44 (1.87-3.19)	<0.001	2.78 (1.77-4.36)	<0.001
PHQ-9 0-27	1.15 (1.08-1.21)	<0.001		

Abstract THU0367 – Figure 1. Association between clinical characteristics of patients with SpA and physical performance

REFERENCE:

- [1] Van Weely SFE, Dekker J, Steultjens MP, van Denderen JC, Nurmo-hamed MT, Dijkmans BAC, van der Horst-Bruinsma IE. Objective evaluation of physical functioning after tumor necrosis factor inhibitory therapy in patients with ankylosing spondylitis: a selection of 3 feasible performance-based tests. J Rheumatol. 2015 Apr;42(4):623-9.

Disclosure of Interests: Eerik Ahomaa: None declared, Uta Kiltz Grant/research support from: Abbvie, Chugai, Eli Lilly, Grünenthal, Janssen, MSD, Novartis, Pfizer, Roche, and UCB., Consultant for: Abbvie, Chugai, Eli Lilly, Grünenthal, Janssen, MSD, Novartis, Pfizer, Roche, and UCB., Björn Bühring Grant/research support from: GE/Lunar and Kinemed., Consultant for: GE/Lunar and Lilly, Xenofon Baraliakos Grant/research support from: Abbvie, Boehringer Ingelheim, Bristol-Myers Squibb, Celgene, Centocor, Chugai, Janssen, MSD, Novartis, Pfizer Inc, Roche and UCB, Grant/research support from: Abbvie, Pfizer, Merck Sharp & Dohme, UCB Pharma, Novartis, Consultant for: Abbvie, Bristol-Myers Squibb, Boehringer Ingelheim, Celgene, Chugai, Janssen Biologics, Novartis, Pfizer, UCB Pharma, Galapagos, Speakers bureau: Abbvie, Chugai, Janssen, Novartis, Pfizer, UCB Pharma, Salima van Weely: None declared, Juergen Braun Shareholder of: Abbvie, BMS, Celgene, Chugai, Merck, Novartis, Pfizer, UCB, Grant/research support from: Abbvie, BMS, Celgene, Chugai, Merck, Novartis, Pfizer, UCB, Grant/research support from: Abbott, Bristol Myers Squibb, Celgene, Celltrion, Chugai, Johnson & Johnson, MSD, Novartis, Pfizer, Roche, UCB Pharma, Grant/research support from: Abbvie, BMS, Celgene, Chugai, Merck, Novartis, Pfizer, UCB, Grant/research support from: Abbvie (Abbott), Amgen, Baxter, Biogen, BMS, Boehringer, Celgene, Celltrion, Centocor, Chugai, Hexal, Janssen, Lilly, Medac, MSD (Schering-Plough), Mylan, Mundipharma, Novartis, Pfizer (Wyeth, Hospira), Roche, Sanofi-Aventis and UCB, Consultant for: Abbvie (Abbott), Amgen, Baxter, Biogen, BMS, Boehringer, Celgene, Celltrion, Centocor, Chugai, Hexal, Janssen, Lilly, Medac, MSD (Schering-Plough), Mylan, Mundipharma, Novartis, Pfizer (Wyeth, Hospira), Roche, Sanofi-Aventis and UCB, Speakers bureau: Abbvie (Abbott), Amgen, Baxter, Biogen, BMS, Boehringer, Celgene, Celltrion, Centocor, Chugai, Hexal, Janssen, Lilly, Medac, MSD (Schering-Plough), Mylan, Mundipharma, Novartis, Pfizer (Wyeth, Hospira), Roche, Sanofi-Aventis and UCB, Speakers bureau: Abbvie, BMS, Celgene, Chugai, Merck, Novartis, Pfizer, UCB, Consultant for: Abbott, Bristol Myers Squibb, Celgene, Celltrion, Chugai, Johnson & Johnson, MSD, Novartis, Pfizer, Roche, UCB Pharma, Speakers bureau: Abbvie, BMS, Celgene, Chugai, Merck, Novartis, Pfizer, UCB, Grant/research support from: Abbvie (Abbott), Amgen, Baxter, Biogen, BMS, Boehringer, Celgene, Celltrion, Centocor, Chugai, Hexal, Janssen, Lilly, Medac, MSD (Schering-Plough), Mylan, Mundipharma, Novartis, Pfizer (Wyeth, Hospira), Roche, Sanofi-Aventis and UCB, Speakers bureau: Abbvie, BMS, Celgene, Chugai, Merck, Novartis, Pfizer, UCB

DOI: 10.1136/annrheumdis-2019-eular.4527