

and final scores of VAS and WOMAC were significantly higher ( $p < 0.05$ ) in study group.

**Conclusion:** The results of the present study showed that virtual reality game-based exercise programs performed better results than conventional treatment program in patients with knee osteoarthritis.

#### REFERENCES:

- [1] Bennell KL, Dobson F, Hinman RS. Exercise in osteoarthritis: moving from prescription to adherence. *Best Pract Res Clin Rheumatol*. 2014; 28(1): 93-117.
- [2] Ju SB, Park GD, Kim SS. Effects of proprioceptive circuit exercise on knee joint pain and muscle function in patients with knee osteoarthritis. *J Phys Ther Sci*. 2015; 27(8): 2439-41.
- [3] Molina KI, Ricci NA, de Moraes SA, Perracini MR. Virtual reality using games for improving physical functioning in older adults: a systematic review. *J Neuroeng Rehabil*. 2014; 11: 156.

**Disclosure of Interests:** None declared

**DOI:** 10.1136/annrheumdis-2019-eular.3749

THU0422

#### COMPARISON OF PRP DERIVED GROWTH FACTOR (PRGF) VERSUS HYALURONIC ACID (HA) IN MILD TO MODERATE KNEE OSTEOARTHRITIS; A SINGLE BLIND ONE YEAR RANDOMIZED CLINICAL TRIAL STUDY

Seyed Ahmad Raeissadat<sup>1</sup>, Azadeh Gharooee Ahangar<sup>2</sup>, Seyed Mansoor Rayegani<sup>1</sup>, Pegah Yavari<sup>1</sup>. <sup>1</sup>Shahid Beheshti University of medical sciences, physical medicine and rehabilitation, Tehran, Iran (Islamic Republic of); <sup>2</sup>Zanjan University of medical Sciences, physical medicine and rehabilitation, Zanjan, Iran (Islamic Republic of)

**Background:** Osteoarthritis (OA) is the most common joint disease with characteristics of progressive loss of joint cartilage.

**Objectives:** Aim of this study was to evaluate clinical outcomes of intra-articular injection of PRP derived growth factor (PRGF) versus hyaluronic acid (HA) in patients with knee osteoarthritis.


**Methods:** 102 patients with grade II or grade III knee OA were randomly assigned to 2 intra-articular injections of PRGF 3 weeks apart or 3 weekly injections of HA. Primary outcome was the mean change from baseline until 2, 6 and 12 months post intervention in scores of visual analog scale (VAS), WOMAC and Lequesne index.

**Results:** The mean age of patients was 57.08±7.3 in PRGF compared to 58.63±7.09 in HA group. In PRGF group, VAS decreased from 7.8±1.5 to 4.5±1.7, and from 7.8±1.1 to 6.1±1.8 in the HA group after 12 months ( $P < 0.0001$ ). Total WOMAC score decreased from 41.96±11.71 to 27.10±12.3 ( $P = 0.02$ ), and from 39.71±10.4 to 32.41±11.8 after 12 months, respectively ( $P > 0.05$ ). In Lequesne index, all scores were significantly decreased after 12 months in PRGF group in comparison to HA group ( $P < 0.001$ ).

**Conclusion:** Although PRGF and HA were both effective, but PRGF injection resulted in significantly higher satisfaction, lower VAS and Lequesne pain score and improvement in function of patients with symptomatic mild to moderate knee osteoarthritis.

#### REFERENCES:

- [1] Raeissadat SA, Rayegani SM, Tabibian E, Rahimi Dehgolan S. Intra-articular ozone injection efficacy in knee osteoarthritis: A systematic review with meta-analysis. *Annals of Physical and Rehabilitation Medicine*. 2018;61:e132-e3.
- [2] Raeissadat SA, Rayegani SM, Hassanabadi H, Fathi M, Ghorbani E, Babaei M, et al. Knee Osteoarthritis Injection Choices: Platelet-Rich Plasma (PRP) versus Hyaluronic Acid (A one-year randomized clinical trial). *Clinical Medicine Insights: Arthritis and Musculoskeletal Disorders*. 2015;8:CMAMD.S17894.
- [3] Raeissadat SA, Rayegani SM, Ahangar AG, Abadi PH, Mojjani P, Ahangar OG. Efficacy of Intra-articular Injection of a Newly Developed Plasma Rich in Growth Factor (PRGF) Versus Hyaluronic Acid on Pain and Function of Patients with Knee Osteoarthritis: A Single-Blinded Randomized Clinical Trial. *Clinical medicine insights Arthritis and musculoskeletal disorders*. 2017;10:1179544117733452.



**Abstract THU0422 – Figure 1.** Mean values of Lequesne total at baseline, 2, 6 and 12 months after injection. PRGF indicates plasma rich in growth factor

**Disclosure of Interests:** None declared

**DOI:** 10.1136/annrheumdis-2019-eular.2294

THU0423

#### ANTERIOR TIBIOTALAR FAT PAD MORPHOLOGY AND SIGNAL INTENSITY ON MAGNETIC RESONANCE IMAGING ARE CORRELATED WITH PATIENT CHARACTERISTICS AND JOINT PATHOLOGY

Stijn Arnaert<sup>1</sup>, Paul Byttebier<sup>2</sup>, Sam Van Rossum<sup>3</sup>, Evie Vereecke<sup>4</sup>, Ilse Jonkers<sup>3</sup>, Edwin Oei<sup>5</sup>, S.M.A. Bierma-Zeinstra<sup>6</sup>, Rik Lories<sup>7,8</sup>, Marienke Van Middelkoop<sup>6</sup>, Stefan Clockaerts<sup>2,7</sup>. <sup>1</sup>KU Leuven, Leuven, Belgium; <sup>2</sup>AZ Groeninge, Orthopaedic Surgery and Traumatology, Kortrijk, Belgium; <sup>3</sup>KU Leuven, Human Movement Biomechanics Research Group, Leuven, Belgium; <sup>4</sup>KU Leuven (KULAK), Development and Regeneration, Kortrijk, Belgium; <sup>5</sup>Erasmus MC Medical University Rotterdam, Radiology, Rotterdam, Netherlands; <sup>6</sup>Erasmus MC Medical University Rotterdam, General Practice, Rotterdam, Netherlands; <sup>7</sup>KU Leuven, Skeletal Biology and Engineering Research Center, Leuven, Belgium; <sup>8</sup>University Hospitals Leuven, Rheumatology, Leuven, Belgium

**Background:** Ankle sprains are the most frequent form of trauma in the ankle and up to 33% of patients experience ongoing pain 1 year after the initial trauma.<sup>1</sup> In the ankle, trauma is the primary etiology of osteoarthritis with an overwhelming proportion of 70-78%.<sup>2</sup> Recently, our group completed a small pilot study that suggested that the anterior tibiotalar fat pad (ATFP) should be investigated as a source of inflammation and pain.<sup>3</sup>

**Objectives:** In this study, we tried to investigate the innovative concept of the ATFP as missing link in the pathogenesis of persistent complaints and potential source driving inflammation in the development of osteoarthritis.

**Methods:** The present study is a secondary analysis of an observational case control study by Van Ochten et al.<sup>4</sup> We included 106 patients with a Kellgren & Lawrence score of 0 in the tibiotalar joint on x-ray. T1 MRI scans were assessed for the signal intensity and area of the ATFP by mapping the fat pad in 'Mimics 18.0'. After importing those mapped scans in the program 'MATLAB', quantitative values of intensity and area were generated. Those values were statistically tested for correlations with patient characteristics and structural abnormalities by univariate and multivariate linear regression.

**Results:** MRI signal intensity of the ATFP is associated with BMI ( $p = 0.03$ ), sex ( $p < 0.01$ ) and age ( $p = 0.01$ ). ATFP area is correlated with sex ( $p < 0.01$ ) and presence of pre-OA signs in the subtalar joint ( $p = 0.01$ ). After multivariate analysis, correcting for sex, subtalar pre-OA signs and BMI, persistent complaints were associated with ATFP area ( $p = 0.04$ ).

**Conclusion:** This study demonstrates the involvement of the ATFP in hindfoot joint pathology. ATFP MRI characteristics were also influenced by patient characteristics. Further research should confirm these findings in a more elaborate population including OA patients, focus on histological validation and determine underlying pathogenic processes that may explain the observed correlations.

#### REFERENCES:

- [1] van Rijn, R. M. et al. What Is the Clinical Course of Acute Ankle Sprains? A Systematic Literature Review. *American Journal of Medicine* 121, (2008).
- [2] Valderrabano, V., Horisberger, M., Russell, I., Dougall, H. & Hintermann, B. Etiology of ankle osteoarthritis. *Clinical Orthopaedics and Related Research* 467, 1800–1806 (2009).
- [3] Clockaerts, S. et al. The anterior tibiotalar fat pad as a source of pain and inflammation in osteoarthritis of the ankle: anatomy, histology and imaging. *Osteoarthr. Cartil.* 26, S120 (2018).

- [4] Van Ochten, J. M. et al. Structural abnormalities and persistent complaints after an ankle sprain are not associated: An observational case control study in primary care. *Br. J. Gen. Pract.* 64, e545–e553 (2014).

**Disclosure of Interests:** Stijn Arnaert: None declared, Paul Byttebier: None declared, Sam Van Rossum: None declared, Evie Vereecke: None declared, Ilse Jonkers: None declared, Edwin Oei: None declared, S.M.A. Bierma-Zeinstra: None declared, Rik Lories Consultant for: Abbvie, Celgene, Eli-Lilly, Janssen, Merck, Novartis, Pfizer, UCB, Marienke Van Middekoop: None declared, Stefan Clockaerts: None declared

DOI: 10.1136/annrheumdis-2019-eular.3650

**THU0424 THE NEW TREATMENT APPROACH IN KNEE OSTEOARTHRITIS: EFFICACY OF CELLULAR MATRIX COMBINATION OF PLATELET RICH PLASMA WITH HYALURONIC ACID VERSUS TWO DIFFERENT TYPES OF HYALURONIC ACID (HA) (PROSPECTIVE, RANDOMIZED, DOUBLE BLIND CONTROL STUDY)**

Branko Barac, Nemanja Damjanov, Ana Zekovic. *Institute of rheumatology, Belgrade, Serbia*

**Background:** Osteoarthritis pathogenesis is a complex process associated with decreased ability to regenerate cartilage mainly due to lack of physiological vascularization. One of the most commonly affected joints is the knee.

**Objectives:** The aim of this study was to compare the efficacy of intra-articular (IA) injections of platelet rich plasma (PRP) combined with hyaluronic acid (HA) prepared with the Cellular Matrix device versus IA injections with two different types of hyaluronic acid for treatment of knee osteoarthritis.

**Methods:** This is a prospective, randomized, double-blind, controlled study on 53 patients (90 knees) suffering from knee osteoarthritis, divided in 3 groups. The first group comprised 19 patients (30 knees) treated with 3 IA injections, one every second week, of Cellular Matrix (CM) PRP-HA combination. The second group of 19 patients (30 knees) was treated with 3 weekly IA injections of 2% non-cross-linked sodium hyaluronate and the third group of 15 patients (30 knees) treated with 3 weekly IA injections of 2% non-cross-linked sodium hyaluronate with mannitol. All groups were homogeneous concerning gender, age and Kellgren Lawrence scale (I to III). For all patients visual analog pain scale (VAS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Knee Injury and Osteoarthritis Outcome Score (KOOS), The International Knee Documentation Committee (IKDC) score ("well-being" scale for all 4 scores between 0 and 100) and ultrasound (US) cartilage thickness on lateral, trochlear, and medial compartments, with normal range values from 2 to 2.5 mm, were measured at the beginning of the treatment (baseline) and at each follow up visit, that is at 2, 6 and 12 months after the last injection.

**Results:** A statistically significant difference ( $p < 0.05$ ) in the CM group was found compared to AV and OP group in the values of VAS, WOMAC, KOOS and IKDC after two months, although an improvement, compared to baseline values, was observed for the indicated parameters in all groups. A high statistically significant difference ( $p < 0.01$ ) was obtained in the CM group compared to the AV and OP group for VAS, WOMAC, KOOS and IKDC after 6 and 12 months. In both groups of patients treated with hyaluronic acid, a deterioration of values for VAS, WOMAC, KOOS and IKDC score was seen at 12 months in relation to values at 6 months. The CM treated group showed statistically significant improvement ( $p < 0.05$ ) of the cartilage thickness after 2, 6 and 12 months in the medial and highly statistically significant improvement ( $p < 0.01$ ) in the lateral segments of knee cartilage in comparison to baseline values.

**Conclusion:** The Cellular Matrix PRP-HA combination might be one of the most potent, safe, fast and novel therapeutic option for osteoarthritis of the knee, as well as a useful tool for postponing arthroplasty surgery when it is necessary.

**REFERENCES:**

- [1] Saturveithan C, Premganes G, Fakhrazzaki S, et al. Intra-articular Hyaluronic Acid (HA) and platelet rich plasma (PRP) injections versus Hyaluronic acid (HA) injections in grade III and IV knee osteoarthritis (OA) patients: A retrospective study on functional outcome. 45th Malaysian Orthopedic Association(2015).
- [2] Chen SH, Khuan TS, Kao MJ, et al. Clinical effectiveness in severe knee osteoarthritis after intra-articular platelet rich plasma therapy in association

with hyaluronic acid injection: three case reports. *Clin. Interv.Aging.* 11, 1213-1219 (2016).

3. Renevier JL, Marc JF, Adam P, et al. "Cellular Matrix TM PRP-HA": A new treatment option with platelet-rich plasma and hyaluronic acid for patients with osteoarthritis having had an unsatisfactory clinical response to hyaluronic acid alone: Results of a pilot multicenter French study with long-term follow-up. *Int J Clin Rheumatology* 2018; 13(4), 226-229

**Disclosure of Interests:** Branko Barac: None declared, Nemanja Damjanov Grant/research support from: AbbVie, Pfizer and Roche, Consultant for: Abbvie, Gedeon Richter, Merck, Novartis, Pfizer and Roche., Speakers bureau: Abbvie, Gedeon Richter, Merck, Novartis, Pfizer and Roche., Ana Zekovic: None declared

DOI: 10.1136/annrheumdis-2019-eular.5018

**THU0425 COST-EFFECTIVE OA TRIALS REQUIRE ENROLMENT OF KNEES WITH DEFINITE JOINT SPACE NARROWING (KELLGREN LAWRENCE 3); DATA FROM 6,939 KNEES FROM THE OSTEOARTHRITIS INITIATIVE**

Michael Bowes<sup>1</sup>, Alan Brett<sup>1</sup>, Philip G. Conaghan<sup>2</sup>. <sup>1</sup>Imorphics Ltd, Manchester, United Kingdom; <sup>2</sup>University of Leeds, Leeds Institute of Rheumatic and Musculoskeletal Medicine, Leeds, United Kingdom

**Background:** The design of clinical trials for osteoarthritis is challenging; structural changes in tissues are quantitatively small and proceed very slowly. No clear guidance exists on how to optimise recruitment. We have previously shown that the use of radiographic joint space width of 2 to 4.5 mm is of major importance for improving responsiveness in clinical trials using MRI bone and cartilage outcomes. However, it can be technically challenging to screen for joint space width this carefully, so we considered whether Kellgren-Lawrence (KL) grade could be used as an enrichment strategy. As no other commonly used covariates have been shown to reliably increase responsiveness, it would be useful to know the numbers needed using ONLY an expert-read KL grade structural inclusion criterion.

**Objectives:** To determine responsiveness of change in femur bone shape and cartilage thickness using a large observational dataset and calculate likely trial cohort sizes per arm for each KL grade.

**Methods:** We used all knees from the Osteoarthritis Initiative which had MR images at baseline, 1 and 2 years, and a baseline KL grade (centrally read and adjudicated by 2 experienced radiologists). Quantitative 3D femur bone shape, and cartilage thickness in the central medial femoral region were used as outcome measures. Responsiveness was assessed using standardised response means "SRM" (CIs were assessed using the bootstrap method of Efron) and derived the number of patients per arm in a putative trial to demonstrate 50% change, at 80% probability,  $\alpha=0.05$ .

**Results:** 6,945 knees (3,667 subjects, 2,085 female) were included (KL 0: 2,798; KL1: 1,338 knees; KL2: 1,879 knees; KL3: 924 knees). Table 1 provides summary results and Figure 1 shows SRM and putative trial cohort numbers by KL grade for bone shape and cartilage thickness. Femur bone shape had higher SRM values at all timepoints, typically twice that of SRM for cartilage thickness in all KL groups. Cohort size when using cartilage thickness increased significantly between KL2 and KL3.

**Conclusion:** Expert-read KL3 inclusion, using 2 independent radiologists, with strict attention to standards provides increased responsiveness for common MRI OA outcomes. It is worth noting that 80% of KL3 knees have OARSI JSN grade 3, and 78% of KL3 knees have radiographic JSW of 2 to 4.5 mm, and similar enrichment can be expected using these alternatives. Few clinical studies can afford cohort sizes of greater than 200 knees, and for these studies, using a large proportion of KL3 knees is critical, especially if cartilage thickness is to be used as the outcome.