- [4] Van Ochten, J. M. et al. Structural abnormalities and persistent complaints after an ankle sprain are not associated: An observational case control study in primary care. *Br. J. Gen. Pract.* 64, e545–e553 (2014).

Disclosure of Interests: Stijn Arnaert: None declared, Paul Byttebier: None declared, Sam Van Rossum: None declared, Evie Vereecke: None declared, Ilse Jonkers: None declared, Edwin Oei: None declared, S.M.A. Bierma-Zeinstra: None declared, Rik Lories Consultant for: Abbvie, Celgene, Eli-Lilly, Janssen, Merck, Novartis, Pfizer, UCB, Marienke Van Middekoop: None declared, Stefan Clockaerts: None declared

DOI: 10.1136/annrheumdis-2019-eular.3650

THU0424

THE NEW TREATMENT APPROACH IN KNEE OSTEOARTHRITIS: EFFICACY OF CELLULAR MATRIX COMBINATION OF PLATELET RICH PLASMA WITH HYALURONIC ACID VERSUS TWO DIFFERENT TYPES OF HYALURONIC ACID (HA) (PROSPECTIVE, RANDOMIZED, DOUBLE BLIND CONTROL STUDY)

Branko Barac, Nemanja Damjanov, Ana Zekovic. *Institute of rheumatology, Belgrade, Serbia*

Background: Osteoarthritis pathogenesis is a complex process associated with decreased ability to regenerate cartilage mainly due to lack of physiological vascularization. One of the most commonly affected joints is the knee.

Objectives: The aim of this study was to compare the efficacy of intra-articular (IA) injections of platelet rich plasma (PRP) combined with hyaluronic acid (HA) prepared with the Cellular Matrix device versus IA injections with two different types of hyaluronic acid for treatment of knee osteoarthritis.

Methods: This is a prospective, randomized, double-blind, controlled study on 53 patients (90 knees) suffering from knee osteoarthritis, divided in 3 groups. The first group comprised 19 patients (30 knees) treated with 3 IA injections, one every second week, of Cellular Matrix (CM) PRP-HA combination. The second group of 19 patients (30 knees) was treated with 3 weekly IA injections of 2% non-cross-linked sodium hyaluronate and the third group of 15 patients (30 knees) treated with 3 weekly IA injections of 2% non-cross-linked sodium hyaluronate with mannitol. All groups were homogeneous concerning gender, age and Kellgren Lawrence scale (I to III). For all patients visual analog pain scale (VAS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Knee Injury and Osteoarthritis Outcome Score (KOOS), The International Knee Documentation Committee (IKDC) score ("well-being" scale for all 4 scores between 0 and 100) and ultrasound (US) cartilage thickness on lateral, trochlear, and medial compartments, with normal range values from 2 to 2.5 mm, were measured at the beginning of the treatment (baseline) and at each follow up visit, that is at 2, 6 and 12 months after the last injection.

Results: A statistically significant difference ($p < 0.05$) in the CM group was found compared to AV and OP group in the values of VAS, WOMAC, KOOS and IKDC after two months, although an improvement, compared to baseline values, was observed for the indicated parameters in all groups. A high statistically significant difference ($p < 0.01$) was obtained in the CM group compared to the AV and OP group for VAS, WOMAC, KOOS and IKDC after 6 and 12 months. In both groups of patients treated with hyaluronic acid, a deterioration of values for VAS, WOMAC, KOOS and IKDC score was seen at 12 months in relation to values at 6 months. The CM treated group showed statistically significant improvement ($p < 0.05$) of the cartilage thickness after 2, 6 and 12 months in the medial and highly statistically significant improvement ($p < 0.01$) in the lateral segments of knee cartilage in comparison to baseline values.

Conclusion: The Cellular Matrix PRP-HA combination might be one of the most potent, safe, fast and novel therapeutic option for osteoarthritis of the knee, as well as a useful tool for postponing arthroplasty surgery when it is necessary.

REFERENCES:

- [1] Saturveithan C, Premganes G, Fakhrazaki S, et al. Intra-articular Hyaluronic Acid (HA) and platelet rich plasma (PRP) injections versus Hyaluronic acid (HA) injections in grade III and IV knee osteoarthritis (OA) patients: A retrospective study on functional outcome. 45th Malaysian Orthopaedic Association(2015).
- [2] Chen SH, Khuan TS, Kao MJ, et al. Clinical effectiveness in severe knee osteoarthritis after intra-articular platelet rich plasma therapy in association

with hyaluronic acid injection: three case reports. *Clin. Interv. Aging.* 11, 1213-1219 (2016).

3. Renevier JL, Marc JF, Adam P, et al. "Cellular Matrix TM PRP-HA": A new treatment option with platelet-rich plasma and hyaluronic acid for patients with osteoarthritis having had an unsatisfactory clinical response to hyaluronic acid alone: Results of a pilot multicenter French study with long-term follow-up. *Int J Clin Rheumatology* 2018; 13(4), 226-229

Disclosure of Interests: Branko Barac: None declared, Nemanja Damjanov Grant/research support from: AbbVie, Pfizer and Roche, Consultant for: Abbvie, Gedeon Richter, Merck, Novartis, Pfizer and Roche., Speakers bureau: Abbvie, Gedeon Richter, Merck, Novartis, Pfizer and Roche., Ana Zekovic: None declared

DOI: 10.1136/annrheumdis-2019-eular.5018

THU0425

COST-EFFECTIVE OA TRIALS REQUIRE ENROLMENT OF KNEES WITH DEFINITE JOINT SPACE NARROWING (KELGREN LAWRENCE 3); DATA FROM 6,939 KNEES FROM THE OSTEOARTHRITIS INITIATIVE

Michael Bowes¹, Alan Brett¹, Philip G. Conaghan². ¹Imorphics Ltd, Manchester, United Kingdom; ²University of Leeds, Leeds Institute of Rheumatic and Musculoskeletal Medicine, Leeds, United Kingdom

Background: The design of clinical trials for osteoarthritis is challenging; structural changes in tissues are quantitatively small and proceed very slowly. No clear guidance exists on how to optimise recruitment. We have previously shown that the use of radiographic joint space width of 2 to 4.5 mm is of major importance for improving responsiveness in clinical trials using MRI bone and cartilage outcomes. However, it can be technically challenging to screen for joint space width this carefully, so we considered whether Kellgren-Lawrence (KL) grade could be used as an enrichment strategy. As no other commonly used covariates have been shown to reliably increase responsiveness, it would be useful to know the numbers needed using ONLY an expert-read KL grade structural inclusion criterion.

Objectives: To determine responsiveness of change in femur bone shape and cartilage thickness using a large observational dataset and calculate likely trial cohort sizes per arm for each KL grade.

Methods: We used all knees from the Osteoarthritis Initiative which had MR images at baseline, 1 and 2 years, and a baseline KL grade (centrally read and adjudicated by 2 experienced radiologists). Quantitative 3D femur bone shape, and cartilage thickness in the central medial femoral region were used as outcome measures. Responsiveness was assessed using standardised response means "SRM" (CIs were assessed using the bootstrap method of Efron) and derived the number of patients per arm in a putative trial to demonstrate 50% change, at 80% probability, $\alpha=0.05$.

Results: 6,945 knees (3,667 subjects, 2,085 female) were included (KL 0: 2,798; KL1: 1,338 knees; KL2: 1,879 knees; KL3: 924 knees). Table 1 provides summary results and Figure 1 shows SRM and putative trial cohort numbers by KL grade for bone shape and cartilage thickness. Femur bone shape had higher SRM values at all timepoints, typically twice that of SRM for cartilage thickness in all KL groups. Cohort size when using cartilage thickness increased significantly between KL2 and KL3.

Conclusion: Expert-read KL3 inclusion, using 2 independent radiologists, with strict attention to standards provides increased responsiveness for common MRI OA outcomes. It is worth noting that 80% of KL3 knees have OARSI JSN grade 3, and 78% of KL3 knees have radiographic JSW of 2 to 4.5 mm, and similar enrichment can be expected using these alternatives. Few clinical studies can afford cohort sizes of greater than 200 knees, and for these studies, using a large proportion of KL3 knees is critical, especially if cartilage thickness is to be used as the outcome.

Abstract THU0425 – Figure 1. SRM and cohort numbers by KL grade for bone shape and cartilage thickness in 6,939 knees. Top row shows SRM values at 1 year (left) and 2 years for femur bone shape and central medial femur cartilage thickness, with 95% confidence limits calculated using a bootstrap method. Bottom row shows numbers needed per cohort arm, assuming 50% change, 80% probability, alpha 0.05. Units are in vector units for bone shape, and mm for cartilage thickness.

Abstract THU0425 – Table 1. Responsiveness of bone shape and cartilage thickness by KL Grade

B score Change (femur)		1 year				2 year			
	SRM (±95% CI)	Cohort Size	Mean Change	SD	SRM (±95% CI)	Cohort Size	Mean Change	SD	
KL 0	0.21 (0.17, 0.24)	1177 (851, 1808)	0.030	0.147	0.48 (0.44, 0.51)	218 (188, 258)	0.090	0.189	
KL 2	0.34 (0.29, 0.39)	421 (324, 578)	0.064	0.186	0.56 (0.51, 0.61)	156 (132, 188)	0.158	0.281	
KL 2	0.55 (0.52, 0.59)	162 (141, 185)	0.125	0.225	0.75 (0.71, 0.79)	88 (79, 98)	0.259	0.344	
KL 3	0.78 (0.68, 0.85)	82 (69, 106)	0.251	0.322	0.96 (0.9, 1.03)	54 (47, 62)	0.459	0.477	

Cartilage Thickness Change (cmf region)		1 year				2 year			
	SRM (±95% CI)	Cohort Size	Mean Change	SD	SRM (±95% CI)	Cohort Size	Mean Change	SD	
KL 0	-0.05 (-0.09, -0.01)	17857 (51065, 6130)	-0.004	0.068	-0.24 (-0.27, -0.2)	875 (1202, 663)	-0.019	0.079	
KL 1	-0.17 (-0.22, -0.12)	1765 (969, 1063)	-0.016	0.093	-0.32 (-0.36, -0.27)	488 (683, 378)	-0.037	0.117	
KL 2	-0.21 (-0.17, -0.25)	1132 (1798, 787)	-0.023	0.110	-0.35 (-0.31, -0.39)	409 (514, 328)	-0.049	0.140	
KL 3	-0.34 (-0.4, -0.28)	433 (636, 314)	-0.054	0.161	-0.52 (-0.58, -0.46)	186 (235, 148)	-0.107	0.207	

Detailed breakdown of SRM values, cohort size, mean change, and SD of that change for Figure 1. Units are in vector units for bone shape, and mm for cartilage thickness.

Disclosure of Interests: Michael Bowes Shareholder of: Stryker Corporation, Employee of: Stryker Corporation, Alan Brett Shareholder of: Stryker Corp, Employee of: Stryker Corp, Philip G Conaghan Consultant for: Flexion Therapeutics, AbbVie, Medivir, Merck Serono, Novartis, GlaxoSmithKline

DOI: 10.1136/annrheumdis-2019-eular.6992

THU0426 EFFECTS OF SODIUM SUCCINATE AND HYALURONIC ACID IN KNEE OSTEOARTHRITIS TREATMENT

Oleksandr Burianov¹, Liudmyla Khimion², Taras Omelchenko¹, Halyna Havryliuk².
¹Bogomolets National Medical University, Traumatology and Orthopedic, Kyiv, Ukraine; ²Shupyk National Medical Academy of Postgraduate Education, Family Medicine, Kyiv, Ukraine

Background: At present hyaluronic acid (HA) is rather widely used in treatment of patients with osteoarthritis (OA). HA normalizes the properties of the synovial fluid; has a protective effects; promotes the cartilage nutrition and so improves the signs of OA and function of the joints. Sodium succinate (the salt of the succinic acid) helps to normalize intracellular metabolism and tissue respiration in hypoxic conditions via mitochondrial mechanism of action; normalizes acidic – alkaline balance; takes part in K⁺ and Ca²⁺ transportation and provides antioxidant defense, - so is a promising compound for cartilage treatment

Objectives: To investigate the clinical efficacy of combination of hyaluronate and sodium succinate in treatment of early OA stages

Methods: The study included 126 patients with knee OA (stages I-II, Kelgren and Lawrence), mean age (54.3 ± 2.7) years, among them - 75 women (60%) and 51 men - (40%). All enrolled patients had OA exacerbation (without clinically evident synovitis) and received standard OA treatment (NSAIDs, exercises, orthopedic devices) for 15 days; Gr.1 patients (58) also got 5 intra-articular injections of 1.1% hyaluronic acid, stabilized with sodium succinate (2 ml, once a week); patients of Gr.2 (68) in addition to standard treatment received 5 intra-articular injections of 1.1% solution of non-stabilized HA (2 ml, once a week). Clinical observation and evaluation of the results were performed at the beginning of the treatment, at 6th, 12th and 24th week after the study beginning

Results: During the treatment period, patients in both groups showed the positive changes in clinical signs and symptoms of OA which led to the lowering of the general WOMAC index (from (78.3 ± 4.1) in Gr. 1 and (75.4 ± 3.8) in Gr. 2 at the beginning of the study to (27.9 ± 2.6) and (29.8 ± 1.9) accordingly at week 12 (p<0,05 for both groups). The VAS score in both groups indicated a significant pain reduction, but the stability and duration of the clinical effect in the groups was different. In patients of Gr.1, the pain syndrome continued to decrease after 12 weeks till 24th week, whereas in Gr.2 after the treatment course there was no significant changes in further pain regression after 6th week point. The changes in Lisholm score were also significantly better in Gr.1 than in Gr. 2 (before treatment (21.7 ± 4.6) and (22.6 ± 5.3), at week 6 - (86.4 ± 5.7) and (71.3 ± 4.8), at week 12 - (87.6 ± 6.2) and (63.8 ± 5.3), respectively, p<0,05 between groups at week 12th

Conclusion: Combination of the hyaluronic acid and sodium succinate biochemical and physiologic properties in early stages of knee OA (as intra-articular injections) allows to increase the treatment efficacy, achieve better pain control, and more stable remission comparing to use of the hyaluronic acid alone

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.6120

THU0427 THE LIFETIME RISK OF KNEE AND HIP REPLACEMENT FOLLOWING A GP DIAGNOSIS OF OSTEOARTHRITIS: REAL WORLD COHORT DATA FROM CATALONIA INCLUDING 48,311 PARTICIPANTS WITH UP TO 9 YEARS OF FOLLOW-UP

Edward Burn¹, David W Murray², Gillian A Hawker^{3,4,5}, Rafael Pinedo-Villanueva¹, Daniel Prieto-Alhambra^{1,6}. ¹Centre for Statistics in Medicine, NDORMS, University of Oxford, Oxford, United Kingdom; ²NDORMS, University of Oxford, Oxford, United Kingdom; ³Women's College Hospital, Women's College Research Institute, Toronto, Canada; ⁴Institute for Clinical Evaluative Sciences, Toronto, Canada; ⁵Department of Medicine, University of Toronto, Toronto, Canada; ⁶GREMPAL Research Group, Idiap Jordi Gol and CIBERFes, Universitat Autònoma de Barcelona and Instituto de Salud Carlos III, Barcelona, Spain

Background: The lifetime risk of knee and hip replacement following a primary care diagnosis of knee or hip osteoarthritis is unknown and overly negative beliefs about prognosis act as a barrier to implementing recommended management strategies.

Objectives: To estimate lifetime risk of knee and hip replacement following a GP diagnosis of osteoarthritis and assess how this risk varies with patient characteristics.

Methods: Primary care and linked hospital data from Catalonia, covering 2006 to 2015, were used. Study participants had a newly recorded GP diagnosis of knee or hip osteoarthritis. Parametric survival models were specified for risk of knee/hip replacement and death following diagnosis. Survival models were combined and extrapolated using a Markov model and lifetime risk estimated for the average patient profile. The effects of age at diagnosis, sex, comorbidities, socioeconomic status, body mass index (BMI), and smoking on risk were assessed.

Results: 48,311 individuals diagnosed with knee osteoarthritis were included with a median follow-up of 4.3 years (IQR: 2.1 to 6.5) and of whom 2,561 underwent knee replacement. Respective figures for hip osteoarthritis were 15,105 individuals diagnosed with a median follow-up of 3.8 years (IQR: 1.8 to 6.1) and 1,247 hip replacements. The average participant's lifetime risk for knee replacement was 30% (95% CI: 25% to 36%) and for hip replacement was 14% (10% to 19%). Notable patient characteristics influencing lifetime risk were age at diagnosis for knee and hip replacement, sex for hip replacement, and BMI for knee replacement. BMI increasing from 25 to 35 was associated with lifetime risk of knee replacement increasing from 24% (20% to 28%) to 32% (26% to 37%) for otherwise average patients.

Conclusion: Knee and hip replacement are not bound to happen for most after a GP diagnosis of osteoarthritis, with average lifetime risks of less than a third and a sixth, respectively. Patient characteristics influence lifetime risks with, most notably higher BMI associated with a meaningfully increased risk of knee replacement.

Disclosure of Interests: : Edward Burn: None declared, David W Murray Grant/research support from: Grants from Zimmer Biomet, Consultant for: Personal fees from Zimmer Biomet, Gillian A Hawker: None declared, Rafael Pinedo-Villanueva: None declared, Daniel Prieto-Alhambra Grant/research support from: Grants from Amgen, UCB Biopharma and Servier outside the submitted work, Consultant for: UCB Biopharma, Speakers bureau: Amgen

DOI: 10.1136/annrheumdis-2019-eular.5401