FRI0036

EFFECTS OF DISEASE ACTIVITY, ILLNESS PERCEPTION, SOCIAL SUPPORT AND COPING METHODS ON QUALITY OF LIFE IN THE PATIENTS WITH RHEUMATOID ARTHRITIS AND FIBROMYALGIA SYNDROME

Hakan Kesmen¹, Ahmet Kivanc Cengiz², Ayhan Bilgili². ¹Samsun Education and Research Hospital, Physical Medicine and Rehabilitation, Samsun, Turkey; ²19 Mayıs University Faculty of Medicine, Physical Medicine and Rehabilitation, Samsun, Turkey

Background: In chronic diseases, social support, illness perception and coping methods have important effect on quality of life.

Objectives: In the present study, we aimed to compare the quality of life (QOL) of the patients with rheumatoid arthritis (RA) with fibromyalgia (FMS) patients and healthy controls and to assess the relationship between QOL and illness perception, social support, disease coping methods in the patients with RA and FMS.

Methods: Fifty-eight patients with RA, 50 patients with FMS and 50 healthy controls were enrolled in the study. Pain (Visual Analogue Scale, VAS), QOL (Short Form SF-36), illness perception (Revised Illness Perception Scale, IPQ-R), social support (Multidimensional Perceived Social Support Scale, MSPSS) were assessed in the patients. In RA patients, disease activity was evaluated with DAS 28 and Clinical Disease Activity (CDAI), and functional status was evaluated with Health Assessment Questionnaire (HAQ). Fibromyalgia Impact Questionnaire (FIQ) scale was used in the clinical assessment of FMS patients.

Results: While RA and FMS patients had higher COPE-emotional and COPE-problem scores than the healthy controls ($p < 0.05$), there was no significant difference between the patients with RA and FMS ($p > 0.05$). Regarding the all MSPSS scores, there was no significant difference between the three groups ($p > 0.05$). FMS patients had lower scores than the RA patients and healthy controls regarding the physical function, pain, social functioning and mental health subscales of SF-36 ($p < 0.05$).

In RA patients, MSPSS-friend and MSPSS-special one scores were positively correlated with all subscales of SF-36. IPQ-R consequences scores were negatively associated with all subscales of SF-36 ($p < 0.01$). There were positive correlations between IPQ-R treatment control and SF-36 physical function and social function subscales ($p < 0.05$).

In FMS patients, IPQ-R-timeline acute/chronic score was negatively correlated with physical function, pain and social function subscales ($p < 0.05$). While IPQ-R timeline cyclical was negative correlated with SF-36 mental health subscale, COPE-emotional was negatively correlated with SF-36 general health.

Conclusion: In RA and FMS patients, QOL and physical functions were found to be related with illness perception, social support and coping methods as much as disease activity.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.8039

FRI0037

DIAGNOSTIC PERFORMANCE OF CURRENT CVR SCORES FOR THE DIAGNOSIS OF CAROTID PLAQUE IN MEXICAN-MESTIZO RA SUBJECTS

Dionicio Ángel Galarza-Delgado¹, Iris Jazmin Colunga-Pedraza¹, José Ramón Azpiri-López², Karla Paola Cuéllar-Calderón¹, Ileana Cecilia Reynosa-Silva¹, Marielva Castro-González¹, Carolina Marlene Martínez-Flores², José Dionisio Castillo-Ortiz³, Alberto Cárdenas¹. ¹Hospital Universitario "Dr. José Eleuterio González", UANL, Rheumatology, Monterrey, Mexico; ²Hospital Universitario "Dr. José Eleuterio González", UANL, Cardiology, Monterrey, Mexico; ³Unidad de Artritis y Reumatismo, Guadalajara, Mexico

Background: Rheumatoid arthritis (RA) is an inflammatory disease with an increased mortality compared to the general population. The measurement of carotid intima media thickness (CIMT) is used as an independent marker of subclinical atherosclerosis.

Objectives: To determine the prevalence of carotid plaque (CP) in a cohort of Mexican RA subjects and diagnostic performance of cardiovascular risk (CVR) calculators for the prediction of augmented CIMT and CP.

Methods: A cross-sectional study of subjects aged from 40-75 years that fulfilled the 2010 ACR/EULAR RA criteria. Exclusion criteria: previous CV disease and overlap syndromes. CVR was calculated by 6 scales: ACC/AHA 2013, Framingham-lipids (FRS-lipids), Framingham-Body Mass Index (FRS-BMI), Reynolds Risk Score (RRS), QRISK2 and SCORE. Carotid ultrasound (US) was performed by a board-certified radiologist and reviewed by other two. We defined an augmented CIMT as ≥ 0.9 mm and CP as CIMT ≥ 1.2 mm. We categorized subjects in 3 groups: 1)

normal carotid US, 2) augmented CIMT, and 3) CP. We used Kruskal-Wallis test to compare the groups.

Results: 130 patients were included, 124 (95.4%) were females with median age of 57 years old and disease duration of 9.7 years. Almost 50% of the subjects with CP were in the low-risk strata of every CVR calculator. The presence of CP was higher in subjects categorized as moderate/high risk using ACC/AHA 2013 CVR score ($p = 0.013$), moderate/high risk strata of QRISK2 ($p = 0.004$), and high-risk strata of ACC/AHA 2013 CVR score ($p = 0.02$). Of these 3, the highest sensitivity was obtained by the moderate risk stratum of QRISK2 with 51%; the highest specificity was obtained by the high-risk stratum of the ACC/AHA 2013 CVR score with 84%.

Table 1. Comparison groups

	CP presence N=39	CIMT ≥ 0.9 mm N= 29	Normal CIMT N= 62	p
Age years	59 (55-66)	60 (53-70)	51 (44-58)	0.00
BMI kg/m ²	27.3 (25.3-29.9)	27.8 (24.7-31.3)	29.9 (26.6-34.5)	0.022
ACC/AHA 2013	4.6 (2.3-10.2)	5.6 (2.2-11.8)	1.7 (0.7-3.5)	0.00
FRS-LIPIDS	7.1 (5.3-10.3)	9.25 (6.1-13.4)	3.8 (2.1-6.1)	0.00
FRS-BMI	10.4 (7.7-15.4)	12.2 (8.7-21.6)	4.8 (2.7-9.3)	0.00
SCORE	1 (0.5-2)	1 (0-2)	0 (0-1)	0.00
QRISK2	10.5 (5.2-16.7)	11.6 (5.6-17.1)	3.1 (1.9-6.3)	0.00
RRS	2 (1.3-4)	3 (1.5-7)	1 (1-2)	0.00

Variables reported as median, (q25 - q75)

Table 2. Diagnostic test performance

CVR score moderate and high risk	CP (n=39)	No CP (n=91)	p	HR (95% CI)	Sens (%)	Spec (%)	PPV (%)	NPV (%)
ACC/AHA 2013, n (%)	19 (48.7)	24 (26.4)	0.01	2.6 (1.2-5.7)	0.48	0.73	0.44	0.77
FRS-Lipids, n (%)	11 (28.2)	17 (18.7)	0.22	1.71 (0.7-4)	0.28	0.81	0.39	0.72
FRS-BMI, n (%)	20 (51.3)	31 (34.1)	0.06	2.03 (0.9-4.3)	0.51	0.65	0.39	0.75
QRISK2, n (%)	20 (51.3)	23 (25.3)	0.00	3.1 (1.4-6.8)	0.51	0.74	0.46	0.78
RRS, n (%)	2 (5.1)	2 (2.2)	0.37	2.4 (0.3-17.7)	0.05	0.97	0.50	0.70

Conclusion: Increased CIMT or CP might be unidentified if the risk is exclusively determined by CVR calculators. Patients with a CIMT ≥ 0.9 mm had an average score that placed them in the low risk strata, leading to an underestimation of their real CVR. Calculators can be helpful to estimate the CVR, however, they don't seem to be an adequate screening tool for an increased CIMT or CP. Meanwhile, to assess CVR precisely, a carotid US should be performed in all RA patients.

REFERENCES:

- [1] Peters M, et al. EULAR evidence-based recommendations for cardiovascular risk management in patients with rheumatoid arthritis and other forms of inflammatory arthritis. ARD 2010.
- [2] Meune C, et al. Cardiovascular mortality in patients with rheumatoid arthritis over 50 years: a systematic review and meta-analysis of cohort studies. Rheumatology 2009.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.7454

FRI0038

HERPES ZOSTER IN RHEUMATOID ARTHRITIS. PROSPECTIVE SINGLE CENTER STUDY OF 390 RA PATIENTS FOR 5YEARS

Lucia Domínguez¹, Vanesa Calvo-Rio¹, Paz Rodríguez-Cundin², Virginia Portilla¹, Nuria Vegas-Revenga¹, Francisco Manuel Antolin-Juarez², María Henar Rebollo Rodríguez², Alfonso Corrales¹, D. Prieto-Peña¹, Monica Calderón-Goercke¹, Miguel A. González-Gay¹, Ricardo Blanco¹. ¹Hospital Universitario Marqués de Valdecilla, Rheumatology, Santander, Spain; ²Hospital Universitario Marqués de Valdecilla, Preventive Medicine, Santander, Spain

Background: Immunosuppressed patients such as Rheumatoid Arthritis (RA) patients have a greater risk (1.5-2 times) of presenting herpes zoster (HZ). Both, the disease itself and the use of immunosuppressive

Case	Sex	Age (years)	Duration of RA (months)	RF +/-	APCA +/-	Erosive RA	HZ Localization	Previous Herpes Viral serology	HZ Treatment	Concomitant treatment for RA
1	♀	74	339	+	+	Yes	Left arm	IgG+ IgM NR	Famciclovir	Corticoids/MTX/TCZ
2	♀	86	142	+	-	Yes	-	IgG+ IgM -	None (topic)	ETN
3	♀	80	90	-	-	Yes	Gluteus (right)	IgG + IgM NR	Brivudina	HCC
4	♀	83	313	+	-	Yes	D9-D10 metamer	IgG + IgM -	Brivudina	Corticoids/MTX/RTX
5	♀	52	222	+	-	Yes	Generalized	IgG + IgM NR	Aciclovir	Corticoids
6	♀	70	171	+	-	No	Left arm	IgG+ IgM NR	Aciclovir	Corticoids/MTX/ADA
7	♀	68	1	+	-	Yes	Intercostal (left)	IgG + IgM -	Brivudina	Corticoids/HCC
8	♀	61	40	-	-	No	-	IgG + IgM NR	-	Corticoids/LFN/TCZ
9	♂	58	34	+	+	No	Intercostal (left)	IgG + IgM NR	Famciclovir	None
10	♀	62	67	+	+	Yes	Intercostal (left)	IgG + IgM NR	Aciclovir	LFN/ETN
11	♀	50	5	-	-	No	Intercostal (left)	IgG + IgM NR	Aciclovir	Corticoids/MTX/ADA
12	♀	66	2	+	+	No	Gluteus (left)	IgG + IgM -	Aciclovir	None

TABLE General features of 12 RA patients at HZ infection

drugs, are involve in this increased risk. Furthermore, in these patients is more frequent a disseminated presentation of zoster involving several dermatomes.

Objectives: In a series of RA patients our aim was analyzed HZ infections and to assess a) HZ prevalence and b) HZ general features

Methods: Prospective Single center study of 390 RA patients included in the vaccination program of the Preventive and Rheumatology department of our hospital between october 2011 and october 2016. The follow-up was made until December 2017. HZ vaccination is not included in our program.

RA was diagnosed according to the ACR/EULAR 2010 proposed criteria (Arthritis Rheum 2010; 62: 2569–2581)

The diagnose of HZ was made according to the clinical manifestations and was confirmed by a dermatologist. These manifestations were characteristic skin rash and blisters, paresthesia and local pain, in one (localized) or more dermatomes (generalized)

Results: We studied 390 patients (307♀/83♂), average age 61.28±12.9 years that were included in the vaccination program and followed-up.

HZ infection was observed in 12 of 390 (3.07%) in the follow-up (TABLE).

The 12 RA patients (11 women/1 man) with a mean±SD age of 67.5 ±11.67. More than half of patients, 7 (58.33%) were taking corticosteroids. 8 patients (66.66%) were receiving conventional disease modifying drugs (DMARDs) methotrexate (33.33%), leflunomide (16.66%) and hydroxychloroquine (16.66%). Besides corticosteroids and conventional DMARDs, 7 patients (58.33%) were in treatment with biologic drugs, tocilizumab (n=2), etanercept (n=2), adalimumab (n=2), and rituximab (n=1).

Conclusion: Herpes zoster is a relative frequent viral infection in RA patients non-vaccine for HZ. The female sex, older age, more aggressive RA and treatment with corticosteroids were more frequent. Probably in this group of patients HZ vaccination may be useful.

Abbreviations: MTX: methotrexate, LFN: leflunomide, HCC: hydroxychloroquine, TCZ: tocilizumab, ADA: adalimumab, ETN: etanercept

Disclosure of Interests: Lucia Domínguez: None declared, Vanesa Calvo-Río: None declared, Paz Rodríguez-Cundin: None declared, Virginia Portilla: None declared, Nuria Vegas-Revenga: None declared, Francisco Manuel Antolin-Juarez: None declared, Maria Henar Rebollo Rodríguez: None declared, Alfonso Corrales: None declared, D. Prieto-Peña: None declared, Monica Calderón-Goercke: None declared, Miguel A González-Gay Grant/research support from: Prof. MA Gonzalez-Gay received grants/research supports from Abbvie, MSD, Jansen and Roche., Speakers bureau: Consultation fees/participation in company sponsored speaker's bureau from Pfizer, Lilly, Sobi, Celgene, Novartis, Roche and Sanofi., Ricardo Blanco Grant/research support from: Abbvie, MSD, and Roche, Consultant for: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen, Speakers bureau: Abbvie, Pfizer, Roche, Bristol-Myers, Janssen

DOI: 10.1136/annrheumdis-2019-eular.2482

with a higher mortality and identification of safe and effective drugs is essential.

Objectives: Characterize lung involvement in a RA cohort; evaluate rituximab (RTX) effectiveness and safety in RA associated interstitial lung disease (ILD).

Methods: Retrospective analysis of the electronic records from RA cohort followed at University College Hospital. Lung involvement was based on high resolution computed tomography. Demographic data, smoking status, complementary exams at baseline/follow-up and therapies used were analysed. A sub-analysis of pts treated with rituximab (RTX) evaluated response at 12, 24 and 36 months. Declines of 15% in gas transfer from baseline and/or 10% in forced vital capacity were recognized.

Results: From 1129 RA pts, 87 (7.7%) had documented lung involvement. Mean age at last follow-up was 68.3±12 years, 74.7% were female and 85.1% Caucasian. Median disease duration was 14 (IQR 8-29) years. 54% of pts had erosive disease. Rheumatoid factor was positive in 88.1% and anti-citrullinated protein antibodies in 87.8%. 23 pts had positive antinuclear antibodies (25 missing data) and 4 anti-Ro, 2 anti-Scl70 and 2 anti-PL12. Secondary Sjögren's syndrome occurred in 6.9%, cutaneous rheumatoid nodules in 5.7% and cutaneous vasculitis in 1.1%. 11.5% and 43.7% were current and previous smokers, respectively. Mean interval between onset of articular and pulmonary symptoms was 12.3 years; 2 pts had lung disease as a prior manifestation. Types of lung involvement are shown in table 1.

TABLE 1 - Characterisation of different types of lung involvement

	Female	Age at lung disease diagnosis (years, mean±SD)	Interval between RA and lung disease diagnosis (years, median [IQR])	Smoking habits	Positive RF	Positive ACPA
NSIP (n=18)	11 (61.1%)	66.3±12.5	1 [0-15]	13 (76.5%; 1 missing)	16 (88.9%)	15 (88.2%; 1 missing)
UIP (n=18)	11 (61.1%)	66.8±13	1 [0-13.5]	9 (52.9%; 1 missing)	14 (87.5%; 2 missing)	16 (94.1%; 1 missing)
OP (n=4)	4 (100%)	57±7.1	10 [4.25-21]	1 (25%)	4 (100%)	4 (100%)
Unclassifiable ILD pattern (n=5)	4 (80%)	65.2±12.4	4 [2.5-8.5]	4 (80%)	5 (100%)	5 (100%)
Isolated bronchiectasis (n=31)	27 (87.1%)	63.7±9.3	17 [5.5-24.5]	14 (56%; 6 missing)	25 (83.3%; 1 missing)	23 (79.3%; 2 missing)
Obliterative bronchiolitis (n=2)	1 (50%)	74.5±3.5	30.5	1 (50%)	1 (50%)	2 (100%)
Panbronchiolitis (n=2)	2 (100%)	48.5±15.5	8.5	1 (50%)	2 (100%)	2 (100%)
Small airways disease without any specific pattern (n=1)	1 (100%)	57	4	0 (0%)	1 (100%)	1 (100%)
Rheumatoid nodules (n=4)*	4 (100%)	45.8±11.4	6.5 [2.5-15.75]	3 (75%)	4 (100%)	3 (75%)
Pleural effusion (n=3)*	2 (66.7%)	41±9.3	13	2 (66.7%)	3 (100%)	3 (100%)

*1 patient had concomitant pleural effusion and rheumatoid nodules

Legend: RF – Rheumatoid factor; ACPA – anti-citrullinated protein antibodies; NSIP – non-specific interstitial pneumonia; UIP – usual interstitial pneumonia; OP – organising pneumonia;

FRI0039 THE LUNG IN AN ENGLISH COHORT OF RHEUMATOID ARTHRITIS PATIENTS – AN OVERVIEW OF DIFFERENT TYPES OF INVOLVEMENT AND TREATMENT

Ana Catarina Duarte¹, Joanna Porter², Maria José Leandro³. ¹Hospital Garcia de Orta, Rheumatology Department, Almada, Portugal; ²University College London, Respiratory Medicine, London, United Kingdom; ³University College London, Rheumatology Department, London, United Kingdom

Background: Lung disease is described in 5–20% of patients (pts) with Rheumatoid arthritis (RA) and affects parenchyma, pleura, airways and vasculature; drug-induced pulmonary disease also occurs. It is associated

At last follow-up appointment 22 pts were still on methotrexate (MTX) and 27 had previously received it. MTX-acute pneumonitis occurred in 2 pts, both in the 1st year of treatment.

RTX was used in 26 pts (57.8%) with ILD (14 nonspecific interstitial pneumonia-NSIP, 8 usual interstitial pneumonia-UIP, 2 organising pneumonia-OP). The mean number of cycles was 4 (range 1-12). Two pts were concomitantly receiving mycophenolate mofetil and 1 azathioprine. RTX treatment outcomes are shown in table 2.