support from: AbbVie, Genentech, Lilly, Novartis, Pfizer, Consultant for: AbbVie, Amgen, BMS, Genentech, Lilly, Regeneron, Sanofi, Pfizer
DOI: 10.1136/annrheumdis-2019-eular.3082

FILGOTINIB, A SELECTIVE JANUS KINASE 1 (JAK1) INHIBITOR, MODULATES DISEASE-ASSOCIATED CYTOKINES IN PATIENTS WITH ACTIVE RHEUMATOID ARTHRITIS

Peter C. Taylor¹, Emon Elboudwarej², Li Wanying², Rachael E. Hawtin², Jinfeng Liu², Amer M. Mirza². ¹Botnar Research Centre, University of Oxford, Oxford, United Kingdom; ²Gilead Sciences, Inc., Foster City, United States of America

Background: Filgotinib (FIL), an oral selective JAK1 inhibitor, was safe and effective in FINCH2, a randomized, double-blind, placebo-controlled, phase 3 study in patients with active rheumatoid arthritis (RA) who had an inadequate response to biologic disease-modifying anti-rheumatic drugs (bDMARDs).¹

Objectives: A longitudinal study of cytokines from patients in FINCH2 was conducted to identify RA-associated biomarkers related to bone biology, immune cell migration, and inflammation that are altered by FIL therapy; and FIL-associated biomarkers that correlated with clinical response (DAS28 [CRP], swollen and tender joint counts, pain, and fatigue).

Methods: Plasma, serum, and urine samples from RA patients (n=449) who received FIL (100 mg, 200 mg) or placebo (PBO) once daily plus methotrexate were analyzed at baseline (BL) and week 12 for 42 disease-relevant cytokines using validated, commercially available single or multiplex assays. PBO-corrected on-treatment changes in cytokine levels from BL to week 12 were compared between treatment arms (Wilcoxon rank sum test). Spearman rank correlation was used to compare changes in cytokine level from BL to week 12 and clinical response. P-values <0.05 were considered significant.

Results: At week 12, 18 of 42 cytokines significantly decreased with FIL 100 mg treatment relative to PBO; FIL 200 mg decreased these cytokines to a similar or greater degree. An additional 6 cytokines were significantly decreased by FIL 200 mg. Conversely, 2 cytokines increased relative to PBO with FIL 100 mg, and 5 cytokines increased with FIL 200 mg (sIL-6R, IL-10, IL-2, leptin, and IL-17A). Biomarkers most significantly modulated by FIL 200 mg (p<0.0001) included markers related to bone biology (MMP1 [-22.8%], MMP3 [-24.7%], CTX1 [-27.4%], and NTX [-16.4%]), immune cell migration (VCAM1 [-20.0%], ICAM1 [-14.2%], CXCL13 [-45.0%], and CXCL10 [-32.3%]), and inflammation (TNFRI [-20.7%], CRP [-77.4%], SAA [-61.8%], IL-6 [-13.6%], and resistin [-20.2%]).

Hierarchical clustering of BL biomarker levels revealed distinct groups of cytokines that were strongly correlated with each other. Among them, SAA, IL-6, and CXCL10 were significantly positively correlated with each other (rho >0.6) and with RA disease activity (DAS28[CRP]) at BL (rho >0.3). Biomarkers, including CRP (IL-6, SAA), PainVAS (CRP, SAA), and SJC28 (CRP, IL-6, CXCL10), were also significantly correlated with individual components of DAS28[CRP]. Several biomarkers associated with RA disease activity at BL were decreased with FIL at week 12 relative to PBO (FIL 100 mg: CRP [-48.7%], SAA [-36.9%], and IL-6 [-2.6%]; and FIL 200 mg: CRP [-77.4%], SAA [-61.8%], IL-6 [-13.6%], and CXCL10 [-32.3%]), suggesting FIL impacts these disease activities at a molecular level.

Conclusion: Twelve weeks of FIL treatment significantly reduced 24 disease-relevant cytokines associated with bone biology, immune cell migration, and inflammation in patients with active RA. These effects were dose-dependent and suggest a shift toward a restored immune homeostasis. These findings are consistent with the clinical efficacy of FIL in FINCH2.

REFERENCES:

[1] Genovese MC, et al. ACR 2018. Abstract L06.

Disclosure of Interests: Peter C. Taylor Grant/research support from: Celgene, Galapagos, Eli Lilly, UCB, Consultant for: AbbVie, Galapagos, Gilead, Eli Lilly, Pfizer Inc, Emon Elboudwarej Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., Wanying Li Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc., MyoKardia, Rachael E. Hawtin Shareholder of: Gilead, Employee of: Gilead, Jinfeng Liu Shareholder of: Gilead Sciences, Inc., Roche, Employee of: Gilead Sciences, Inc., Amer M. Mirza Shareholder of: Gilead Sciences, Inc., Employee of: Gilead Sciences, Inc.

DOI: 10.1136/annrheumdis-2019-eular.2510

FRIO134

A PHASE 3 STUDY OF THE EFFICACY AND SAFETY OF PEFICITINIB (ASP015K) IN PATIENTS WITH RHEUMATOID ARTHRITIS WHO HAD AN INADEQUATE RESPONSE TO DMARDS

Yoshiya Tanaka¹, Tsutomu Takeuchi², Sakae Tanaka³, Atsushi Kawakami⁴, Manabu Iwasaki⁵, Yeong Wook Song⁶, Yi-Hsing Chen⁷, Mitsuhiko Rokuda⁸, Hiroyuki Izutsu⁸, Satoshi Ushijima⁸, Yuichiro Kaneko⁸, Teruaki Shiomi⁸, Emi Yamada⁸. ¹University of Occupational and Environmental Health, Kitakyushu, Japan; ²Keio University, Tokyo, Japan; ³University of Tokyo, Tokyo, Japan; ⁴Nagasaki University, Nagasaki, Japan; ⁵Yokohama City University, Yokohama, Japan; ⁶Seoul National University Hospital, Seoul, Korea, Rep. of (South Korea); ⁷Taichung Veterans General Hospital, Taichung City, Taiwan, Republic of China; ⁸Astellas Pharma, Inc., Tokyo, Japan

Background: Peficitinib (ASP015K), a novel oral JAK inhibitor, demonstrated efficacy as once-daily monotherapy in patients with moderate to severe rheumatoid arthritis (RA) in a phase 2b study (NCT01649999).

Objectives: To evaluate the efficacy and safety of peficitinib alone or in combination with disease-modifying antirheumatic drugs (DMARDs) in patients with RA who had an inadequate response to DMARDs.

Table 1 Primary and selected secondary efficacy endpoints at week 12/ET

Result at week 12/ET	PBO	Peficitinib 100 mg/day	Peficitinib 150 mg/day	Etanercept 50 mg/week (open-label arm)
ACR20 ^a , n/N (%)	31/101 (30.7)	60/104 (57.7)***	76/102 (74.5)***	167/200 (83.5)
ACR50 ^a , n/N (%)	9/101 (8.9)	32/104 (30.8)***	43/102 (42.2)***	105/200 (52.5)
ACR70 ^a , n/N (%)	1/101 (1.0)	14/104 (13.5) ^{b,c}	28/102 (27.5)***	61/200 (30.5)
Mean (SD) CRP change from baseline, mg/dL	0.022 (1.498)	-1.056 (1.908)***	-1.734 (1.906)***	-1.207 (2.694)
Mean (SD) ESR change from baseline, mm/h	-1.96 (17.82)	-12.96 (21.51)***	-23.92 (21.26)***	-20.92 (21.66)
DAS28-CRP <2.6 ^a , n/N (%)	5/100 (5.0)	25/102 (24.5)***	35/101 (34.7)***	91/200 (45.5)
Mean (SD) DAS28-CRP change from baseline	-0.64 (1.20)	-1.62 (1.41)***	-2.17 (1.14)***	-2.42 (1.11)
Mean (SD) change from baseline in patient's assessment of pain, 100 mm VAS	-6.64 (24.61)	-23.78 (30.75)***	-30.65 (27.91)***	-30.82 (27.68)
SDAI remission (SDAI score ≤3.3) ^a , n/N (%)	0/100 (0.0)	9/102 (8.8) ^b	9/101 (8.9) ^b	37/200 (18.5)

Last Observation Carried Forward imputation method was used. Statistical testing was performed for ASP015K 100 mg and 150 mg, compared with PBO. ***p<0.001 vs PBO according to Wald's Chi-square test for ACR20, ACR50, ACR70, DAS28-CRP <2.6, and SDAI remission, and analysis of covariance for CRP, ESR, DAS28-CRP and patient's assessment of pain. Closed testing procedure was used for multiplicity adjustment in the primary analysis.

^aBased on logistic regression model: Response (responder, non-responder) = treatment + the prior biologic-DMARD-IR (no, yes) + concomitant DMARD use (no, yes) + study region (Japan, Korea, and Taiwan).

^bNot estimable owing to the small number of responders.

^cIn a post-hoc analysis, based on the logistic regression model in which the only independent variable was treatment, the odds ratio between PBO and peficitinib 100 mg for ACR70 was estimated and p=0.009.

SD=standard deviation; SDAI=simplified disease activity index; VAS=visual analog scale.

Table 2 Treatment-emergent adverse events

	PBO (N=101)	Peficitinib 100 mg/day (N=104)	Peficitinib 150 mg/day (N=102)	Peficitinib 100 mg/day + 150 mg/day (N=206)	Etanercept 50 mg/week (open-label arm) (N=200)
Event, n (%), week 0-12					
Any AE	54 (53.5)	59 (56.7)	55 (53.9)	114 (55.3)	119 (59.5)
Drug-related AE ^a	29 (28.7)	33 (31.7)	38 (37.3)	71 (34.5)	75 (37.5)
Death	0	0	0	0	0
SAE	4 (4.0)	3 (2.9)	2 (2.0)	5 (2.4)	4 (2.0)
Drug-related SAE ^a	3 (3.0)	2 (1.9)	1 (1.0)	3 (1.5)	4 (2.0)
Grade ≥3 AE ^a	8 (7.9)	6 (5.8)	3 (2.9)	9 (4.4)	6 (3.0)
AE leading to permanent discontinuation of study drug or reference drug	4 (4.0)	6 (5.8)	3 (2.9)	9 (4.4)	5 (2.5)
Serious infection	0	0	1 (1.0)	1 (0.5)	1 (0.5)
Herpes zoster-related disease	0	1 (1.0)	0	1 (0.5)	2 (1.0)
Malignancy	0	0	0	0	0
Incidence rate per 100 patient-years (95% confidence interval), overall study period^b					
Serious infection ^c	0.0	1.1 (0.2, 8.1)	2.2 (0.5, 8.7)	2.0 (0.8, 4.9)	2.0 (0.8, 5.5)
Herpes zoster-related disease	0.0	5.8 (2.4, 13.8)	4.4 (1.7, 11.7)	5.8 (3.4, 9.8)	2.6 (1.1, 6.2)
Malignancy	0.0	2.3 (0.6, 9.1)	0.0	1.2 (0.4, 3.8)	0.5 (0.1, 3.6)

^aPossibly or probably related to study drug, as assessed by the investigator, or records where relationship is missing.

^bBased on NCI-CTCAE grade: grade 3-severe or medically significant, grade 4-life threatening, grade 5-death related to AE.

^cPatient-years covers from initial dose up to first incidence of AE for patients who had at least one event; otherwise, the duration of the patients are summed as from initial dose through follow-up. Incidence rate was calculated as (100 × number of patients who had at least one incidence) / total patient-years.

^dIncludes AEs that occurred after initial peficitinib dosing, and after switching from PBO to peficitinib 100 mg/day or 150 mg/day.

^eDefined as an AE belonging to the system organ class of "Infections and infestations" and regarded as serious.

AE=adverse event; SAE=serious adverse event.

Methods: This multicentre, randomised, double-blind, parallel-group, placebo (PBO)-controlled phase 3 study (NCT02308163) was conducted in Japan, Korea and Taiwan. All patients had RA diagnosed according to 1987 ACR or 2010 ACR/EULAR criteria. Patients with active RA (defined as ≥ 6 tender and painful joints and ≥ 6 swollen joints, using 68 and 66-joint assessment respectively, and CRP >0.50 mg/dL) and inadequate response to DMARDs (administered for ≥ 90 days) were randomised in a 1:1:1:2 ratio to 52 weeks' treatment with PBO, peficitinib 100 mg/day, peficitinib 150 mg/day or etanercept 50 mg/week (open-label reference arm). At week 12, patients initially assigned to PBO were switched (under blinded conditions) to either peficitinib 100 mg/day or peficitinib 150 mg/day until end of treatment. Concomitant stable dose of DMARDs was permitted. The primary efficacy variable was ACR20 response rate at week 12/early termination (ET).

Results: In total, 507 patients were randomised and treated: PBO ($n=101$), peficitinib 100 mg/day ($n=104$), peficitinib 150 mg/day ($n=102$) and etanercept ($n=200$). Regarding efficacy at week 12/ET, significant differences were observed with peficitinib 100 mg/150 mg vs PBO ($p<0.001$) in the proportion of patients achieving ACR20, ACR50, ACR70 (150 mg/day dose only) and DAS28-CRP <2.6 , and for change from baseline to week 12/ET in DAS28-CRP (Table 1). Week 0–12 safety results were similar between treatment groups and serious adverse events were more common with PBO than other study treatments (Table 2). For the overall study period, the incidence rate of serious infections per 100 patient-years was higher with peficitinib 100 mg/150 mg than PBO (Table 2). There were no deaths during the study.

Conclusion: In patients with RA who had an inadequate response to DMARDs, 100 mg/day and 150 mg/day peficitinib doses significantly reduced RA symptoms according to clinical and patient assessment scores. The proportion of patients achieving the primary efficacy variable (ACR20 at week 12/ET) was significantly greater for both peficitinib doses versus PBO. Peficitinib 100 mg/day and 150 mg/day showed acceptable safety and tolerability, with no new safety signals detected compared with other JAK inhibitors.

Acknowledgement: This study was sponsored by Astellas Pharma, Inc. Medical writing support was provided by Rhian Harper Owen of Cello Health MedErgy and funded by Astellas Pharma, Inc.

Disclosure of Interests: Yoshiya Tanaka Grant/research support from: Abbvie, Astellas, Bristol-Myers Squibb, Chugai, Daiichi-Sankyo, Eisai, Mitsubishi-Tanabe, MSD, Ono, Taisho-Toyama, Takeda, Speakers bureau: Abbvie, Asahi-kasei, Astellas, Bristol-Myers Squibb, Chugai, Daiichi-Sankyo, Eli Lilly, Eisai, Glaxo-SmithKline, Janssen, Mitsubishi-Tanabe, Novartis, Pfizer Japan Inc, Sanofi, Takeda, UCB, YL Biologics, Tsutomu Takeuchi Grant/research support from: Astellas Pharma Inc, Chugai Pharmaceutical Co., Ltd., Daiichi Sankyo Co., Ltd., Takeda Pharmaceutical Co., Ltd., AbbVie GK, Asahikasei Pharma Corp., Mitsubishi Tanabe Pharma Co., Pfizer Japan Inc., Eisai Co., Ltd., AYUMI Pharmaceutical Corporation, Nipponkayaku Co. Ltd., Novartis Pharma K.K., Grant/research support from: AbbVie, Asahi Kasei, Astellas, AstraZeneca, AYUMI, Bristol-Myers Squibb, Chugai, Daiichi Sankyo, Eisai, Eli Lilly Japan, Janssen, Mitsubishi Tanabe, Nippon Kayaku, Novartis, Pfizer Japan Inc, Taiho, Taisho Toyama, Takeda, Teijin, Grant/research support from: Astellas Pharma Inc., Bristol Myers Squibb, Chugai Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., Pfizer Japan Inc., Santen Pharmaceutical Co., Ltd., Takeda Pharmaceutical Co., Ltd., Teijin Pharma Ltd., AbbVie GK, Asahi Kasei Pharma Corp., Taisho Toyama Pharmaceutical Co., Ltd., SymBio Pharmaceuticals Ltd., Janssen Pharmaceutical K.K., Celltrion Inc., Nipponkayaku Co. Ltd., and UCB Japan, Consultant for: Astra Zeneca K.K., Eli Lilly Japan K.K., Novartis Pharma K.K., Mitsubishi Tanabe Pharma Co., Abbvie GK, Nipponkayaku Co.Ltd, Janssen Pharmaceutical K. K., Astellas Pharma Inc., Taiho Pharmaceutical Co. Ltd., Chugai Pharmaceutical Co. Ltd., Taisho Toyama Pharmaceutical Co. Ltd., GlaxoSmithKline K.K., UCB Japan Co. Ltd., Consultant for: AbbVie, Asahi Kasei, Astellas, AstraZeneca, AYUMI, Bristol-Myers Squibb, Chugai, Daiichi Sankyo, Eisai, Eli Lilly Japan, Janssen, Mitsubishi Tanabe, Nippon Kayaku, Novartis, Pfizer Japan Inc, Taiho, Taisho Toyama, Takeda, Teijin, Consultant for: Astra Zeneca K.K., Eli Lilly Japan K.K., Novartis Pharma K.K., Mitsubishi Tanabe Pharma Co., Asahi Kasei Medical K.K., AbbVie GK, Daiichi Sankyo Co., Ltd., Bristol Myers Squibb, and Nipponkayaku Co. Ltd., Speakers bureau: Astellas Pharma Inc., Bristol Myers Squibb, Chugai Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., Pfizer Japan Inc., Santen Pharmaceutical Co., Ltd., Takeda Pharmaceutical Co., Ltd., Teijin Pharma Ltd., AbbVie GK, Asahi Kasei Pharma Corp., Taisho Toyama Pharmaceutical Co., Ltd., SymBio Pharmaceuticals Ltd., Janssen Pharmaceutical K.K., Celltrion Inc., Nipponkayaku Co. Ltd., and UCB Japan, Speakers bureau: AbbVie, Asahi Kasei, Astellas, AstraZeneca, AYUMI, Bristol-Myers Squibb, Chugai, Daiichi Sankyo, Eisai, Eli Lilly Japan, Janssen, Mitsubishi Tanabe, Nippon Kayaku, Novartis, Pfizer Japan Inc, Taiho, Taisho Toyama, Takeda, Teijin, Speakers bureau: AbbVie GK., Bristol-Myers K.K., Chugai Pharmaceutical Co. Ltd., Mitsubishi Tanabe Pharma Co., Pfizer Japan Inc., Astellas Pharma Inc,

Daiichi Sankyo Co. Ltd., Eisai Co. Ltd., Sanofi K.K., Teijin Pharma Ltd., Takeda Pharmaceutical Co. Ltd., Novartis Pharma K.K., Sakae Tanaka Grant/research support from: KYOCERA Corporation and Asahi Kasei Corporation, Consultant for: Amgen Astellas BioPharma K.K., KYOCERA Corporation, Pfizer and Daiichi Sankyo Co., Ltd., Speakers bureau: Asahi Kasei Corporation, Astellas Pharma Inc, Ayumi Pharmaceutical Corporation, Eisai Co., Ltd., Ono Pharmaceutical Co., Ltd., Daiichi Sankyo Co., Ltd, Taisho Toyama Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Corporation, Chugai Pharmaceutical Co., Ltd., Teijin Pharma Ltd., Eli Lilly, Hisamitsu Pharmaceutical Co, Inc., Pfizer, Bristol-Myers., Atsushi Kawakami Grant/research support from: Astellas Pharma, Consultant for: Astellas Pharma, Speakers bureau: Astellas Pharma, Manabu Iwasaki: None declared, Yeong Wook Song: None declared, Yi-Hsing Chen Speakers bureau: Astellas Pharma, Mitsuhiro Rokuda Employee of: Astellas Pharma, Inc., Hiroyuki Izutsu Employee of: Astellas Pharma, Inc., Satoshi Ushijima Employee of: Astellas Pharma, Inc., Yuichiro Kaneko Employee of: Astellas Pharma, Inc., Teruaki Shiomi Employee of: Astellas Pharma, Inc., Emi Yamada Employee of: Astellas Pharma, Inc.

DOI: 10.1136/annrheumdis-2019-eular.1448

FRI0135

INTRAVENOUS PEGYLATED LIPOSOMAL PREDNISOLONE SODIUM PHOSPHATE MORE EFFECTIVE THAN INTRAMUSCULAR INJECTION OF METHYLPREDNISOLONE ACETATE IN TREATING RA PATIENTS WHO ARE EXPERIENCING A FLARE: A PHASE III, RANDOMIZED, DOUBLE-BLIND, DOUBLE DUMMY, ACTIVE CONTROLLED, MULTI-CENTER STUDY

Johannes W.J. Bijlsma^{1,2}, Bart Metselaar³, Leonie Middelink⁴, Cees Wortel⁵, Reinhard Bos⁶, Jacob M. van Laar¹, Harald Vonkeman⁷, Rene Westhovens⁸, Siu Long Yao⁹, Mudgal Kotheekar⁹, Atul Raut⁹. ¹University Medical Center Utrecht, Utrecht, Netherlands; ²University Medical Center Utrecht, Rheumatology and Clinical Immunology, Utrecht, Netherlands; ³Enceladus, Utrecht, Netherlands; ⁴Middelinc BV, Utrecht, Netherlands; ⁵Accelovance, Boston, United States of America; ⁶Medisch Centrum Leeuwarden, Leeuwarden, Netherlands; ⁷Medisch Spectrum Twente, Enschede, Netherlands; ⁸Universitair Ziekenhuis Leuven, Leuven, Belgium; ⁹SUN Pharma, Mumbai, India

Background: Intravenous pegylated liposomal prednisolone sodium phosphate is developed to deliver liposomal entrapped prednisolone in the bloodstream; the lipid vesicles allow selective accumulation of glucocorticoids in inflamed tissues by locally increased permeability of blood vessel walls, while limiting systemic exposure.

Objectives: To assess efficacy and safety of liposomes with prednisolone in patients with active rheumatoid arthritis (RA) who are experiencing a flare in comparison to a standard of care medication (methylprednisolone injections).

Methods: The study enrolled active RA patients (≥ 18 years) who were experiencing a flare of their disease, defined as a recent switch from a period with well documented remission or low disease activity to an active disease phase, as determined by a change in the Disease Activity Score in 28 Joints (DAS28 ≥ 3.2). Patients were randomized 1:1:1 into liposomal prednisolone 75 mg, liposomal prednisolone 150 mg and methylprednisolone 120 mg groups. Treatment was administered on Day 1 and Day 15. Patients treated with liposomal prednisone IV received IM placebo injections; patients treated with methylprednisolone IM received IV placebo infusions. Evaluations were performed weekly for the first 4 weeks, and every second week thereafter. The primary endpoint was EULAR response rate (good and moderate combined) at Day 8. The primary analysis involved two comparisons: 1) liposomal prednisone 150 mg vs. methylprednisolone 120 mg, 2) liposomal prednisone 75 mg vs. methylprednisolone 120 mg. Safety results (AEs, vital signs, physical examinations, laboratory, ECG) were also evaluated. A total of 172 patients were screened and 150 were randomized to liposomal prednisone 75 mg ($N=49$), liposomal prednisone 150 mg ($N=52$) and methylprednisolone 120 mg ($N=49$). 144 (96%) patients completed the study; mean study duration was 82.9 days; adverse events or intercurrent illness were primary reasons for discontinuation.

Results: EULAR response rate at Week 1 (Day 8) was 90.0% for liposomal prednisone 150 mg and 85.7% for liposomal prednisone 75 mg vs. 66.7% for methylprednisolone 120 mg treated patients (p -values of 0.007 and 0.018, resp.). Other secondary endpoints supported the primary endpoint results showing significant or clinically meaningful improvements in other EULAR response evaluations, as well as DAS28, VAS, ACR20/50/70, SF-36, HAQ and FACIT-F assessments. Similar numbers of patients reported at least one adverse event (AE) in each treatment group; 42 (86 %), 46 (89 %) and 39 (80 %), resp., for the liposomal prednisone 75 mg, liposomal prednisone 150 mg and methylprednisolone 120 mg groups. Most commonly reported AEs were nausea in the liposomal prednisone groups and headache in all 3 treatment arms. Approximately 8%