Methods: This was a post hoc analysis of 3-, 6- and 12-month data from the active treatment arms of Phase (P)3 study, ORAL Standard (NCT00853385; n=609) and P3b/4 study, ORAL Strategy (NCT02187055; n=1152) in patients with active RA refractory to methotrexate (MTX). Both studies included tofacitinib 5 mg BID + MTX and ADA (40 mg SC q2w) + MTX treatment arms. ORAL Standard also included a tofacitinib 10 mg BID + MTX treatment arm and ORAL Strategy, a tofacitinib 5 mg BID treatment arm. A paired joint pathology score (PJPS), a combination of bilateral tender/swollen joint counts ranging from 0 (neither side swollen/tender) to 4 (both sides swollen/tender) was calculated. Data for 33 (ORAL Standard) and 14 (ORAL Strategy) joint pairs were available. The percentage change from baseline (%Δ) in PJPS for each joint was presented in a heat map to visualise responses. To show tofacitinib-specific effects at each joint, difference in mean PJPS for each tofacitinib regimen at 3 months minus the respective ADA + MTX group was calculated.

Results: Across all treatment arms of both studies, baseline joint involvement ranged from 15.7% (5th distal interphalangeal joint of the hand) to 93.3% (wrist). All joints showed a treatment response, with %ΔPJPS ranging from -36 to -94% (Figure). The heat map showed lower response rates in the wrist and knee vs most other joints in both studies persisting to 12 months, regardless of treatment arm. Similarly, in ORAL Standard, the ankle (not assessed in ORAL Strategy) showed a lower response vs most other joints. Tofacitinib-specific responses vs ADA + MTX (not shown) were seen (mean differences in PJPS ranged from -0.4 to 0.2) with no clear pattern in the combination therapy groups. In the tofacitinib 5 mg BID arm of ORAL Strategy, a pattern was observed in the metacarpophalangeal (MCP) joints, with tofacitinib responsiveness vs ADA + MTX increasing progressively (mean difference in PJPS increased from 0.16 in MCP1 to -0.08 in MCP5).

Figure: Heat map showing the percentage change from baseline in PJPS for each joint

Conclusion: There is variation in the responsiveness of RA joints to tofacitinib ± MTX and ADA + MTX, with the wrist, knee and ankle responding less well vs other joints, irrespective of treatment mode of action. Concomitant MTX use in most treatment arms may mask joint-specific actions of tofacitinib and ADA. Observed responsiveness to tofacitinib monotherapy vs ADA + MTX increased from MCP1-5, which may be tentative evidence of differential sensitivity to JAK inhibition across embryologically imprinted functional genetic gradients. Further analyses of the monotherapy arms of randomised trials may yield additional evidence of joint-specific responsiveness, and provide a basis for personalised RA treatment.

REFERENCES:

- [1] Frank-Bertoncelj M, et al. Nat Commun 2017; 8: 14852.
- [2] Ai R, et al. Nat Commun 2016; 7: 11849.

Acknowledgement: Study sponsored by Pfizer Inc. Medical writing support was provided by Kirsten Woolcott of CMC Connect and funded by Pfizer Inc.

Disclosure of Interests: Adrian Ciurea Consultant for: AbbVie, Celgene, Janssen-Cilag, MSD, Eli Lilly, Novartis, Pfizer, UCB, Speakers bureau: AbbVie, Celgene, Janssen-Cilag, MSD, Eli Lilly, Novartis, Pfizer, UCB, Oliver Distler Grant/research support from: Prof. Distler received research funding from Actelion, Bayer, Boehringer Ingelheim and Mitsubishi Tanabe to investigate potential treatments of scleroderma and its complications, Consultant for: Prof. Distler has/had consultancy relationship within the last 3 years with Actelion, AnaMar, Bayer, Boehringer Ingelheim, Chemo-mAb, espeRare foundation, Genentech/Roche, GSK, Inventiva, Italfarmaco, iQvia, Lilly, medac, MedImmune, Mitsubishi Tanabe Pharma, Pharmacyclics, Novartis, Pfizer, Sanofi, Serodapharm and UCB in the area of potential treatments of scleroderma and its complications. In addition, he

had/had consultancy relationship within the last 3 years with A. Menarini, Amgen, Abbvie, GSK, Mepha, MSD, Pfizer and UCB in the field of arthritides and related disorders, Tim Killeen Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Ermeg Akybekova Employee of: E Akybekova is an employee of IQVIA, which is a paid contractor to Pfizer Inc, Kenneth Kwok Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Lisy Wang Shareholder of: Pfizer Inc, Employee of: Pfizer Inc, Caroline Ospelt: None declared, Mojca Frank-Bertoncelj Grant/research support from: AbbVie

DOI: 10.1136/annrheumdis-2019-eular.1196

FR10145

FACTORS ASSOCIATED WITH NEAR REMISSION DIFFER BETWEEN AUTOANTIBODY-POSITIVE AND -NEGATIVE PATIENTS WITH EARLY RHEUMATOID ARTHRITIS TREATED WITH CONVENTIONAL SYNTHETIC DISEASE-MODIFYING ANTIRHEUMATIC DRUGS

Ludovico De Stefano, Serena Bugatti, Francesca Benaglio, Garifallia Sakellariou, Antonio Manzo, Roberto Caporali, Carlomaurizio Montecucco. Division of Rheumatology, IRCCS Policlinico San Matteo Foundation, University of Pavia, Pavia, Italy

Background: Disease remission is recommended as the treatment target in patients with early rheumatoid arthritis (RA). According to the criterion, the proportion of patients classified as in remission considerably varies, with the Boolean-based definition being the most restrictive [1]. A considerable proportion of patients with established RA misses Boolean remission solely because of one of the 4 criteria (near remission) [2]. The frequency and the limiting variables to Boolean remission in real-life cohorts of patients with early RA are however unknown.

Objectives: To assess the frequency and the limiting factors for fulfilling Boolean remission in early RA patients treated with csDMARDs according to a treat-to-target strategy.

Methods: The study population consisted of 578 early RA patients (<12 months of symptoms) consecutively recruited at our Early Arthritis Clinic, treatment-naïve at inclusion and prospectively followed-up upon initiation of therapy with methotrexate (MTX) aimed at the achievement of low disease activity (28-joints disease activity score [DAS28] ≤3.2). After 6 months of treatment, patients were classified as in remission or near remission according to the Boolean criterion as follows: (i) remission – swollen joint count (SJC28), tender joint count (TJC28), C-reactive protein (CRP) and patient global assessment of disease activity (PGA) all ≤1; (ii) near remission – just 3 of the 4 core items. Patients characteristics between remission and near remission were compared, and reasons for missing Boolean remission were analysed in relation to disease variables.

Results: Complete follow-up data after 6 months of treatment were available for 89.3% of the patients. Disease remission was achieved in 25.6% of the cases according to the DAS28, 16.7% according to the simplified disease activity index (SDAI) and 11.6% according to the Boolean criterion. One-hundred and thirty-three patients (25.8%) failed to reach Boolean remission solely because of one of the 4 criteria. Of these, 45.9% were in DAS28 remission and 22.6% in SDAI remission. No differences in baseline demographic and clinical characteristics were found between patients in Boolean remission and patients in near remission. Use of prednisone co-medication and MTX starting dose were also comparable. Patients negative for both rheumatoid factor (RF) and anti-citrullinated protein autoantibodies (ACPA) were more frequently in near remission rather than in Boolean remission compared to autoantibody-positive patients (Fig. 1A). The missing criterion in near remission was SJC28 >1 in 49% of the cases, PGA >1 in 41%, and TJC28 >1 or CRP >1 mg/dl in the remaining 10%. Collectively, failure to achieve Boolean remission because of SJC28 >1 was more common in RF and/or ACPA positive patients. Rather, autoantibody-negative patients more frequently missed remission due to PGA (Fig. 1B).

Conclusion: In patients with early RA treated with csDMARDs, near remission is up to two times more frequent than Boolean remission. Collectively, autoantibody-negative patients more commonly miss complete remission solely because of one criterion compared to autoantibody-

positive patients. Furthermore, in these patients, failure to achieve remission is mostly related to high PGA rather than to the persistence of joint inflammation. Altogether, these findings implicate that autoantibody-negative RA patients failing to achieve remission would require adjunctive tailored interventions rather than reinforcement of DMARD therapy.

REFERENCES:

- [1] Mack ME, et al. *Arthritis Rheumatol* 2017;69:518-28.
- [2] Studenic P, et al. *Ann Rheum Dis* 2012;71:1702-5.

Disclosure of Interests: Ludovico De Stefano: None declared, Serena Bugatti Speakers bureau: Bristol-Myers Squibb, Celgene, Lilly, Novartis, Sanofi, Janssen, Francesca Benaglio: None declared, Garifallia Sakellariou: None declared, Antonio Manzo: None declared, Roberto Caporali Speakers bureau: AbbVie, Bristol-Myers Squibb, Celgene, Roche, Genzyme, Lilly, MSD, Pfizer, UCB, Carlomaurizio Montecucco Speakers bureau: AbbVie, Bristol-Myers Squibb, Celgene, Sanofi, Genzyme, Lilly, MSD, Pfizer, UCB

DOI: 10.1136/annrheumdis-2019-eular.5967

FRI0146 SURVIVAL OF JAK INHIBITORS: REAL-WORLD DATA FROM THE BIOBADASER III REGISTRY

Valentina Emperiale^{1,2}, Carlos Sánchez-Piedra¹, Eduardo Cuende², Paloma Vela-Casasempere³, María del Carmen Castro Villegas⁴, Sara Manrique Arijas⁵, Cristina Campos Fernández⁶, Javier del Pino⁷, Manuel Pombo⁸, Fernando Sánchez-Alonso¹, Juan Jesus Gomez-Reino⁸. ¹Spanish Society of Rheumatology, Research Unit, Madrid, Spain; ²HU Príncipe de Asturias, Alcalá de Henares, Spain; ³HGU de Alicante, Alicante, Spain; ⁴HU Reina Sofía, Córdoba, Spain; ⁵HR de Málaga, Málaga, Spain; ⁶HGU de Valencia, Valencia, Spain; ⁷H de Salamanca, Salamanca, Spain; ⁸HCU de Santiago, Santiago de Compostela, Spain

Background: JAK inhibitors (JAKi) have been recently authorized in the European Union. JAKi long-term performance in the clinical practice setting still needs to be assessed. Drug survival reflects a drug's effectiveness, safety, and tolerability; thus, it is an important measurement for real-world studies.

Objectives: To describe the pattern of use of JAKi in real-world dataset, and to analyze the drug survival as a group (tofacitinib+baricitinib).

Methods: Information was obtained from BIOBADASER III, an ongoing observational longitudinal multicenter cohort of patients with rheumatic diseases treated with biologic or targeted synthetic (b/ts) DMARDs in Spain. All subjects with current or previous JAKi use were included and followed until September 2018. Clinical characteristics of the patients were analyzed. Drug survival (overall, and for biologic-naïve and biologic-experienced patients) was determined using Kaplan-Meier analysis.

Table 1. Diagnoses of patients with JAKi treatment

Diagnosis (patients)	n (%)
Rheumatoid arthritis	138 (92.6)
Idiopathic juvenile arthritis*	3 (2.0)
Psoriatic arthritis	2 (1.3)
Undifferentiated spondyloarthritis*	2 (1.3)
Seronegative polyarthritis*	2 (1.3)
Ankylosing spondylitis*	1 (0.7)
Enteropathic arthritis*	1 (0.7)
Total	149 (100)

*off-label use.

Results: 149 patients, 75.2% women, were treated with JAKi, receiving a total of 152 cycles of treatment (50.7% tofacitinib, 49.3% baricitinib; 3 patients had both). The most frequent diagnosis was rheumatoid arthritis (RA, 92.6%), and there is a small number of off-label uses (6%), depicted at Table 1. Use on psoriatic arthritis is scarce (1.3%). Concomitant use of methotrexate (MTX) was registered in 68 patients (45.6%). Previous use of bDMARDs was high (n=124, 81.6%); drug survival rate for biologic-experienced patients was 81.7% and 78.7% at 6 and 12 months. None of the JAKi treatments in biologic-naïve patients (n=18, 18.4%) were discontinued during follow-up. Pooled survival rate of JAKi was 85.0% and 82.5% at 6 and 12 months (Figure 1). Discontinuation was seen in 19 treatments (12.5%); the reasons were inefficacy (n=15, 9.9%) or adverse events (n=4, 2.6%).

Figure 1. Overall survival of JAKi and for biologic-naïve and biologic-experienced patients

Conclusion: The current use of JAKi in Spain is mainly in RA and as 2nd line after bDMARDs. The use of JAKi in psoriatic arthritis is still scarce and a small group of patients are treated off-label. Less than half use combination therapy with MTX. Overall survival of JAKi is superior to 80% at 12 months. A longer follow-up is needed to continue analyzing the survival of JAKi in a real-world context.

Acknowledgement: We thank all researchers from the BIOBADASER III group.

Disclosure of Interests: Valentina Emperiale: None declared, Carlos Sánchez-Piedra: None declared, Eduardo Cuende: None declared, Paloma Vela-Casasempere Grant/research support from: UCB, Abbvie, Pfizer, Roche, Bristol-Myer-Squibb (another research, not BIOBADASER related), Consultant for: UCB, Lilly, Pfizer, Roche, Bristol-Myer-Squibb, Speakers bureau: Roche, UCB, MSD, Pfizer, GSK, BMS, Lilly, María del Carmen Castro Villegas Paid instructor for: MSD, Abbvie, Pfizer, Janssen, Lilly, Roche, Sara Manrique Arijas Speakers bureau: Abbvie, MSD, Janssen, Lilly, Roche, Pfizer, Novartis, Cristina Campos Fernández: None declared, Javier del Pino: None declared, Manuel Pombo: None declared, Fernando Sánchez-Alonso: None declared, Juan Jesus Gomez-Reino: None declared

DOI: 10.1136/annrheumdis-2019-eular.211

FRI0147 SAFETY AND EFFECTIVENESS OF UPADACITINIB OR ADALIMUMAB IN PATIENTS WITH RHEUMATOID ARTHRITIS: RESULTS AT 48 WEEKS FROM THE SELECT-COMPARE STUDY

Roy Fleischmann¹, Jeffrey Enejesa², In-Ho Song³, Eduardo Mysler⁴, Louis Bessette⁵, Charles Peterfy⁶, Patrick Durez⁷, Andrew Ostor⁸, Li Yihan³, Yijie Zhou³, Mark C. Genovese⁹. ¹Univ of Texas Southwestern Med Center, Metrolplex Clinical Research Center, Dallas, TX, United States of America; ²AbbVie, Inc., N Chicago, IL, United States of America; ³AbbVie, Inc., N Chicago, IL, United States of America; ⁴Organización Médica de Investigación, Buenos Aires, Argentina; ⁵Laval Univ, Quebec, Canada; ⁶Spire Sciences Inc, Boca Raton, FL, United States of America; ⁷Pôle de Recherche en Rhumatologie, Institut de Recherche Expérimentale et Clinique, UCL Saint-Luc, Brussels, Belgium; ⁸Cabrini Medical Center, Malvern, Australia; ⁹Stanford Univ School of Medicine, Palo Alto, CA, United States of America

Background: In the SELECT-COMPARE study in rheumatoid arthritis (RA) patients (pts) with inadequate response to methotrexate (MTX), upadacitinib (UPA), a JAK1- selective inhibitor, was superior to placebo (PBO) or adalimumab (ADA) for treatment of signs & symptoms and for inhibition of radiographic progression vs PBO up to 26 weeks (wks). All pts were on stable background MTX. ¹

Objectives: To report safety and efficacy of UPA plus stable background MTX up to 48 wks in this phase 3 study.

Methods: Pts were randomized to once-daily (QD) UPA 15mg, PBO, or ADA 40mg every other wk, with all patients continuing background MTX. The study was double-blind for 48 wks. Between Wks14-26, pts were rescued (from PBO to UPA, UPA to ADA, or ADA to UPA) if there was <20% improvement in tender/swollen joint count (Wks 14/18/22) or if Clinical Disease Activity Index (CDAI) was >10 (Wk26); all PBO pts who were not rescued were switched to UPA at Wk 26. Efficacy was analyzed by randomized group. Non-responder imputation (NRI) was used for binary endpoints for rescue prior to Wk26. Last observation carried