positive patients. Furthermore, in these patients, failure to achieve remission is mostly related to high PGA rather than to the persistence of joint inflammation. Altogether, these findings implicate that autoantibody-negative RA patients failing to achieve remission would require adjunctive tailored interventions rather than reinforcement of DMARD therapy.

REFERENCES:

- [1] Mack ME, et al. *Arthritis Rheumatol* 2017;69:518-28.
- [2] Studenic P, et al. *Ann Rheum Dis* 2012;71:1702-5.

Disclosure of Interests: Ludovico De Stefano: None declared, Serena Bugatti Speakers bureau: Bristol-Myers Squibb, Celgene, Lilly, Novartis, Sanofi, Janssen, Francesca Benaglio: None declared, Garifallia Sakellariou: None declared, Antonio Manzo: None declared, Roberto Caporali Speakers bureau: AbbVie, Bristol-Myers Squibb, Celgene, Roche, Genzyme, Lilly, MSD, Pfizer, UCB, Carlomaglio Montecucco Speakers bureau: AbbVie, Bristol-Myers Squibb, Celgene, Sanofi, Genzyme, Lilly, MSD, Pfizer, UCB

DOI: 10.1136/annrheumdis-2019-eular.5967

FRI0146 SURVIVAL OF JAK INHIBITORS: REAL-WORLD DATA FROM THE BIOBADASER III REGISTRY

Valentina Emperiale^{1,2}, Carlos Sánchez-Piedra¹, Eduardo Cuende², Paloma Vela-Casasempere³, María del Carmen Castro Villegas⁴, Sara Manrique Arijia⁵, Cristina Campos Fernández⁶, Javier del Pino⁷, Manuel Pombo⁸, Fernando Sánchez-Alonso¹, Juan Jesus Gomez-Reino⁸. ¹Spanish Society of Rheumatology, Research Unit, Madrid, Spain; ²HU Príncipe de Asturias, Alcalá de Henares, Spain; ³HGU de Alicante, Alicante, Spain; ⁴HU Reina Sofía, Córdoba, Spain; ⁵HR de Málaga, Málaga, Spain; ⁶HGU de Valencia, Valencia, Spain; ⁷H de Salamanca, Salamanca, Spain; ⁸HCU de Santiago, Santiago de Compostela, Spain

Background: JAK inhibitors (JAKi) have been recently authorized in the European Union. JAKi long-term performance in the clinical practice setting still needs to be assessed. Drug survival reflects a drug's effectiveness, safety, and tolerability; thus, it is an important measurement for real-world studies.

Objectives: To describe the pattern of use of JAKi in real-world dataset, and to analyze the drug survival as a group (tofacitinib+baricitinib).

Methods: Information was obtained from BIOBADASER III, an ongoing observational longitudinal multicenter cohort of patients with rheumatic diseases treated with biologic or targeted synthetic (b/ts) DMARDs in Spain. All subjects with current or previous JAKi use were included and followed until September 2018. Clinical characteristics of the patients were analyzed. Drug survival (overall, and for biologic-naïve and biologic-experienced patients) was determined using Kaplan-Meier analysis.

Table 1. Diagnoses of patients with JAKi treatment

Diagnosis (patients)	n (%)
Rheumatoid arthritis	138 (92.6)
Idiopathic juvenile arthritis*	3 (2.0)
Psoriatic arthritis	2 (1.3)
Undifferentiated spondyloarthritis*	2 (1.3)
Seronegative polyarthritis*	2 (1.3)
Ankylosing spondylitis*	1 (0.7)
Enteropathic arthritis*	1 (0.7)
Total	149 (100)

*off-label use.

Results: 149 patients, 75.2% women, were treated with JAKi, receiving a total of 152 cycles of treatment (50.7% tofacitinib, 49.3% baricitinib; 3 patients had both). The most frequent diagnosis was rheumatoid arthritis (RA, 92.6%), and there is a small number of off-label uses (6%), depicted at Table 1. Use on psoriatic arthritis is scarce (1.3%). Concomitant use of methotrexate (MTX) was registered in 68 patients (45.6%). Previous use of bDMARDs was high (n=124, 81.6%); drug survival rate for biologic-experienced patients was 81.7% and 78.7% at 6 and 12 months. None of the JAKi treatments in biologic-naïve patients (n=18, 18.4%) were discontinued during follow-up. Pooled survival rate of JAKi was 85.0% and 82.5% at 6 and 12 months (Figure 1). Discontinuation was seen in 19 treatments (12.5%); the reasons were inefficacy (n=15, 9.9%) or adverse events (n=4, 2.6%).

Figure 1. Overall survival of JAKi and for biologic-naïve and biologic-experienced patients

Conclusion: The current use of JAKi in Spain is mainly in RA and as 2nd line after bDMARDs. The use of JAKi in psoriatic arthritis is still scarce and a small group of patients are treated off-label. Less than half use combination therapy with MTX. Overall survival of JAKi is superior to 80% at 12 months. A longer follow-up is needed to continue analyzing the survival of JAKi in a real-world context.

Acknowledgement: We thank all researchers from the BIOBADASER III group.

Disclosure of Interests: Valentina Emperiale: None declared, Carlos Sánchez-Piedra: None declared, Eduardo Cuende: None declared, Paloma Vela-Casasempere Grant/research support from: UCB, Abbvie, Pfizer, Roche, Bristol-Myer-Squibb (another research, not BIOBADASER related), Consultant for: UCB, Lilly, Pfizer, Roche, Bristol-Myer-Squibb, Speakers bureau: Roche, UCB, MSD, Pfizer, GSK, BMS, Lilly, María del Carmen Castro Villegas Paid instructor for: MSD, Abbvie, Pfizer, Janssen, Lilly, Roche, Sara Manrique Arijia Speakers bureau: Abbvie, MSD, Janssen, Lilly, Roche, Pfizer, Novartis, Cristina Campos Fernández: None declared, Javier del Pino: None declared, Manuel Pombo: None declared, Fernando Sánchez-Alonso: None declared, Juan Jesus Gomez-Reino: None declared

DOI: 10.1136/annrheumdis-2019-eular.211

FRI0147 SAFETY AND EFFECTIVENESS OF UPADACITINIB OR ADALIMUMAB IN PATIENTS WITH RHEUMATOID ARTHRITIS: RESULTS AT 48 WEEKS FROM THE SELECT-COMPARE STUDY

Roy Fleischmann¹, Jeffrey Enejoza², In-Ho Song³, Eduardo Mysler⁴, Louis Bessette⁵, Charles Peterfy⁶, Patrick Durez⁷, Andrew Ostro⁸, Li Yihan³, Yijie Zhou³, Mark C. Genovese⁹. ¹Univ of Texas Southwestern Med Center, Metrolplex Clinical Research Center, Dallas, TX, United States of America; ²AbbVie, Inc., N Chicago, IL, United States of America; ³AbbVie, Inc., N Chicago, IL, United States of America; ⁴Organización Médica de Investigación, Buenos Aires, Argentina; ⁵Laval Univ, Quebec, Canada; ⁶Spire Sciences Inc, Boca Raton, FL, United States of America; ⁷Pôle de Recherche en Rhumatologie, Institut de Recherche Expérimentale et Clinique, UCL Saint-Luc, Brussels, Belgium; ⁸Cabrini Medical Center, Malvern, Australia; ⁹Stanford Univ School of Medicine, Palo Alto, CA, United States of America

Background: In the SELECT-COMPARE study in rheumatoid arthritis (RA) patients (pts) with inadequate response to methotrexate (MTX), upadacitinib (UPA), a JAK1- selective inhibitor, was superior to placebo (PBO) or adalimumab (ADA) for treatment of signs & symptoms and for inhibition of radiographic progression vs PBO up to 26 weeks (wks). All pts were on stable background MTX. ¹

Objectives: To report safety and efficacy of UPA plus stable background MTX up to 48 wks in this phase 3 study.

Methods: Pts were randomized to once-daily (QD) UPA 15mg, PBO, or ADA 40mg every other wk, with all patients continuing background MTX. The study was double-blind for 48 wks. Between Wks14-26, pts were rescued (from PBO to UPA, UPA to ADA, or ADA to UPA) if there was <20% improvement in tender/swollen joint count (Wks 14/18/22) or if Clinical Disease Activity Index (CDAI) was >10 (Wk26); all PBO pts who were not rescued were switched to UPA at Wk 26. Efficacy was analyzed by randomized group. Non-responder imputation (NRI) was used for binary endpoints for rescue prior to Wk26. Last observation carried