Conclusion: This study showed positive effects on disease activity and blood lipid levels of a diet combining foods with anti-inflammatory properties, but the results must be confirmed in larger studies.

REFERENCES:

- [1] Agca R, et al. EULAR recommendations for cardiovascular disease risk management in patients with rheumatoid arthritis and other forms of inflammatory joint disorders: 2015/2016 update. *Annals of the rheumatic diseases*. 2017;76(1):17-28.
- [2] Piepoli MF, et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *European heart journal*. 2016;37(29):2315-81.
- [3] Sködstam L, et al. An experimental study of a Mediterranean diet intervention for patients with rheumatoid arthritis. *Ann Rheum Dis*. 2003;62(3):208-14.
- [4] SLV. [Available from: <https://www.livsmedelverket.se/matvanor-halsamiljo/kostrad-och-matvanor/matvanor-undersokningar/riksmaten-2010-11-vuxna>.
- [5] Tedeschi SK, et al. Diet and Rheumatoid Arthritis Symptoms: Survey Results From a Rheumatoid Arthritis Registry. *Arthritis Care Res (Hoboken)*. 2017;69(12):1920-5.

Disclosure of Interests: Anna Vadell: None declared, Erik Hulander: None declared, Linnea Barebring: None declared, Inger Gjerstsson: None declared, Helen Lindqvist Employee of: Yes, for Janssen-Cilag AB in 2001-2003, Before I was a PhD-student and within a different field (dementia). My research field is within diet and Rheumatology., Anna Winkvist: None declared

DOI: 10.1136/annrheumdis-2019-eular.4522

FRI0169

DOES INITIATING TOCILIZUMAB LEAD TO BETTER DISEASE CONTROL COMPARED TO INITIATING MTX WITH LOW-MODERATE DOSE PREDNISONE IN EARLY RHEUMATOID ARTHRITIS; AN INDIRECT COMPARISON OF U-ACT-EARLY AND CAMERA-II TREAT-TO-TARGET TRIALS

Maxime Verhoeven¹, Janneke Tekstra¹, M Jacob. van Laar¹, Johannes WJ Bijlsma¹, Attila Pethoe-Schramm², Michelle Borm³, Floris Laféber¹, Paco Welsing¹. ¹University Medical Center Utrecht, Utrecht, Netherlands; ²Hoffmann-La Roche, Basel, Switzerland; ³Roche Nederland BV, Woerden, Netherlands

Background: Treatment with methotrexate (MTX), often with concomitant glucocorticoids, is the cornerstone of early rheumatoid arthritis (RA) therapy. However, it may be less effective compared to (expensive) biological disease modifying anti-rheumatic drugs, such as tocilizumab (TCZ). Hitherto, the effectiveness and safety of MTX in combination with glucocorticoids have never been compared to TCZ with or without MTX.

Objectives: To compare effectiveness and safety of initiating TCZ, or TCZ with MTX (TCZ+MTX) to initiation of MTX with 10mg prednisone (MTX+Pred) all in a step-up treat-to-target treatment strategy in early RA patients.

Methods: Individual patient data of the U-Act-Early ($n=317$) and CAMERA-II ($n=236$) trials were used. Both were 2-year, double-blind, randomised, placebo-controlled studies evaluating step-up tight-control, treat-to-target treatment strategies with the opportunity to taper, in case of sustained remission, TCZ and/or MTX.^{1,2} Using MTX ($n=108+119$) as the reference strategy, TCZ+MTX ($n=106$) and TCZ ($n=103$) were compared with MTX+Pred ($n=117$): primary outcome was the disease activity score (DAS28) over time. Secondary outcomes were remission, defined as DAS28<2.6, and the ConReW score (cumulative occurrence of remission and sustainment of remission). To assess the influence of acute phase reactants (APRs) on the results a disease activity outcome without APRs was also analysed (i.e. CDAI, modified due to lack of VAS physician in CAMERA-II). Multiple imputation was used for missing baseline data: HAQ, rheumatoid factor (RF) and smoking status. Multi-level models were used to account for clustering of patients within trials and for repeated measurements within patients over time. All models were corrected for baseline DAS28, HAQ, RF-status and smoking using fixed (and random) effects.

Results: Differences between U-Act-Early and CAMERA-II for RF seropositivity and DAS28 at baseline were observed; respectively 73% vs. 60%

($p<0.01$) and 5.2 vs. 5.7 ($p=0.01$). DAS28 was statistically significantly lower over time for TCZ+MTX compared to MTX+Pred (mean difference: -0.62 [95%CI -1.14 to -0.10]), but not for TCZ, Table 1. Remission occurred more often in TCZ+MTX and TCZ compared to MTX+Pred: relative risk 1.11 [95%CI 1.02 to 1.22] and 1.09 [1.00 to 1.20], respectively. ConReW scores were in line but not statistically significant different, Table 1. When using modified-CDAI, TCZ strategies did not show better control of disease activity over time than MTX+Pred (mean difference in log transformed values: TCZ+MTX vs. MTX+Pred: 0.10 [95%CI -0.16 to 0.36]; TCZ vs. MTX+Pred: 0.24 [95%CI -0.04 to 0.52]), Table 1. No differences in safety outcomes could be established, Table 2.

	TCZ+MTX vs MTX+Pred		TCZ vs MTX+Pred	
Efficacy	Mean difference (95% CI)	P-value	Mean difference (95% CI)	P-value
DAS28	-0.62 (-1.14 to -0.10)	0.01	-0.44 (-1.00 to 0.12)	0.08
ConReW	3.02 (-1.12 to 6.17)	0.14	4.73 (-0.41 to 9.86)	0.08
Modified CDAI ^a	0.10 (-0.16 to 0.36)	0.48	0.24 (-0.04 to 0.52)	0.09
	Relative Risk (95% CI)		Relative Risk (95% CI)	
Remission	1.11 (1.02 to 1.22)	0.01	1.09 (1.00 to 1.20)	0.05

^alog transformed values. Multiplying the above regression coefficients by 100 leads to a difference on a log scale of percentage.

modified percentage scale

Table 2 Differences in safety outcomes over 24 months between TCZ (U-Act-Early) and MTX+Pred (CAMERA-II) strategy arms

	TCZ+MTX vs MTX+Pred		TCZ vs MTX+Pred	
Safety*	Relative Risk (95% CI)	P-value	Relative Risk (95% CI)	P-value
Elevated ALT	1.62 (0.66 to 3.97)	0.30	0.92 (0.33 to 2.53)	0.86
Elevated AST	1.48 (0.35 to 6.17)	0.60	0.55 (0.11 to 2.66)	0.48
Infections	0.99 (0.11 to 9.03)	1.00	1.05 (0.11 to 9.58)	0.96
Drop out due to AE	1.18 (0.38 to 3.83)	0.78	1.30 (0.43 to 3.80)	0.60

* occurrence of 1 time

Conclusion: In early RA patients, TCZ-based strategies resulted in better DAS28 over time compared to MTX+Pred, as well as higher percentage of remission, part of these effects may be due to a specific effect of TCZ on APRs.

REFERENCES:

- [1] Bijlsma JWJ, et al. *Lancet*. 2016;388:343-55.
- [2] Bakker MF, et al. *Ann Intern Med*. 2012;156:329-339.

Disclosure of Interests: Maxime Verhoeven: None declared, Janneke Tekstra: None declared, Jacob M. van Laar Grant/research support from: Genentech, Consultant for: F. Hoffmann-La Roche, Johannes WJ Bijlsma Grant/research support from: The department of the author who included patients (JWJB) in the U-Act-Early trial received reimbursements from Roche Nederland BV. JWJB reported grants and fees from Roche, AbbVie, Bristol-Myers Squibb, Merck Sharp & Dohme, Pfizer, and UCB University Medical Center Utrecht, Utrecht University, Consultant for: SUN Pharma, Speakers bureau: Lilly, Roche, Attila Pethoe-Schramm Shareholder of: F. Hoffmann-La Roche, Employee of: F. Hoffmann-La Roche, Michelle Borm Employee of: An employee of Roche Nederland BV, Floris Laféber Shareholder of: ArthroSave, Grant/research support from: FOREUM; Dutch Arthritis Society, Paco Welsing: None declared

DOI: 10.1136/annrheumdis-2019-eular.3276

FRI0170

PHYSICAL ACTIVITY LEVEL IN RHEUMATOID ARTHRITIS: A SYSTEMATIC REVIEW

Charlotte Schouler¹, Jean Francis Maillefer², Jean Marie Casillas³, Daniel Wendling¹, Frank Verhoeven⁴. ¹CH Montceau, Rheumatology, Montceau Les Mines, France; ²CHU Dijon, Rheumatology, Dijon, France; ³CHU Dijon, Physical rehabilitation, Dijon, France; ⁴CHRU Besançon, Rheumatology, Besançon, France

Background: Rheumatoid arthritis (RA) is one of the most common inflammatory rheumatism characterized by an increased cardiovascular risk. Regarding the last EULAR recommendations, physical activity is an important part of the management of RA. The evaluation of Physical activity level is needed to know RA patients practices. However, to our knowledge, there is no consensual measure tool of the physical activity level for patients suffering RA.

Objectives: The aim of this study is to evaluate the different methods of measurement of physical activity levels in RA.

Methods: This is a systematic review of literature realized on the pubmed and Cochrane databases and meeting the PRISMA recommendations. We used the following key words: « physical activity », « physical activity level » AND « rheumatoid arthritis ». We included only article written in English language and with RA patients older than 18 years old.

Results: We identified 190 studies with the key words, 51 were selected on title and 23 articles have been identified as eligible. Finally, 19 studies, 13567 RA patients were evaluated on their level of physical activity. There were 73,4% female with a mean age of 56,1 years. In 10 studies, the BMI was available with results between 25 and 30 kg/m². Two methods for measuring physical activity levels have been identified.