REFERENCES:

- [1] Yamaguchi Y, Noriuchi N, Kaburaki J, Kobayashi K, Matsuura E, Kuwana M. Excessive exposure to anionic surfaces maintains autoantibody response to B2-glycoprotein I in patients with antiphospholipid syndrome. *Blood* 2007;110:4312-18.
- [2] Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, Derksen RH, DE Groot PG, Koike T, Meroni PL, Reber G, Shoenfeld Y, Tincani A, Vlachoyiannopoulos PG, Krilis SA. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome. *J Thromb Haemost* 2006; 4:295-306.
- [3] Alijotas-Reig J, Vilardell-Tarrés M. Is obstetric antiphospholipid syndrome a primary nonthrombotic, proinflammatory, complement-mediated disorder related to antiphospholipid antibodies? *Obstet Gynecol Surv* 2010; 65:39-45.
- [4] Berman J, Girardi G, Salmon JE. TNF-alpha is a critical effector and a target for therapy in antiphospholipid antibody-induced pregnancy loss. *J Immunol* 2005;174:1222-1226.
- [5] Meroni PL, Borghi MO, Raschi E, Tedesco F. Pathogenesis of antiphospholipid syndrome: understanding the antibodies. *Nat Rev Rheumatol* 2011;7:330-39.
- [6] Brogin Morel J, Cirino Ruocco AM, Vernini JM, Rudge MV, Calderon IM. Interleukin 10 and tumor necrosis factor-alpha in pregnancy: aspects of interest in clinical obstetrics. *ISRN Obstet Gynecol*. 2012;2012:230742.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.1279

FRI0192

A SYSTEMATIC LITERATURE REVIEW TO INFORM THE 2019 UPDATE OF THE EULAR RECOMMENDATIONS FOR THE TREATMENT OF SYSTEMIC LUPUS ERYTHEMATOSUS

Antonis Fanouriakis¹, Myrto Kostopoulou², Alessia Alunno³, Martin Aringer⁴, Ingeborg Bajema⁵, John N. Boletis⁶, Ricard Cervera⁷, Andrea Doria⁸, Caroline Gordon⁹, Marcello Govoni¹⁰, Frederic Houssiau¹¹, David Jayne¹², Marios Kouloumas¹³, Annegret Kuhn¹⁴, Janni Lisander Larsen¹⁵, Kirsten Lerstrom¹⁶, Gabriela Moroni¹⁷, Marta Mosca¹⁸, Matthias Schneider¹⁹, Josef S. Smolen²⁰, Elisabet Svenungsson²¹, Vladimir Tesar²², Angela Tincani²³, Anne Trolldborg²⁴, Ronald van Vollenhoven²⁵, Jörg Wenzel²⁶, George Bertias²⁷, Dimitrios Boumpas¹. ¹"Attikon" University Hospital, Athens, Greece; ²G. Gennimatas" General Hospital, Athens, Greece; ³University of Perugia, Perugia, Italy; ⁴University Medical Center and Faculty of Medicine Carl Gustav Carus at the TU Dresden, Dresden, Germany; ⁵Leiden University Medical Center, Leiden, Netherlands; ⁶"Laikon" Hospital, National and Kapodistrian University of Athens, Medical School, Athens, Greece; ⁷Hospital Clinic, Barcelona, Spain; ⁸University of Padova, Padova, Italy; ⁹Institute of Inflammation and Ageing, University of Birmingham, Birmingham, United Kingdom; ¹⁰University of Ferrara, Azienda Ospedaliero-Universitaria Sant'Anna Ferrara, Ferrara, Italy; ¹¹Université catholique de Louvain, Brussels, Belgium; ¹²University of Cambridge, Cambridge, United Kingdom; ¹³Cyprus League Against Rheumatism, Aglantzia, Cyprus; ¹⁴University Hospital Muenster, Muenster, Germany; ¹⁵Rheumatology and Spine Diseases Centre, Rigshospitalet, Copenhagen, Denmark; ¹⁶Lupus Europe, Farum, Denmark; ¹⁷Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ¹⁸University of Pisa, Pisa, Italy; ¹⁹UKD, Heinrich-Heine University, Düsseldorf, Düsseldorf, Germany; ²⁰Medial University of Vienna, Vienna, Austria; ²¹Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden; ²²1st Faculty of Medicine and General University Hospital, Charles University, Prague, Czech Republic; ²³University of Brescia, Brescia, Italy; ²⁴Aarhus University Hospital, Aarhus, Denmark; ²⁵Amsterdam University Medical Centers, Amsterdam, Netherlands; ²⁶University Hospital Bonn, Bonn, Germany; ²⁷University Hospital of Heraklion, Heraklion, Greece

Background: Culminating evidence over the past decade regarding management of systemic lupus erythematosus (SLE) called for an update of the 2008 EULAR recommendations for the treatment of the disease.

Objectives: Systematic review of the literature (SLR) to inform the 2019 EULAR recommendations for the management of SLE.

Methods: SLR of Pubmed from 01/2007 to 12/2017 for questions (selected through Delphi exercise) regarding: i) efficacy/safety of different drugs used in SLE, ii) treatment of specific manifestations, iii) monitoring and treatment goals and iv) comorbidities and adjunct therapy. Evidence was categorised based on design and validity of available studies [Oxford Centre for Evidence-Based Medicine 2011 Levels of Evidence (LoE)] and strength of statements was graded [Grading of recommendation (GoR), assessment, development and evaluations, GRADE].

Results: Main topics supported by strong evidence base included: Association of hydroxychloroquine (HCQ) use with favourable outcomes (LoE 1b,

GoR A), belimumab for extrarenal disease (LoE 1a, GoR A), efficacy of antimalarials in skin disease (LoE 1a, GoR A), mycophenolate mofetil (MMF) for induction and maintenance therapy of lupus nephritis (LN) and cyclophosphamide (CYC) in severe LN (LoE 1b, GoR A). Weak evidence supported the value of repeat kidney biopsy in refractory LN (LoE 4, GoR C), all second-line agents for skin disease (LoE 4, GoR C) and efficacy of most first and second-line treatments for thrombocytopenia (LoE 4, GoR C). Moderate LoE was found for all other questions (Table).

Conclusion: A SLR for the treatment of SLE found the highest LoE for benefits of HCQ, efficacy of belimumab for extrarenal disease and MMF and IV-CYC in LN.

Acknowledgement: European League Against Rheumatism

Disclosure of Interests: Antonis Fanouriakis Paid instructor for: Amgen, GSK, Speakers bureau: Abbvie, Enorasis, Genesis Pharma, Myrto Kostopoulou: None declared, Alessia Alunno: None declared, Martin Aringer Grant/research support from: Roche, Consultant for: AstraZeneca and Eli Lilly, Ingeborg Bajema Consultant for: GSK, John N. Boletis: None declared, Ricard Cervera: None declared, Andrea Doria: None declared, Caroline Gordon Grant/research support from: Sandwell and West Birmingham Hospitals NHS Trust have received funding from UCB to support research work done by my research group that was unrelated to any pharmaceutical product or clinical trial., Consultant for: I have provided consultancy advice and taken part in scientific advisory boards on the design and analysis of clinical trials and the management of lupus for GSK, EMD Serono and UCB. I have taken part in adjudication and safety monitoring committees for BMS., Speakers bureau: I have been paid by UCB for speaking at meetings., Marcello Govoni Paid instructor for: Pfizer, Roche, Speakers bureau: Pfizer, Abbvie, MSD, Roche, Eli-Lilly, Celgene, Sanofi, Janssen, Frederic Houssiau: None declared, David Jayne Grant/research support from: David Jayne has received research grants from Chemocentryx, GSK, Roche/Genentech and Sanofi-Genzyme. He has received consultancy fees from Astra-Zeneca, Boehringer-Ingelheim, Chemocentryx, Chugai, GSK, Infla-RX, Insmed and Takeda, Marios Kouloumas: None declared, Annegret Kuhn Grant/research support from: Biogen, Galderma, GlaxoSmithKline, LeoPharma, Speakers bureau: La Roche Posay, Janni Lisander Larsen: None declared, Kirsten Lerstrom: None declared, Gabriela Moroni: None declared, Marta Mosca Paid instructor for: GlaxoSmithKline, Lilly, UCB, Matthias Schneider Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study., Speakers bureau: Chugai, Josef S. Smolen Grant/research support from: AbbVie, Eli Lilly, Janssen, MSD, Pfizer Inc, Roche, Consultant for: AbbVie, Amgen, AstraZeneca, Astro, Celgene, Celtrion, Eli Lilly, GlaxoSmithKline, ILTOO, Janssen, Medimmune, MSD, Novartis-Sandoz, Pfizer Inc, Roche, Samsung, Sanofi, UCB, Speakers bureau: AbbVie, Amgen, AstraZeneca, Astro, Celgene, Celtrion, Eli Lilly, GlaxoSmithKline, ILTOO, Janssen, Medimmune, MSD, Novartis-Sandoz, Pfizer Inc, Roche, Samsung, Sanofi, UCB, Elisabet Svenungsson: None declared, Vladimir Tesar: None declared, Angela Tincani Consultant for: UCB, Pfizer, Abbvie, BMS, Sanofi, Roche, GSK, AlphaSigma, Lilly, Janssen, Celgene, Novartis, Anne Trolldborg: None declared, Ronald van Vollenhoven Grant/research support from: AbbVie, Amgen, Bristol-Myers Squibb, GlaxoSmithKline, Pfizer, Roche, and UCB, Consultant for: AbbVie, AstraZeneca, Biotest, Bristol-Myers Squibb, Celgene, Crescendo, GlaxoSmithKline, Janssen, Lilly, Merck, Novartis, Pfizer, Roche, UCB, and Vertex., Speakers bureau: AbbVie, AstraZeneca, Biotest, Bristol-Myers Squibb, Celgene, Crescendo, GlaxoSmithKline, Janssen, Lilly, Merck, Novartis, Pfizer, Roche, UCB, and Vertex., Jörg Wenzel Grant/research support from: GSK, Incyte, Consultant for: Biogen, Leo, Paid instructor for: Novartis, George Bertias: None declared, Dimitrios Boumpas: None declared

DOI: 10.1136/annrheumdis-2019-eular.6348

Table. Level of evidence for different items regarding treatment of systemic lupus erythematosus

Evidence Base	Topics in SLE treatment
A (strong)	<ul style="list-style-type: none"> Association of HCQ use with favourable outcomes Chronic exposure to GC and risk of organ damage Gonadal toxicity of CYC Efficacy and safety of belimumab in extrarenal SLE Efficacy of antimalarials in skin disease MMF for induction and maintenance therapy of LN and cyclophosphamide for severe LN Immunosuppressive therapy for inflammatory neuropsychiatric manifestations Association of aPL with adverse outcomes
B (moderate)	<ul style="list-style-type: none"> HCQ cumulative exposure and retinal toxicity Efficacy of MTX and MMF in extrarenal SLE Comparable efficacy and less toxicity of lower-dose GC regimens Efficacy of RTX in renal and extrarenal SLE Efficacy of multitarget therapies for induction treatment in LN Use of CNIs in refractory LN Primary prophylaxis with low-dose aspirin in aPL(+) SLE patients
C/D (weak)	<ul style="list-style-type: none"> Efficacy of AZA, CsA, and CYC in extrarenal SLE Efficacy of second-line treatment in skin lupus Efficacy of most first and second-line treatments for thrombocytopenia Efficacy of vaccinations in SLE patients Efficacy of primary cardiovascular protection with statins and low-dose aspirin in SLE

FRI0193

2019 UPDATE OF THE EULAR RECOMMENDATIONS FOR THE MANAGEMENT OF SYSTEMIC LUPUS ERYTHEMATOSUS

Antonis Fanouriakis¹, Myrto Kostopoulou², Alessia Alunno³, Martin Aringer⁴, Ingeborg Bajema⁵, John N. Boletis⁶, Ricard Cervera⁷, Andrea Doria⁸, Caroline Gordon⁹, Marcello Govoni¹⁰, Frederic Houssiau¹¹, David Jayne¹², Marios Kouloumas¹³, Annegret Kuhn¹⁴, Janni Lisander Larsen¹⁵, Kirsten Lerstrom¹⁶, Gabriela Moroni¹⁷, Marta Mosca¹⁸, Matthias Schneider¹⁹, Josef S. Smolen²⁰, Elisabet Svenungsson²¹, Vladimir Tesar²², Angela Tincani²³, Anne Trolldborg²⁴, Ronald van Vollenhoven²⁵, Jörg Wenzel²⁶, George Bertias²⁷, Dimitrios Boumpas¹. ¹"Attikon" University Hospital, Athens, Greece; ²G. Gennimatas" General Hospital, Athens, Greece; ³University of Perugia, Perugia, Italy; ⁴University Medical Center and Faculty of Medicine Carl Gustav Carus at the TU Dresden, Dresden, Germany; ⁵Leiden University Medical Center, Leiden, Netherlands; ⁶"Laikon" Hospital, National and Kapodistrian University of Athens, Medical School, Athens, Greece; ⁷Hospital Clinic, Department of Autoimmune Diseases, Barcelona, Spain; ⁸University of Padova, Padova, Italy; ⁹Institute of Inflammation and Ageing, University of Birmingham, Birmingham, United Kingdom; ¹⁰University of Ferrara, Azienda Ospedaliero-Universitaria Sant'Anna Ferrara, Ferrara, Italy; ¹¹Université catholique de Louvain, Louvain, Belgium; ¹²Cambridge, Department of Medicine, Cambridge, United Kingdom; ¹³Cyprus League Against Rheumatism, Aglantzia, Cyprus; ¹⁴University Hospital Muenster, Muenster, Germany; ¹⁵Rheumatology and Spine Diseases Centre, Rigshospitalet, Copenhagen, Denmark; ¹⁶Lupus Europe, Farum, Denmark; ¹⁷Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; ¹⁸University of Pisa, Pisa, Italy; ¹⁹UKD, Heinrich-Heine University, Düsseldorf, Germany; ²⁰Medical University of Vienna, Vienna, Austria; ²¹Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden; ²²Charles University, Prague, Czech Republic; ²³University of Brescia, Brescia, Italy; ²⁴Aarhus University Hospital, Aarhus, Denmark; ²⁵Amsterdam University Medical Centers, Amsterdam, Netherlands; ²⁶University Hospital Bonn, Bonn, Germany; ²⁷University Hospital of Heraklion, Heraklion, Greece

Background: Recommendations for the treatment of systemic lupus erythematosus (SLE) were published by EULAR in 2008. Advances in treatment strategies and goals called for an update of these recommendations, capitalizing on strengths and experience from previous projects.

Objectives: To update the EULAR recommendations for the management of SLE.

Methods: Systematic literature review (01/2007-12/2017) followed by Delphi method to form questions, elicit expert opinions and reach consensus.

Results: Treatment in SLE aims at remission or low disease activity and prevention of flares. Hydroxychloroquine is recommended in all lupus patients, at a dose not exceeding 5mg/kg real body weight. During chronic maintenance treatment, glucocorticoids should be minimized to less than 7.5 mg/day (prednisone equivalent) and, when possible, withdrawn. Prompt initiation of immunomodulatory agents (methotrexate, azathioprine, mycophenolate) can expedite tapering/discontinuation of glucocorticoids. In persistently active or flaring extrarenal disease, add-on belimumab should be considered; rituximab may be considered in organ-threatening, refractory disease. Specific recommendations are also provided for the management of cutaneous, neuropsychiatric, haematological

and renal disease (including the use of calcineurin inhibitors). SLE patients should be assessed for their antiphospholipid antibody status, infectious and cardiovascular diseases risk profile, and preventative strategies be tailored accordingly.

Conclusion: Updated EULAR recommendations provide physicians and patients with updated consensus guidance on the management of SLE, combining evidence-base and expert-opinion.

Figure

Acknowledgement: European League against Rheumatism **Disclosure of Interests:** Antonis Fanouriakis Paid instructor for: Amgen, GSK, Speakers bureau: Abbvie, Enorasis, Genesis Pharma, Myrto Kostopoulou: None declared, Alessia Alunno: None declared, Martin Aringer Grant/research support from: Roche, Consultant for: AstraZeneca and Eli Lilly, Ingeborg Bajema Consultant for: GSK, John N. Boletis: None declared, Ricard Cervera: None declared, Andrea Doria: None declared, Caroline Gordon Grant/research support from: Sandwell and West Birmingham Hospitals NHS Trust have received funding from UCB to support research work done by my research group that was unrelated to any pharmaceutical product or clinical trial., Consultant for: I have provided consultancy advice and taken part in scientific advisory boards on the design and analysis of clinical trials and the management of lupus for GSK, EMD Serono and UCB. I have taken part in adjudication and safety monitoring committees for BMS., Speakers bureau: I have been paid by UCB for speaking at meetings., Marcello Govoni Paid instructor for: Pfizer, Roche, Speakers bureau: Pfizer, Abbvie, MSD, Roche, Eli-Lilly, Celgene, Sanofi, Janssen, Frederic Houssiau: None declared, David Jayne Grant/research support from: David Jayne has received research grants from Chemocentryx, GSK, Roche/Genentech and Sanofi-Genzyme. He has received consultancy fees from Astra-Zeneca, Boehringer-Ingelheim, Chemocentryx, Chugai, GSK, Infla-RX, Insmed and Takeda, Marios Kouloumas: None declared, Annegret Kuhn Grant/research support from: Biogen, Galderma, GlaxoSmithKline, LeoPharma, Speakers bureau: La Roche Posay, Janni Lisander Larsen: None declared, Kirsten Lerstrom: None declared, Gabriela Moroni: None declared, Marta Mosca Paid instructor for: GlaxoSmithKline, Lilly, UCB, Matthias Schneider Grant/research support from: GlaxoSmithKline and UCB Pharma for performing the LuLa-study., Speakers bureau: Chugai, Josef S. Smolen Grant/research support from: AbbVie, Eli Lilly, Janssen, MSD, Pfizer Inc, Roche, Consultant for: AbbVie, Amgen, AstraZeneca, Astro, Celgene, Celtrion, Eli Lilly, GlaxoSmithKline, ILTOO, Janssen, Medimmune, MSD, Novartis-Sandoz, Pfizer Inc, Roche, Samsung, Sanofi, UCB, Speakers bureau: AbbVie, Amgen, AstraZeneca, Astro, Celgene, Celtrion, Eli Lilly, GlaxoSmithKline, ILTOO, Janssen, Medimmune, MSD, Novartis-Sandoz, Pfizer Inc, Roche, Samsung, Sanofi, UCB, Elisabet Svenungsson: None declared, Vladimir Tesar: None declared, Angela Tincani Consultant for: UCB, Pfizer, Abbvie, BMS, Sanofi, Roche, GSK, AlphaSigma, Lilly, Janssen, Celgene, Novartis, Anne Trolldborg: None declared, Ronald van Vollenhoven Grant/research support from: AbbVie, Amgen, Bristol-Myers Squibb, GlaxoSmithKline, Pfizer, Roche, and UCB, Consultant for: AbbVie, AstraZeneca, Biotest, Bristol-Myers Squibb, Celgene, Crescendo, GlaxoSmithKline, Janssen, Lilly, Merck, Novartis, Pfizer, Roche, UCB, and Vertex., Jörg Wenzel Grant/research support from: GSK, Incyte, Consultant for: Biogen, Leo, Paid instructor for: Novartis, George Bertias: None declared, Dimitrios Boumpas: None declared

DOI: 10.1136/annrheumdis-2019-eular.3117