with iRPF from four tertiary medical units in Greece and b) to evaluate factors potentially associated with disease relapse.

Methods: Medical records of patients diagnosed with RPF from 2000-2018 in four rheumatology units (Laiko, Euroclinic, Sismanoglion and Sotiria Hospitals) were retrospectively evaluated. Sixty-seven patients with iRPF were included in the study.

Results: The median age at diagnosis was 56 years (IQR:52.0–60.0), with median disease duration 8.0 years (IQR:3.0–11.0), 58% were smokers and 73% males. Patients more often presented with constitutional symptoms (57%), low back pain (63%), raised inflammatory markers (78%), anemia (43%) and compromised renal function (15%). Commonest imaging findings were periaortic-periliac mass (46%), periaortic mass (33%), periaortitis (25%) and hydronephrosis (36%) with envelopment of one (31%) or both ureters (18%). Tissue biopsy was requested in all patients, but was performed in only 10, with 3 having marked numbers of IgG4-positive plasma cells. Serum IgG4 was measured in 36/67 and 36% had elevated levels at diagnosis (median 224 mg/dL, IQR:174–328). Clinical/laboratory/radiological presentation did not differ between patients with elevated and normal serum IgG4 levels. Steroids were first-line treatment in 93% of patients. Other immunosuppressives used as steroid-sparing agents were azathioprine (70%), cyclophosphamide (19%), mycophenolate-mofetil (18%), D-penicillamine (12%) and methotrexate (8%). Relapse occurred in 19% of patients at a median of 36 months (IQR:18–66) after diagnosis with 69% of them being under therapy. Relapse did not correlate to initial imaging findings or to any treatment modality, yet patients with increased serum IgG4 tended to have higher relapse rate (26% vs 11%, $p=0.071$).

Conclusion: Diagnosis of iRPF was mostly based on imaging studies in our cohort. Steroids were used as first-line treatment. Relapse occurred in one-fifth of patients independently of initial clinical/radiographic presentation or treatment modality used. RPF patients with initially elevated serum IgG4 levels tended to have a higher relapse rate.

REFERENCES:

- [1] Vaglio A, Salvarani C, Buzio C. Retroperitoneal fibrosis. *Lancet* 2006; 367:241–51.
- [2] Brandt AS, Kamper L, Kukuk S, Haage P, Roth S. Associated findings and complications of retroperitoneal fibrosis in 204 patients: results of a urological registry. *J Urol*. 2011; 185:526–31.
- [3] Khosroshahi A, Carruthers MN, Stone JH, Shinagare S, Sainani N, Hasserjian RP, Deshpande V. Rethinking Ormond's disease: "idiopathic" retroperitoneal fibrosis in the era of IgG4-related disease. *Medicine (Baltimore)*. 2013; 92:82–91.

Disclosure of Interests: Aliki Venetsanopoulou: None declared, Evangelia Zampeli Speakers bureau: Roshe, Astrazeneca, Sophia Christaki: None declared, Ourania Argyropoulou: None declared, Kyriaki A. Boki: None declared, Menelaos N. Manoussakis: None declared, Fotini N. Skopouli: None declared, Athanasios Tzioufas Grant/research support from: ABBVIE, PFIZER, AMGEN, NOVARTIS, GSK, Haralampos M. Moutsopoulos: None declared

DOI: 10.1136/annrheumdis-2019-eular.4231

FRI0298

ABNORMALITY OF PERCENTAGES AND ABSOLUTE NUMBERS OF CD4+ T SUBSETS IN PATIENTS WITH ANCA-ASSOCIATED VASCULITIS AND ITS CORRELATIONS WITH CLINICAL INDICATORS

Yanlin Wang, Xiang-Cong Zhao, Jing Luo. *The Second Hospital of Shanxi Medical University, Taiyuan, China*

Background: ANCA-associated vasculitis (AAV) is a heterogeneous autoimmune disease with unknown etiology [1-2]. During the last decade, a panel of CD4+ T subsets have been identified. However, the exact role and quantitative status of these subsets in AAV patients remains unclear. **Objectives:** We therefore investigated these T cell subsets in AAV patients.

Methods: AAV patients ($n = 54$) and healthy controls (HCs) ($n = 19$) were enrolled. Of them, ten patients initially presenting with active disease were assessed again after remission was achieved. In addition, 38 patients were renal vasculitis. Proportions and absolute numbers of peripheral CD4+ T cell subsets and expression of multiplex cytokines were determined by flow cytometry (FCM). Correlations of clinical indicators with the CD4+ T cell subsets were systematically analyzed.

Results: Percentages of naive T cells (TN) ($p<0.001$), terminally differentiated effector (TEMRA) T cells ($p=0.027$) and activated Treg cells (aTreg) in AAV patients were decreased, but those of effector memory T-cell

subpopulation (TEM) ($p<0.001$), regulatory T cell (Treg) cells and Fox-P3lowCD45RA- T cells were increased. Similar results were observed when we compared absolute numbers of the above corresponding cells in AAV patients and HCs, except TEM. Furthermore, the percentage of aTreg ($p=0.043$) was decreased while that of Th17 cells ($p=0.027$) was increased in renal vasculitis patients. A significant correlation was observed between the ratio of Th17 to Treg subset and creatinine or BUN, as well as the ratio of Th17/aTreg was significantly increased in active and renal vasculitis patient. In addition, we found that cytokine IL-2 and IL-4 exhibited a downward while IL-6, IL-10, TNF, IFN- γ and IL-17A trend upward in AAV patients.

Conclusion: There were abnormally quantitative changes in CD4+ T subsets and cytokines in AAV patients, especially the decrease in the relative and absolute number of aTreg (activated Tregs), which indicates an imbalance of pro- and anti-inflammatory T cells. These T subsets might be associated with the ANCA-related autoimmune response and can be used as diagnosis markers for disease activity and as targets for potentially powerful therapy of AAV.

REFERENCES:

- [1] Hutton H L, Holdsworth S R, Kitching A R. ANCA-Associated Vasculitis: Pathogenesis, Models, and Preclinical Testing [J]. *Seminars in Nephrology*, 2017, 37(5):418.
- [2] Kerstein A, Silke Schüller, Otávio Cabral Marques, et al. Environmental factor and inflammation-driven alteration of the total peripheral T-cell compartment in granulomatosis with polyangiitis [J]. *Journal of Autoimmunity*, 2016, 78:79.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4667

FRI0299

THE PREVALENCE OF NON-VASCULAR PULMONARY MANIFESTATIONS IN PATIENTS WITH TAKAYASU' ARTERITIS: A RETROSPECTIVE MULTI-CENTERED COHORT STUDY

Ayten Yazici¹, Ayse Cefle¹, Sema Kaymaz Tahra², Nilüfer Alpay Kanitez³, Mete Kara⁴, Önay Gerçik⁵, Handan Yarkan Tuğsal⁶, Fatos Onen⁶, Servet Akar⁵, Kenan Aksu⁴, Gökhan Keser⁴, Fatma Alibaz-Oner², Haner Direskeneli². ¹Kocaeli University School of Medicine, Rheumatology, Kocaeli, Turkey; ²Marmara University School of Medicine, Rheumatology, Istanbul, Turkey; ³Koc University School of Medicine, Rheumatology, Istanbul, Turkey; ⁴Ege University School of Medicine, Rheumatology, Izmir, Turkey; ⁵Katip Celebi University School of Medicine, Rheumatology, Izmir, Turkey; ⁶Dokuz Eylul University School of Medicine, Rheumatology, Izmir, Turkey

Background: Takayasu' arteritis (TAK) is a rare vasculitis characterized by inflammation and obliteration of intermediate to large-size arteries. Even though more than 50% of patients with TAK have pulmonary artery involvement, non-vascular involvement and symptoms are uncommon.

Objectives: We aimed to investigate the frequency of non-vascular pulmonary involvement in TAK.

Methods: We assembled a retrospective cohort of patients with TAK from six different centers in Turkey. The demographics, clinical characteristics, treatment and outcomes of patients were abstracted from medical records, and the computed tomography findings were evaluated for pulmonary manifestations.

Results: As of January 2019, 197 TAK patients were recruited (mean age: 42.7±13.9 years [min-max: 17-75]) to the cohort, and 88.3% of them were female. Twenty-four patients had cough and/or dyspnea and four had hemoptysis as pulmonary symptoms. In CT assessment, parenchymal infiltrations were present in four (2%), pleural effusion in five (2.5%), nodule/cavity in one (0.5%), and pulmonary hemorrhage in one patient (0.5%). The patient who had pulmonary hemorrhage had also pleural effusion at the same time. In the whole cohort, 11.2% of patients ($n=22$) had pulmonary hypertension (PAH), three of them had cough and/or dyspnea and four of them had hemoptysis as a pulmonary symptom. Among patients with PAH, any pulmonary involvement in CT was more frequent compared to the rest of the patients (22.7% vs 5.1%, $p<0.0001$) (Table 1).

Conclusion: In this first assessment of Turkish TAK cohort, we observed non-vascular pulmonary involvement in about 5% of our patients and half of them were pleural effusions. The second most common manifestation was parenchymal infiltration with a frequency of 2%. Although rare, non-vascular pulmonary manifestations should also be investigated in TAK patients, especially in patients with pulmonary hypertension.

REFERENCES:

- [1] Nakajima N. Takayasu Arteritis: Consideration of pulmonary involvement. *Ann Vasc Dis* 2008;1(1):7-10.
- [2] Elsasser S, Soler M, Bollinger CT, et al. Takayasu disease with predominant pulmonary involvement. *Respiration* 2000;67(2):213-5.

Table 1. The frequencies of non-vascular pulmonary manifestations in patients with Takayasu' arteritis

n (%)	All Patients n=197	Patients with PAH n=22	p
Symptoms	24 (12.2)	3 (13.6)	0.049
Cough/Dyspnea	4 (2)	3 (13.6)	0.000
Hemoptysis			
Pulmonary involvement in CT	10 (5.1)	5 (22.7)	0.000
	4 (2)	1 (4.5)	0.000
Pulmonary infiltrates	1 (0.5)	1 (4.5)	0.000
Nodules/cavities	1 (0.5)	1 (4.5)	0.000
Pulmonary hemorrhage	5 (2.5)	2 (9.1)	0.000
Pleural effusion			

(PAH: pulmonary arterial hypertension, CT: computed tomography)

Disclosure of Interests: Ayten Yazici: None declared, Ayse Cefle: None declared, Sema Kaymaz Tahra: None declared, Nilüfer Alpay Kanitez: None declared, Mete Kara: None declared, Önay Gerçik: None declared, Handan Yarkan Tugsal: None declared, Fatos Onen: None declared, Servet Akar Grant/research support from: MSD, Abbvie, Roche, UCB, Novartis, Pfizer, Amgen, Consultant for: MSD, Abbvie, Roche, UCB, Novartis, Pfizer, Amgen, Speakers bureau: Pfizer, Kenan Aksu: None declared, Gökhan Keser: None declared, Fatma Alibaz-Oner: None declared, Haner Direskeneli: None declared

DOI: 10.1136/annrheumdis-2019-eular.3685

Scleroderma, myositis and related syndromes

FRI0300 IMPACT OF INTERSTITIAL LUNG DISEASE IN SYSTEMIC SCLEROSIS IN A COMPLETE, NATIONWIDE COHORT

Anna-Maria Hoffmann-Vold¹, Håvard Fretheim¹, Anna-Kristine H. Halse², Marit Seip³, Helle Bitter⁴, Marianne Wallenius⁵, Anne Salberg⁶, Torhild Garen¹, May Brit Lund¹, Trond M. Aaløkken¹, Øyvind Midtvedt¹, Øyvind Molberg¹. ¹Oslo University Hospital, Oslo, Norway; ²Haukeland University Hospital, Bergen, Norway; ³University Hospital of Northern Norway, Tromsø, Norway; ⁴Hospital of Southern Norway, Kristiansand, Norway; ⁵St. Olav's University Hospital, Trondheim, Norway; ⁶Lillehammer Hospital, Lillehammer, Norway

Background: Interstitial lung disease (ILD) represents a clinical challenge in systemic sclerosis (SSc) and associates with high mortality. The presence of severe lung fibrosis is a strong predictor for early mortality. There is substantial progress in SSc-ILD research, but precise, population-based data on cumulative incidence, range of severity and predictive value of clinical risk factors are lacking. Such data are vital for clinical decision making, and highly warranted as background information for appropriate development of screening and management strategies for SSc-ILD.

Objectives: To assess cumulative incidence of ILD, range of ILD severity and mortality risk predicted by baseline pulmonary function tests (PFT) and ILD extent by CT in a complete, nationwide SSc cohort.

Methods: The Norwegian SSc cohort study (Nor-SSc) includes all the 630 incident and 185 prevalent SSc patients from 2000-2012 meeting SSc classification criteria. A baseline PFT was recorded in 703 (86%) patients, and 650 (80%) had high resolution computed tomography (HRCT) images available for analyses. Extent of fibrosis was scored on 10 sections from every HRCT and expressed as percentage of total lung volumes. For the survival and mortality analyses, all Nor-SSc patients diagnosed from 2000-2012 (the 630 incident cases) were included and compared with 15 age- and gender matched controls per patient drawn from the national population registry. Vital status was available for all patients and controls at study end (January 2018). Descriptive statistics and standardized mortality rates (SMR) were estimated.

Results: Of the 815 patients in the total Nor-SSc cohort, 682 (84%) were female and 629 (77%) had limited cutaneous SSc. Mean age at SSc diagnosis was 53 yrs, with mean time from SSc onset to diagnosis of 3.8 yrs. We observed ILD on HRCT in 324/650 patients (50%), and the majority of these had <5% lung fibrosis (Figure 1A). Mean FVC at baseline was 94% of expected value, and nearly half of the patients (42%) had an FVC>100% (Figure 1B). Proportionate distribution of FVC

values in patients with no lung fibrosis, <10% lung fibrosis of total lung volume and >10% lung fibrosis is shown in Figure 1C. During the mean 8.6 yrs observation period of this study, 148 of the 630 incident SSc patients died, corresponding to an overall SMR of 2.4. Separate analyses of the 650 patients with baseline HRCT data showed that the SMR correlated with presence and extent of lung fibrosis, from SMR 2.2 in patients with no fibrosis to SMR 8 in patients with >25% lung fibrosis (Figure 1D). Correspondingly, we found that the SMR changed across patient groups stratified by baseline FVC%, with increased mortality evident already in the FVC 90-100% group (Figure 1E).

Conclusion: The results from this population based SSc cohort study provide new, unbiased data regarding the impact of ILD. Our results indicate a dose-response relationship between lung fibrosis extent and SMR; and between FVC% and SMR. Importantly, this relationship was evident even in groups with limited lung fibrosis and groups with normal range FVC%, strongly suggesting that all SSc patients should be screened with PFT and HRCT at baseline, to diagnose ILD early and tailor further management.

Disclosure of Interests: Anna-Maria Hoffmann-Vold Grant/research support from: Received research funding or other remuneration from Boehringer Ingelheim, GSK, and Actelion, Consultant for: Received consulting fees or other remuneration from Boehringer Ingelheim, GSK, and Actelion, Speakers bureau: Actelion and Boehringer Ingelheim, Håvard Fretheim Consultant for: Received consulting fees or other remuneration from GSK, and Actelion, Anna-Kristine H. Halse: None declared, Marit Seip: None declared, Helle Bitter: None declared, Marianne Wallenius: None declared, Anne Salberg: None declared, Torhild Garen: None declared, May Brit Lund: None declared, Trond M. Aaløkken: None declared, Øyvind Midtvedt: None declared, Øyvind Molberg: None declared

DOI: 10.1136/annrheumdis-2019-eular.6743

FRI0301 GASTROINTESTINAL ADVERSE EVENTS IN PATIENTS WITH SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD) TREATED WITH NINTEDANIB: DATA FROM THE SENSICIS TRIAL

Toby Maher¹, Kristin Highland², Martina Gahlemann³, Arata Azuma⁴, Aryeh Fischer⁵, Maureen Mayes⁶, Ganesh Raghu⁷, Wiebke Sauter⁸, Mannaig Girard⁹, Margarida Alves¹⁰, Emmanuelle Clerisme-Beatty¹⁰, Veronika Kohlbrenner¹¹, Masataka Kuwana¹², Oliver Distler¹³, SENSICIS trial investigators. ¹National Heart and Lung Institute, Imperial College London, United Kingdom, and National Institute for Health Research Clinical Research Facility, Royal Brompton Hospital, London, United Kingdom; ²Cleveland Clinic, Cleveland, Ohio, United States of America; ³Boehringer Ingelheim (Schweiz) GmbH, Basel, Switzerland; ⁴Department of Pulmonary Medicine and Oncology, Graduate School of Medicine, Nippon Medical School, Tokyo, Japan; ⁵University of Colorado School of Medicine, Denver, Colorado, United States of America; ⁶Division of Rheumatology and Clinical Immunogenetics, University of Texas McGovern Medical School, Houston, Texas, United States of America; ⁷University of Washington, Seattle, United States of America; ⁸Boehringer Ingelheim Pharma GmbH and Co. KG, Biberach an der Riss, Germany; ⁹Boehringer Ingelheim France S.A.S., Reims, France; ¹⁰Boehringer Ingelheim International GmbH, Ingelheim am Rhein, Germany; ¹¹Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, Connecticut, United States of America; ¹²Department of Allergy and Rheumatology, Nippon Medical School Graduate School of Medicine, Tokyo, Japan; ¹³Department of Rheumatology, University Hospital Zurich, Zurich, Switzerland

Background: In patients with idiopathic pulmonary fibrosis (IPF), nintedanib has a manageable adverse event (AE) profile characterised predominantly by gastrointestinal (GI) events. The efficacy and safety of