Conclusion: a significant improvement in endothelial function and microvascular involvement was achieved after three months of HCQ treatment. It is a novel finding in SSc which can represent a new therapeutic possibility for the prevention of microvascular complications of the disease

REFERENCES:

- [1] Fernández-Codina A, Walker KM, Pope JE; Scleroderma Algorithm Group. Treatment Algorithms for Systemic Sclerosis According to Experts. *Arthritis Rheumatol.* 2018;70:1820-1828. Doi: 10.1002/art.40560.
- [2] Virdis A, Tani C, Duranti E, Vagnani S, Carli L, Kühl AA, et al. Early treatment with hydroxychloroquine prevents the development of endothelial dysfunction in a murine model of systemic lupus erythematosus. *Arthritis Res Ther.* 2015;17:277.
- [3] Andracco R, Irace R, Zaccara E, Vettori S, Maglione W, Riccardi A, et al. The cumulative number of micro-haemorrhages and micro-thromboses in nailfold videocapillaroscopy is a good indicator of disease activity in systemic sclerosis: a validation study of the NEMO score. *Arthritis Res Ther.* 2017;19:133.

Disclosure of Interests: Fabio Basta: None declared, Rosaria Irace: None declared, Alessia Borgia: None declared, Valentina Messiniti: None declared, Antonella Riccardi: None declared, Gabriele Valentini Grant/research support from: MSD, Pfizer, Consultant for: MSD, Pfizer, biogen, Speakers bureau: MSD, amgen, biogen, lilly, sanofi, pfizer, Antonella Afeltra Grant/research support from: MSD, PFIZER, ABBVIE, ROCHE, UCB, Speakers bureau: MSD, PFIZER, BMS, ROCHE, SANOFI
DOI: 10.1136/annrheumdis-2019-eular.850

FRI0313 DIFFERENTIAL PERFORMANCE OF NAILFOLD VIDEO CAPILLAROSCOPIC PARAMETERS IN THE DIAGNOSIS AND PROGNOSIS OF SYSTEMIC SCLEROSIS

Vasiliki-Kalliopi Bournia¹, Konstantinos Kottas¹, Stylianos Panopoulos¹, George Konstantonis¹, Alexios Iliopoulos², Athanasios Tzioufas³, Petros Sfrikakis¹, Panayiotis Vlachoyiannopoulos³. ¹National and Kapodistrian University of Athens, Medical School, First Department of Propaedeutic and Internal Medicine, Athens, Greece; ²Veterans Administration Hospital (NIMTS), Department of Rheumatology, Athens, Greece; ³National and Kapodistrian University of Athens, Medical School, Pathophysiology, Athens, Greece

Background: The role of Nailfold Video Capillaroscopy (NVC) in the identification of patients with Raynaud phenomenon (RP) at risk to develop systemic sclerosis (SSc) is well established. However, it is not clear if certain capillaroscopic indices perform better than others at predicting SSc development and which NVC parameters have a prognostic value in established SSc.

Objectives: To comparatively assess the performance of different NVC parameters in predicting development of SSc, very early diagnosis of SSc (VEDOSS), or mixed connective tissue disease (MCTD) in patients with RP. Also, to longitudinally examine the consistency of clinical correlations of NVC parameters in SSc patients at two different time points and evaluate the prognostic capacity of NVC in SSc.

Methods: Consecutive RP patients referred to our department for NVC (138 with SSc, 12 with VEDOSS, 6 with MCTD, 36 with primary RP, and 50 with non-SSc secondary RP) were evaluated at baseline, both clinically and capillaroscopically; 175 were reevaluated after a mean±SD of 3.34±1.48 years. Sixty-two healthy volunteers served as controls. Qualitative assessment of NVC images permitted categorization of patients to a normal, early, active or late capillaroscopic pattern. Capillary loss, dilated, giant or ramified capillaries and micro-hemorrhages were evaluated by a semi-quantitative score (0-3), derived as the mean of three 1mm fields in each of the 2nd, 3rd, 4th and 5th finger of both hands. FVC and DLCO were recorded, if performed within 6 months of NVC. FVC and DLCO deterioration were considered clinically significant if >10% and >15%, respectively. Skin thickening was measured using the modified Rodnan Skin Score (mRSS). mRSS deterioration was considered clinically significant if >3.5. First occurrence of digital ulcers in patients with no prior such history and vital status were also recorded at follow-up.

Results: Capillary loss score had the highest diagnostic accuracy at discriminating patients with an SSc-spectrum disorder from patients with RP of different etiology and from controls, as defined by ROC curve analysis [AUC (95% CI)=0.925 (0.893-0.956)], followed by dilatation score [AUC (95% CI)=0.904 (0.807- 0.938)] and giant score [AUC (95% CI)=0.856 (0.810-0.902)]. By contrast, micro-hemorrhages [AUC (95% CI)=0.727

(0.669-0.786)] and ramification scores [AUC (95% CI)=0.588 (0.521-0.654)] did not perform equally well. Notably, clinical correlations of capillaroscopic parameters in SSc were found not to be consistent over time, when longitudinally assessed at two different time points by univariate and multivariate analysis. Binary logistic regression analysis indicated that baseline capillaroscopic pattern could predict occurrence of a combined adverse disease outcome (FVC deterioration>10% and/or DLCO deterioration>15% and/or mRSS deterioration>3.5 and/or first occurrence of digital ulcers and/or death), after a mean±SD follow up of 3.28±1.45 years in 94 SSc patients with available follow-up data (OR=3.43, p=0.031 for active versus early pattern, OR=8.778, p=0.007 for late versus early pattern).

Conclusion: Dilatation score performs best of all semi-quantitative NVC parameters in diagnosing SSc and although clinical correlations of capillaroscopic findings change over time, an active or late capillaroscopy pattern at baseline is associated with an adverse prognosis.

Disclosure of Interests: Vasiliki-Kalliopi Bournia: None declared, Konstantinos Kottas: None declared, Stylianos Panopoulos: None declared, George Konstantonis: None declared, Alexios Iliopoulos: None declared, Athanasios Tzioufas Grant/research support from: ABBVIE, PFIZER, AMGEN, NOVARTIS, GSK, Petros Sfrikakis: None declared, PANAYIOTIS VLA-CHOYIANNPOULOS: None declared

DOI: 10.1136/annrheumdis-2019-eular.4393

FRI0314 SENSITIVITY TO CHANGE AND RESPONSIVENESS TO TREATMENT OF RENAL RESISTIVE INDEX (RRI) IN SYSTEMIC SCLEROSIS (SSC)

Cosimo Bruni¹, Edoardo Rosato², Antonietta Gigante², Vanessa Maestripietri¹, Giulia Tesi¹, Marco Chiostrì¹, Gemma Lepri¹, Silvia Bellando Randone¹, Serena Guiducci¹, Sergio Castellani¹, Maria Boddi¹, Marco Matucci-Cerinic¹. ¹Careggi University Hospital, Firenze, Italy; ²Sapienza University of Rome, Roma, Italy

Background: RRI may identify any problem with the blood flow in the renal artery and in the parenchyma. It was shown to be increased in SSc patients and being in relationship with both vascular and fibrotic SSc-related manifestations.

Objectives: to test sensitivity to change and responsiveness to treatment of RRI in SSc.

Methods: patients fulfilling the 2013 ACR/EULAR classification criteria for SSc were enrolled from two SSc-care units, if RRI was determined at least twice since being diagnosed. Data regarding SSc clinical manifestations, instrumental and laboratory evaluation for renal, cardiac and cardiovascular involvement, as well as ongoing immunosuppressive and vasoactive/vasodilating treatment, were collected both at baseline and follow-up RRI measurements.

Results: 230 patients [aged 57 (48-67) years, 12.6% males] were enrolled in the study, with baseline RRI value of 0.68 (IQR 0.07). In a mean 3.4 years follow-up, RRI showed a median change (ΔRRI) of +0.02 (IQR 0.05). ΔRRI was significantly correlated with ΔsPAP (R=0.173, p=0.023) and it was significantly higher in patient with new onset of pulmonary arterial hypertension (0.08 ± 0.02 vs 0.03 ± 0.05; p=0.038). On Cox univariate regression analysis, time from disease onset, ΔssPAP and ΔRRI predicted new PAH, while ΔsPAP was the only independent predictors at multivariate regression analysis. Regarding treatment, Sildenafil exposure determined a significantly lower increase in ΔRRI, with a progressive long-term effect. Conversely, CCBs and Iloprost treated patients showed a significantly higher increase of ΔRRI, which was determined by a higher DU burden. No difference was seen when immunosuppressive treatment was evaluated.

Conclusion: RRI is sensitive to change and reflects, in particular, the worsening of cardio-pulmonary vascular involvement (increase in ΔsPAP and new PAH onset). Moreover, it shows a possible protective effect of Sildenafil in reducing pulmonary vasculature manifestations.

Disclosure of Interests: Cosimo Bruni: None declared, Edoardo Rosato: None declared, Antonietta Gigante: None declared, Vanessa Maestripietri: None declared, Giulia Tesi: None declared, Marco Chiostrì: None declared, Gemma Lepri: None declared, Silvia Bellando Randone: None declared, Serena Guiducci: None declared, Sergio Castellani: None declared, Maria Boddi: None declared, Marco Matucci-Cerinic Grant/research support from: Actelion, MSD, Pfizer, BMS, Chemomab, Sanipedia, Speakers bureau: Actelion, BMS; MSD, Janssen

DOI: 10.1136/annrheumdis-2019-eular.3515