patients (30%) manifested thrombocytopenia and 4 patients (20%) manifested anemia.

- Cardiac symptoms included shortness of breath (n=18), palpitation (n=6), edema of lower extremity (n=4), fatigue (n=5) and/or cough (n=3). All patients were found to have cardiomegaly by echocardiography (UCG) or chest radiography. Left ventricular ejection fraction ($47 \pm 15\%$) were decreased. Ventricular wall motion abnormality was found in 5 patients (25%) by UCG. 15 patients (75%) manifested cardiac arrhythmia in electrocardiogram, with frequent premature ventricular contractions were the most common (n=10, 50%). Myocardial injury was found in 14 patients (70%) by electrocardiogram. Myocarditis was found in 7 patients (35%) by myocardial enzyme assay.
- Nine patients (45%) died, the causes of death were heart failure (n=4, 45%), sudden death (n=3, 33%) and ventricular tachycardia (n=2, 22%). 5 patients (55.6%) died within 1 month after the onset of cardiac symptoms. 3 patients died within 2 month, and one patient with implantable cardioversion defibrillation (ICD) died 2 years after ICD implanted. Ten patients survived in a mean 6 (1-10) years follow-up.
- Six patients (67%) manifested thrombocytopenia and 4 patients (44%) manifested anemia in the death group. However, only one patient (9.1%) in survival group manifested thrombocytopenia (p=0.007) and anemia (p=0.069). There were more organs and systems involved in the death group, compared to the survival group (4 ± 2 vs 2 ± 1 , p=0.044). 4 patients had SRC in death group, while 2 patients had SRC in survival group. 6 patients had pulmonary fibrosis in death group, while 5 patients in survival group (Table 1).

Table 1. Comparison between the death group and the survival group in Systemic Sclerosis cardiomyopathy patients.

	Death group (n=9)	Survival group (n=11)	p
Thrombocytopenia, n (%)	6(66.7)	1(9.1)	0.007
Anemia, n (%)	4(44.4)	1(9.1)	0.069
Pulmonary fibrosis, n (%)	6(66.7)	5(45.5)	0.343
Pulmonary arterial hypertension, n (%)	3(33.3)	4(36.4)	0.888
Scleroderma renal crisis, n (%)	4(44.4)	2(18.2)	0.202
Esophageal involvement, n (%)	3(33.3)	2(18.2)	0.436
Anti-SCL 70 positive, n (%)	5(55.6)	5(50)	0.653
Myocarditis, n (%)	4(44.4)	3(27.3)	0.423
Numbers of organ involved, mean (SD)	4(2)	2(1)	0.044

Conclusion: Thrombocytopenia and multi-organ involvement might be poor prognostic factors for patients of SSc-cardiomyopathy.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2019-eular.4082

FRI0319 ESOPHAGEAL EROSIONS PREDICT PROGRESSION OF LUNG DISEASE IN PATIENTS WITH SYSTEMIC SCLEROSIS

Enrico De Lorenzis¹, Gerlando Natalello¹, Laura Gigante¹, Lucrezia Verardi¹, Umberto La Porta¹, Giovanni Battista Canestrari¹, Ludovica Berardini², Silvia Laura Bosello³, Elisa Gremese^{1,3}. ¹Catholic University of the Sacred Heart, Institute of Rheumatology, Rome, Italy; ²Fondazione Policlinico Universitario A. Gemelli IRCCS, Department of Respiratory Medicine, Rome, Italy; ³Fondazione Policlinico Universitario A. Gemelli IRCCS, Department of Rheumatology, Rome, Italy

Background: Interstitial lung disease (ILD) is the leading cause of death in Systemic Sclerosis (SSc) but its pathogenesis is not fully understood. Esophageal disease is common in SSc and micro aspiration of both acid and non-acid gastroesophageal reflux could be involved in the pathogenesis of ILD. Esophagogastroduodenoscopy (EGD) is an essential tool to evaluate disease severity of upper gastrointestinal tract involvement in SSc.

Objectives: The objective of the present study is to assess the role of EGD in predicting pulmonary functional deterioration in SSc patients.

Methods: One hundred and fifty patients with SSc and suspected esophageal involvement underwent EGD. Pulmonary function tests were performed at baseline and after a 36-months follow-up. Patients were characterized for disease phenotype, BMI, smoking exposure and medication history. A significant ILD progression was defined as a relative decline $\geq 10\%$ of FVC or a concomitant decline $\geq 5\%$ of FVC and $\geq 15\%$ of DLCO.

Results: One hundred and thirty-six patients (90.5%) were female with a mean age of 55.6 ± 13.8 years and 12.1% were active smokers. Fifty

patients (33.3%) had a diffuse cutaneous disease; 37.4% and 40.8% were positive for anti-centromere and anti-Scl70 antibodies respectively. The mean disease duration from the first non-Raynaud symptom was 5.9 ± 6.7 years. Sixty-one patients (40.8%) showed EGD signs of reflux esophagitis. Among them, 31.3% had an erosive form (9.5% grade A, 15.6% grade B, 4.8% grade C and 1.4% grade D according to Los Angeles classification). At the baseline, 23.1% of the patients had a FVC $\leq 80\%$ and 45.6% had a DLCO $\leq 50\%$. Patients with erosive esophagitis did not differ in terms of sex, age, duration and disease variant, positivity for anti-centromere, skin score values, FVC and DLCO at baseline compared to patients without erosions, but had a lower prevalence of anti-Scl70 (28.3 vs 52.5%, p=0.005) and active smoking (20.0 vs 8.4%, p=0.05). At follow-up, patients with esophageal erosions showed a greater relative decrease in FVC ($3.4 \pm 9.3\%$ vs $1.7 \pm 12.0\%$; p=0.013) without significant differences in the DLCO change. Overall, 11.0% of patients presented pulmonary disease progression. The presence of esophageal erosions was associated with a significantly greater risk of lung disease progression (OR 5.3, 95% CI 1.7-16.8, p=0.004) after paired correction for sex, age, duration of disease, auto-antibodies, skin involvement variant, baseline FVC and DLCO, smoke exposure and therapy with immunosuppressants, proton pump inhibitors, prokinetics, antiplatelet agents and prostanoids.

Conclusion: SSc patients with erosive esophagitis present a higher risk of progression of interstitial lung disease. This evidence supports a role of micro-aspiration of gastric contents in the development of inflammation and fibrosis of the airways.

Disclosure of Interests: Enrico De Lorenzis: None declared, Gerlando Natalello: None declared, Laura Gigante: None declared, Lucrezia Verardi: None declared, Umberto La Porta: None declared, Giovanni Battista Canestrari: None declared, Ludovica Berardini: None declared, Silvia Laura Bosello: None declared, Elisa Gremese Consultant for: AbbVie, BMS, Celgene, Janssen, Lilly, MSD, Novartis, Sanofi, UCB, Roche, and Pfizer, Speakers bureau: BMS, Speakers bureau: Roche, Speakers bureau: AbbVie, BMS, Celgene, Janssen, Lilly, MSD, Novartis, Sanofi, UCB, Roche, and Pfizer

DOI: 10.1136/annrheumdis-2019-eular.6401

FRI0320 CN-1A AUTOANTIBODIES ARE SPECIFIC FOR SPORADIC INCLUSION BODY MYOSITIS

Louise Pyndt Diederichsen^{1,2}, Sine Søndergaard Korsholm¹, Line Vinderslev Iversen³, Christoffer Tandrup Nielsen², Marie-Louise From Hermansen², Søren Jacobsen², Nanna Witting⁴, Markus E. Krogager⁴, Tina Friis⁵. ¹Odense University Hospital, Department of Rheumatology, 5000 Odense, Denmark; ²Rigshospitalet, Department of Rheumatology, DK-2100 Copenhagen, Denmark; ³Bispebjerg Hospital, Department of Dermatology, DK-2400 København NV, Denmark; ⁴Rigshospitalet, Department of Neurology, Copenhagen, Denmark; ⁵Statens Serum Institut, Department of Autoimmunology and Biomarkers, DK-2300 Copenhagen S, Denmark

Background: Sporadic inclusion body myositis (sIBM) is a subgroup of the idiopathic inflammatory myopathies (IIMs) and is characterized by both degenerative and autoimmune features. Unlike other IIMs, myositis-specific autoantibodies had not been found in sIBM patients until recently.

Objectives: We aimed to establish the prevalence and clinical associations of anti-CN-1A in a large Danish cohort with connective tissue diseases (CTD).

Methods: In a cross-sectional study design, a total of 568 participants (183 IIMs (55 sIBM, 128 non-sIBM: dermatomyositis (DM), polymyositis (PM), immune-mediated necrotizing myopathy (IMNM)), 164 systemic lupus erythematosus (SLE), 121 systemic sclerosis (SSc), and 100 blood donors (BD)) were tested for the presence of: a) CN-1A autoantibodies using a commercial Anti-CN-1A ELISA and b) myositis specific and associated autoantibodies (anti-Jo-1, anti-PL-7, anti-PL-12, anti-OJ, anti-EJ, anti-SRP, anti-Mi-2 α / β , anti-MDA5, anti-TIF1 γ , anti-NPX2, anti-SAE1, anti-PM-Scl75, anti-PM-Scl100, anti-Ro52, anti-Ku) using a commercial lineblot kit. The patients were classified according to the ACR classification criteria for IIMs (2017), SLE (1997) and SSc (1980), respectively. Clinical features were compared between anti-CN-1A positive and anti-CN-1A negative sIBM patients using two-sample t-test and Mann-Whitney test for continuous normally and non-normally distributed data and Chi-square test for categorical data, as appropriate.

Results: CN-1A antibodies were predominantly found in IIM patients. In the sIBM cohort, 24 patients (43.6%) were anti-CN-1A positive vs. 25 (19.5%) in the non-sIBM myositis cohort and 17 (10%) in the SLE cohort. None of the participants in the SSc or the BD cohorts had

presence of anti-cN-1A. Anti-cN-1A positivity had a sensitivity of 43.6% and a specificity of 91.8% for sIBM. The positive and negative predictive values were 36.4% and 93.8%, respectively.

There was no significant difference in gender, age at study entry, age at symptom onset, duration of symptoms or max creatine kinase (CK) levels during disease course between the anti-cN-1A positive and negative sIBM patients. Dysphagia was present in 19 (79%) of the anti-cN-1A positive and in 17 (55%) of the anti-cN-1A negative sIBM patients (P = 0.06).

Conclusion: Antibodies against cN-1A are the first and so far the only serological marker for sIBM. Our data showed that cN-1A autoantibodies are specific for sIBM and further corroborate the potential diagnostic role of cN-1A autoantibodies in this distinct subgroup of myositis.

Disclosure of Interests: Louise Pyndt Diederichsen: None declared, Sine Søndergaard Korsholm: None declared, Line Vinderslev Iversen: None declared, Christoffer Tandrup Nielsen: None declared, Marie-Louise From Hermansen: None declared, Søren Jacobsen: None declared, Nanna Witting: None declared, Markus E. Krogager: None declared, Tina Friis Grant/research support from: Anti cN-1A ELISA kits and EUROLINE Auto-immune Inflammatory Myopathies 16 AG kits have been provided for a project free of charge from Euroimmun.
DOI: 10.1136/annrheumdis-2019-eular.7264

FRI0321

PERFORMANCE OF DIFFERENT PULMONARY HYPERTENSION SCREENING ALGORITHMS IN PATIENTS WITH SYSTEMIC SCLEROSIS PATIENTS

Mustafa Erdogan¹, Burcak Kilickiran Avci², Yagmur Ersoy³, Cansu Ebrer², Zeki Ongen², Gul Ongen⁴, Vedat Hamuryudan¹, Gulen Hatemi¹. ¹Istanbul University -Cerrahpasa Medical Faculty, Internal Medicine, Division of Rheumatology, Istanbul, Turkey; ²Istanbul University -Cerrahpasa Medical Faculty, Department of Cardiology, Istanbul, Turkey; ³Istanbul University -Cerrahpasa Medical Faculty, Istanbul, Turkey; ⁴Istanbul University -Cerrahpasa Medical Faculty, Department of Pulmonary Diseases, Istanbul, Turkey

Background: Pulmonary hypertension (PH) is an important cause of morbidity and mortality in patients with systemic sclerosis (SSc). Different screening algorithms have been proposed for identifying patients who have a high probability of PH and require right heart catheterization (RHC), which is the gold standard for diagnosing PH.

Objectives: To compare the performance of PH screening algorithms in our patients with SSc.

Methods: Forty-eight consecutive patients, fulfilling ACR/EULAR 2013 SSC criteria, were screened for PH using the, 2015 ESC/ERS, DETECT and ASIG algorithms. Pulmonary function tests (PFT), diffusing capacity of the lung for carbon monoxide (DLCO), trans-thoracic echocardiography, serum NT-proBNP, uric acid assay and high-resolution computed tomography (HRCT) were performed as needed. Patients with known PH, severe interstitial lung disease and severe left ventricular dysfunction were not included. RHC was performed in all patients with positive screening according to any one of the screening algorithms. Patients with PH were classified according to the updated PH classification criteria. Sensitivity and specificity of the 3 screening algorithms were evaluated according to the established cut-off value of 25 mmHg for mean systolic pulmonary artery pressure and for the recently proposed cut-off value of 20 mmHg.

Results: Among the 48 SSc patients, 15 were excluded due to already diagnosed PH (n=4), left ventricular dysfunction (n=4), no measurable tricuspid regurgitation velocity (TRV) (n=5) and coexisting lung cancer (n=2). Among the remaining 34 patients, 16 required RHC according to at least one of the screening algorithms. Demographic and clinical features of remaining 34 patients were summarized in Table 1. Number of patients who had suspected pulmonary hypertension and required RHC according to ESC/ERS 2015, DETECT and ASIG were 8 (%25), 9 (% 27), and 13 (%41) respectively (Figure 1). Among the 14 who had RHC, PH was present in 3 patients according to the 25-mmHg cut-off (Group1 in 1, Group 2 in 1, Group 3 in 1) and in 8 patients according to the 20-mmHg cut-off (Group 1 in 5, Group 2 in 2, Group 3 in 1). The sensitivity and specificity of each algorithm is presented in Table 2. Sensitivity was similar at 100% for the 3 algorithms, but the ESC/ERS algorithm had better specificity, when PH was diagnosed with the 25-mmHg cut-off. For the 20-mmHg cut-off, both the sensitivity and the specificity were better with the ESC/ERS algorithm.

Conclusion: The ESC/ERS algorithm seems to have a better performance for detecting PH in patients with SSc. A limitation of this study was that RHC was not performed in patients who did not fulfill criteria according to any of the screening algorithms. The sensitivities may be lower than what we propose if there are patients with PH who are asymptomatic and not captured with any of the algorithms.

Table 1. Demographic and Clinical Features of Patients											
	All Patients n=33	ESC/ERS screen positive, n=8 (24%)	ESC/ERS screen negative, n=25 (76%)	DETECT screen positive, n=9 (27%)	DETECT screen negative, n=24 (73%)	ASIG screen positive, n=13 (41%)	ASIG screen negative, n=20 (59%)	Any of the algorithms, n=18 (55%)	Any of the algorithms, n=20 (59%)		
Age, median [IQR]	51 (46-75)	66.5 (59-79.5)	48 (38-75)	66.5 (58-75)	48 (38-51.5)	61 (52-75)	48 (38-51)	57 (50-68)	48 (38-51)		
Male, n (%)	100 (100)	100 (100)	100 (100)	100 (100)	100 (100)	100 (100)	100 (100)	100 (100)	100 (100)		
Female, n (%)	13.5 (15-20.5)	52 (39-57)	65 (51-79.5)	24 (5-38)	12 (5-25)	14 (5-37)	13 (5-19)	14 (5-25)	12 (5-17.5)		
Duration of SSc symptoms, year [IQR]	10 (4-29)	2 (0-75)	8 (0-13)	1 (0-13)	9 (2-27)	4 (1-13)	6 (2-29)	4 (1-29)	6 (2-10)		
SSc pattern, n (%)	24 (34 (71))	6 (8 (75))	18 (26 (69))	7 (8 (87))	15 (24 (63))	8 (13 (60))	8 (23 (61))	10 (31)	14 (20 (70))		
SSc related serology, n (%)	18 (34 (69))	3 (6 (50))	15 (20 (75))	5 (7 (71))	13 (18 (72))	6 (10 (60))	12 (36 (75))	6 (10 (60))	12 (36 (75))		
Anticentromere (+)	3 (14 (19))	3 (7 (43))	0 (0 (0))	2 (9 (40))	3 (20 (9))	0 (0 (0))	0 (0 (0))	0 (0 (0))	0 (0 (0))		
Presence of ILD, n (%)	37 (34 (60))	4 (8 (50))	13 (21 (57))	4 (9 (44))	12 (24 (60))	7 (13 (64))	10 (21 (48))	7 (14 (50))	10 (20 (50))		
Presence of digital ulcer, n (%)	23 (34 (66))	4 (8 (50))	17 (24 (71))	6 (9 (67))	14 (22 (64))	8 (13 (62))	13 (29 (68))	9 (14 (54))	12 (20 (60))		
Presence of GIG involvement, n (%)	14 (34 (41))	6 (8 (75))	8 (12 (36))	4 (9 (44))	12 (24 (60))	7 (13 (64))	7 (21 (33))	7 (14 (50))	7 (20 (35))		
Presence of telangiectasia, n (%)	14 (34 (41))	5 (8 (63))	9 (19 (47))	4 (9 (44))	10 (24 (62))	7 (13 (64))	7 (21 (33))	7 (14 (50))	7 (20 (35))		
sPAP on TTE, median (mmHg) [IQR]	28.5 (23.5-33)	36.5 (29-25-46.5)	27 (21-35-31)	33 (25-44)	27 (21-31)	30 (25.5-40.5)	27 (19.5-31)	30 (25.75-40.25)	27 (18.75-31)		
FVC, median (% predicted) [IQR]	86 (69.5-102.5)	92 (58-115.25)	83 (70-99.5)	92 (63.5-114)	79 (69-100)	92 (67-108.75)	76 (65.5-99)	90 (72-107.5)	74 (69.25-100)		
DLCO, median (% predicted) [IQR]	53.5 (41.25-71)	47 (38.75-56.5)	62 (49-78.5)	55 (46-71)	52 (43-71.5)	50 (35.5-58.5)	63 (46-79)	51 (37.25-62.5)	62 (44-77)		
FVC / DLCO ratio, median [IQR]	1.4 (1.1-1.74)	2.31 (1.58-2.18)	1.3 (1.1-1.4)	1.6 (1.28-2.18)	1.40 (1.13-1.58)	1.78 (1.35-2.12)	1.35 (1.1-1.43)	1.71 (1.29-2.12)	1.36 (1.04-1.44)		
NT-proBNP, median (pg/mL) [IQR]	137 (63-256)	284 (162-778)	120 (52-165)	451 (120-887)	120 (52-158)	284 (124-25-695)	120 (38-158)	284 (124-25-695)	120 (38-158)		
Presence of dyspnea, n (%)	7 (8 (88))	7 (26 (21))	5 (9 (56))	9 (24 (88))	7 (13 (33))	7 (13 (33))	7 (14 (50))	7 (20 (35))	7 (20 (35))		
WHO-FC, n (%)	1 (3 (4))	5 (7 (2))	7 (10 (27))	4 (9 (40))	9 (10 (27))	6 (7 (8))	7 (7 (10))	4 (7 (8))	7 (7 (10))		
Class 1	1 (3 (4))	5 (7 (2))	7 (10 (27))	4 (9 (40))	9 (10 (27))	6 (7 (8))	7 (7 (10))	4 (7 (8))	7 (7 (10))		
Class 2	1 (3 (4))	5 (7 (2))	7 (10 (27))	4 (9 (40))	9 (10 (27))	6 (7 (8))	7 (7 (10))	4 (7 (8))	7 (7 (10))		
Class 3	1 (3 (4))	5 (7 (2))	7 (10 (27))	4 (9 (40))	9 (10 (27))	6 (7 (8))	7 (7 (10))	4 (7 (8))	7 (7 (10))		
Class 4	1 (3 (4))	5 (7 (2))	7 (10 (27))	4 (9 (40))	9 (10 (27))	6 (7 (8))	7 (7 (10))	4 (7 (8))	7 (7 (10))		

Table 2. Performance of the PH screening algorithms

	Performance for 25-mmHg cut-off			Performance for 20-mmHg cut-off		
	ESC/ERS	ASIG	DETECT	ESC/ERS	ASIG	DETECT
Sensitivity, %	100	100	100	75	100	100
Specificity, %	54	9	40	67	17	0
PPV, %	37.5	23	33	75	61	44
NPV, %	100	100	100	67	0	0

Disclosure of Interests: Mustafa Erdogan: None declared, Burcak Kilickiran Avci: None declared, Yagmur Ersoy: None declared, Cansu Ebrer: None declared, Zeki Ongen: None declared, Gul Ongen: None declared, Vedat Hamuryudan Consultant for: Abbvie, Amgen, BMS, Jansen, MSD, Pfizer, UCB, Speakers bureau: Abbvie, Amgen, BMS, Jansen, MSD, Pfizer, UCB, Gulen Hatemi Consultant for: Abbvie, Amgen, BMS, Jansen, MSD, Pfizer, UCB, Speakers bureau: Abbvie, Amgen, BMS, Jansen, MSD, Pfizer, UCB.

DOI: 10.1136/annrheumdis-2019-eular.2662

FRI0322

AUTOANTIBODIES PROFILE INFLUENCE ON SYSTEMIC SCLEROSIS INTERSTITIAL LUNG DISEASE. A TERTIARY SPANISH HOSPITAL EXPERIENCE

Antía García Fernández¹, Iván del Bosque Granero¹, Laura Calvo Sanz¹, Nicolás Almeida Arostegui², Luis Gorospe Sarasua², Carlos De la Puente Bujidos¹. ¹Ramón y Cajal University Hospital, Rheumatology, Madrid, Spain; ²Ramón y Cajal University Hospital, Radiology, Madrid, Spain

Background: Systemic Sclerosis (SSc) is a rare and heterogeneous connective tissue disease (CTD) characterized by skin fibrosis, vasculopathy/vascular damage and potential visceral impairment. Interstitial lung disease (ILD) constitutes the leading cause of mortality and requires close periodical assessment and follow-up. Diffuse cutaneous sclerosis and specific autoantibody profile (anti SCL70, anti Th/To, Anti U3 RNP and anti PmScI) are considered ILD development risk factors. In contrast, positivity to anticentromere has been considered a protective factor to develop a clinical significant ILD.

Objectives: To assess ILD frequency and severity (extension and functional impairment) in SSc patients, analyzing the association with the different autoantibodies.

Methods: Retrospective, descriptive study of patients meeting EULAR/ACR 2013 SSc criteria and had a HRTC performed at a tertiary Spanish hospital from 1975 to 2018. One hundred and eight patients were included. HRTCs were graded by two radiologists according to Goh et al semi-quantitative score. Three groups were established according to the presence of different autoantibodies: anticentromere (ACA), antitopoisomerase (ATA), and positivity to other ANAs (nucleolar pattern and other specificities).

Results: Clinical and laboratory characteristics are presented in table 1. Thirty-three patients had pulmonary involvement, 6 were ACA+, 18 ATA+ and 9 had other ANA specificities. The probability of not having pulmonary involvement among ACA+ and of having pulmonary involvement among ATA+ showed statistic significance (p<0.001 and p<0.001). Usual interstitial Pneumonia (UIP) was the most frequently reported pattern (6 patients), followed by Non-Specific Interstitial Pneumonia (NSIP 11 patients); Six patients did not meet the radiological criteria for neither UIP or NSIP. No statistical significant difference was found among radiological pattern and autoantibody profile. Regarding extension: 15 patients