<0.05, $q = 0.02$), supporting the idea that they should be involved in the pathophysiology of the disease.

Conclusion: These preliminary data confirm that the innate immune cells could play an important role in AxSpA. MAIT cells are at the forefront of the expression of IL-17A before $\gamma\delta$ T, CD4+T and CD8+T. Neutrophils do not appear to participate in the production of IL-17A, but the high expression of AS linked genes in these cells suggests their involvement in AxSpA.

REFERENCES:

- [1] Appel H, et al. Analysis of IL-17(+) cells in facet joints of patients with spondyloarthritis suggests that the innate immune pathway might be of greater relevance than the Th17-mediated adaptive immune response. *Arthritis Res Ther*. 2011
- [2] Kenna TJ, et al. Enrichment of circulating interleukin-17-secreting interleukin-23 receptor-positive $\gamma\delta$ T cells in patients with active ankylosing spondylitis. *Arthritis Rheum*. 2012
- [3] Gracey E, et al. IL-7 primes IL-17 in mucosal-associated invariant T (MAIT) cells, which contribute to the Th17-axis in ankylosing spondylitis. *Ann Rheum Dis*. 2016
- [4] Taylor PR, et al. Activation of neutrophils by autocrine IL-17A-IL-17RC interactions during fungal infection is regulated by IL-6, IL-23, ROR γ t and dectin-2. *Nat Immunol*. 2014

Disclosure of Interests: nicolas rosine: None declared, Surya Koturan: None declared, Hanane Yahia: None declared, Claire Leloup: None declared, Elisabetta Bianchi: None declared, Corinne Miceli Richard Grant/ research support from: MSD, Pfizer, AbbVie, Biogen, UCB, Novartis, Consultant for: Abbvie, Novartis, BMS, Lars Rogge: None declared
DOI: 10.1136/annrheumdis-2019-eular.6245

FRI0362 HIGH-DIMENSIONAL MULTIPARAMETRIC CHARACTERIZATION OF THE REGULATORY T CELLS LANDSCAPE IN SPONDYLOARTHRITIS

Davide Simone, India Brough, Liye Chen, Frank Penkava, Anna Ridley, Hui Shi, Hussein Al Mossawi, Paul Bowness. *NDORMS, University of Oxford, Oxford, United Kingdom*

Background: The Spondyloarthritis (SpA) are immune-mediated conditions characterised by spinal and joint inflammation. The pathogenic role of Th17 lymphocytes has been shown by multiple studies but few reports exist on the phenotype of regulatory T cells (Tregs) and their role in the course of the disease. Studying Tregs is particularly challenging because of their heterogeneous phenotype and potential plasticity at the site of inflammation. Given the complexity of the Treg landscape, a multi-dimensional approach, including both protein and gene analysis, offers novel mechanistic insights that can be leveraged to develop new treatments.

Objectives: 1) To describe the Treg phenotype in SpA patients compared to controls. 2) To identify differential expression of markers, such as trafficking molecules or co-inhibitory molecules, within Tregs at the inflammatory site. 3) To define the gene expression landscape in Tregs in the peripheral blood and synovial fluid in patients with active SpA.

Methods: A total of 61 patients with SpA (38 with Ankylosing Spondylitis (AS), 23 with Psoriatic Arthritis (PsA)) and 16 age-matched healthy controls were recruited. Peripheral blood (PB) and paired synovial fluid (SF) mononuclear cells ($n=8$) were also analyzed. Isolated mononuclear cells were stained with 3 multicolor flow cytometry panels, including a total of 35 surface and intracellular protein markers. Manual gating was done in parallel with unsupervised data analysis using FlowSOM and SPADE. Cells from blood and synovial fluid from three treatment-naïve PsA patients were isolated and their RNA sequenced at a single cell level with the 10x Genomics platform.

Results: Whereas no major difference in the Treg frequency was observed comparing the PB of SpA patients and healthy controls, in the SF we observed a higher Treg frequency, with a striking prevalence of the memory (CD45RA-) compartment (mean: 11.7 vs 4.7%; $p=0.01$) and a higher expression of Foxp3 ($p=0.04$). Trafficking markers demonstrated considerable heterogeneity, as visualized on SPADE analysis, mirroring the classification of T helper cells, but little difference in terms of relative frequency compared to healthy controls. Within the Treg compartment we identified populations of Helios-negative Th17-like cells able to secrete higher levels of IL-17A (mean: 2.8 vs 1.2%; $p=0.02$) while expressing normal levels of Foxp3; and putative regulatory CD8+ T cells expressing classical Treg features (Foxp3, CTLA4).

The analysis of the single cell transcriptomic data confirmed a high degree of heterogeneity, both in the SF and the PB, with consistent

findings across patients. Unsupervised clustering identified different subsets that share a core of regulatory transcripts, onto which additional programs are added, including a very distinct Th17-like module. Of note, various Treg subsets express preferentially different co-inhibitory genes, suggesting a functional specialization.

Conclusion: High-dimensional immunoprofiling in SpA patients shows normal frequency of Tregs in the PB, but increased Tregs with activated phenotype in the inflammatory site. The presence within the Treg population of Th17-like and CD8+ populations are intriguing preliminary findings that require further evaluation. Preliminary transcriptomic analysis confirms the presence of specialized subsets within the Treg compartment.

Disclosure of Interests: Davide Simone: None declared, India Brough: None declared, Liye Chen: None declared, Frank Penkava: None declared, Anna Ridley: None declared, Hui Shi: None declared, Hussein Al Mossawi Grant/research support from: UCB, Paul Bowness Grant/ research support from: Merck, GSK, Celgene

DOI: 10.1136/annrheumdis-2019-eular.1749

FRI0363 AUTOANTIBODIES TO THREE NOVEL PEPTIDES IN EARLY AXIAL SPONDYLOARTHRITIS IN TWO INDEPENDENT COHORTS

Dana Quaden¹, Patrick Vandormael¹, Piet Geusens^{1,2,3}, Johan Vanhooft², Kurt de Vlam^{4,5}, Veerle Somers¹. ¹Hasselt University, Biomedical Research Institute, Diepenbeek, Belgium; ²ReumaClinic, Genk, Belgium; ³Maastricht University Medical Center, Internal Medicine, Rheumatology, Maastricht, Netherlands; ⁴University Hospitals Leuven, Division of Rheumatology, Leuven, Belgium; ⁵University Hospitals Leuven, Skeletal Biology and Engineering Research Center, Department of Development and Regeneration, Leuven, Belgium

Background: Diagnosis of axial spondyloarthritis (axSpA) is challenging since clinical manifestations, such as inflammatory back pain, peripheral arthritis, enthesitis and inflammatory bowel disease, often overlap with other disorders. Current laboratory markers for axSpA, Human Leukocyte Antigen (HLA)-B27 and C-reactive protein (CRP) are not sufficiently specific for diagnosis. Despite being considered a "seronegative" disease, emerging evidence supports the involvement of antibodies in axSpA. In order to identify novel autoantibodies in axSpA patients, we recently screened an axSpA cDNA phage display library for reactivity with immunoglobulin G (IgG) antibodies in plasma of early axSpA patients. This resulted in autoantibodies to 9 novel University of Hasselt (UH) axSpA peptide targets, corresponding to fragments of known proteins and novel linear peptides.

Objectives: The aim of this study was to determine the diagnostic potential of autoantibodies to the 9 novel UH axSpA peptides in axSpA patients and controls from 2 independent cohorts.

Methods: Using enzyme-linked immunosorbent assays (ELISA), presence of antibodies to the 9 UH axSpA peptides expressed on phage particles was determined in 76 early axSpA patients, 75 chronic low back pain patients (LBP), 60 early rheumatoid arthritis patients (RA) and 94 healthy controls (HC) from the UH cohort. Antibody reactivity was further validated in 174 patients from the Leuven Spondyloarthritis (Biologics) Cohort ((Bio)SPAR), including 79 early axSpA patients.

Results: Antibody reactivity against at least one of 9 novel UH axSpA peptides was found in 54% (41/76) of early axSpA patients, 26% (24/94) of HC ($p=0.0002$), 39% (29/75) of LBP ($p=0.0731$) and 38% (23/60) of RA patients ($p=0.0845$) from the UH cohort, as compared to 43% (74/174) of the axSpA patients from the (Bio)SPAR cohort. By combining the three UH axSpA peptides with highest positive likelihood ratios (LR+) into a panel, antibodies against these 3 peptides were detected in 14.2% (22/155) of early axSpA patients from the combined UH and (Bio)SPAR cohorts and in only 5% (4/75) of persons with LBP ($p=0.0484$), resulting in a specificity of 95%. The LR+ for confirming axSpA using antibodies to these 3 UH axSpA peptides was 2.7, which is the same as for the currently used laboratory marker CRP. Assuming a 5% pretest probability of axSpA in persons with LBP, a combination of the presence of inflammatory back pain (LR+ 3.1) and a positive test result for the laboratory markers HLA-B27 (LR+ 9.0) and CRP (LR+ 2.5) provides a disease (posttest) probability of 79%. When we added a positive test result for the presence of antibodies to the 3 UH axSpA peptides (LR+ 2.7), posttest probability could be increased to 91%.

Conclusion: Antibodies to 3 UH axSpA peptides were significantly more present in early axSpA patients compared to LBP and could provide a novel tool for objective diagnosis of a subset of axSpA patients.

Disclosure of Interests: Dana Quaden: None declared, Patrick Vandormael: None declared, Piet Geusens Grant/research support from: Research support, consultant and/or speaker fees from: Pfizer, Abbott, Eli