Both *HLA* class I and II regions identified as genome-wide significant susceptibility loci for adult-onset Still's disease in Chinese individuals

Adult-onset Still's disease (AOSD) is a rare autoinflammatory condition with unclear aetiology and highly heterogeneous manifestations, such as spiking fever, skin rash and arthralgia (or arthritis).^{1,2} AOSD is usually the result of both genetic and environmental factors. However, only a few genetic loci have been associated with AOSD, and none of them have reached the threshold for genome-wide significance (GWS) of $p < 5 \times 10^{-8}$.^{2,3} For example, *HLA-DRB1* was strongly associated with AOSD in previous studies (the *p* value for the most significant association was 8.60×10^{-6}).³ *HLA-DRB1* also influences the risk of systemic juvenile idiopathic arthritis (JIA),⁴ which presents similarly to AOSD but differs in age of onset. In addition, functional variations in periodic fever syndrome genes have been identified in some AOSD patients (Bonferroni corrected *p* values ranged from 2.34×10^{-3} to 2.40×10^{-4}).⁵

In this study, for the first time, we conducted a genome-wide association study (GWAS) to systematically screen genetic factors influencing susceptibility to AOSD (online supplementary text). Principal component analysis was adopted to evaluate population stratification between cases and controls and to detect outliers for removal (online supplementary figure S1). After quality control and imputation, a total of 3 547 931 variants in 264 AOSD cases and 2420 controls (discovery: 247 cases vs 2163 controls; replication: 17 cases vs 257 controls; online supplementary text and table S1) from China were analysed. The genomic inflation, λ_{GC} , was 1.018, indicating the presence of population stratification to a minimal degree (online supplementary figure S2). A total of 1281 variants, from the *HLA* class I and II regions, exceeded GWS in this analysis (online supplementary figure S3).

The most significant association was with rs9268791 (OR=2.363, $p=4.10 \times 10^{-19}$; table 1 and figure 1A), located in the intergenic region between *HLA-DRA* and *HLA-DRB5* (the *HLA* class II region). For the *HLA* class I region, the top GWS signal was observed for rs3094178 (OR=2.139, $p=1.97 \times 10^{-8}$; table 1 and figure 1B), which is located 19 kb downstream of *HLA-G*. Outside the *HLA* region, a suggestive association ($p < 10^{-5}$) was observed at *VEGFC* (rs514410, OR=2.211, $p=1.81 \times 10^{-6}$; online supplementary table S2, figure S4 and text).

We explored regulatory annotations for the GWS SNPs and variants in strong linkage disequilibrium (LD) with them ($r^2 > 0.8$) using HaploReg⁶ and RegulomeDB,⁷ and 94 SNPs had a RegulomeDB score from 1a–1f to 2a–2c, suggesting that they are located in regulatory regions with maximum evidence (online supplementary table S3). Of them, the top SNPs in each region were rs3115628 ($p=2.92 \times 10^{-8}$; *HLA* class I) and rs9268832 ($p=3.41 \times 10^{-18}$; *HLA* class II). In addition, the expression quantitative trait loci analyses suggested that rs9268832 was significantly associated with the expression of several genes (including *HLA-DRB1*, *HLA-DRB6*, *HLA-DQA2* and *HLA-DQB1*; $p < 5 \times 10^{-8}$) and that rs3115628 was significantly correlated with *ZFP57* and four *HLA* pseudogenes ($p < 5 \times 10^{-8}$) in different types of tissues (online supplementary table S4).⁸ Further studies are required to clarify the mechanisms underlying these associations.

Previous GWAS of JIA on primarily European ancestry samples has also identified three GWS SNPs in the *HLA* region.^{9,10} However, none of them overlapped with the AOSD-associated markers identified in this study. Pair-wise LD analysis showed that these SNPs are in low LD ($r^2 < 0.1$, online supplementary table S5). This finding suggests genetic heterogeneity in these diseases and across different populations.

Table 1 Summary statistics for the genome-wide significant association of single nucleotide polymorphisms with adult-onset Still's disease

SNP	Nearby gene(s)	Chr	Position (hg19)	Risk/non-risk allele	Stage*	RAF in cases	RAF in controls	OR	SE	P values	P_het
rs9268791	<i>HLA-DRA HLA-DRB5</i>	6	32 421 073	T/C	Meta	—	—	2.363	0.096	4.10E-19	0.686
					Discovery	0.468	0.273	2.389	0.100	2.85E-18	
					Replication	0.382	0.224	2.054	0.360	4.54E-02	
rs3094178	<i>HLA-G</i>	6	29 818 000	C/G	Meta	—	—	2.139	0.135	1.97E-08	0.806
					Discovery	0.875	0.769	2.116	0.143	1.48E-07	
					Replication	0.767	0.568	2.366	0.432	4.62E-02	

*The discovery data set comprises 247 cases and 2163 controls from northern and central China. The replication data set comprises 17 cases and 257 controls from southern China.

Chr, chromosome; P_het, *p* value for heterogeneity test across different cohorts using Cochran's Q test; RAF, frequency of risk allele; SNP, single nucleotide polymorphism.

Letters

Figure 1 Regional association plots of the adult-onset Still's disease-associated *HLA* class I and II regions. (A) *HLA* class II region and (B) *HLA* class I region. Purple circles represent the most significantly associated single nucleotide polymorphisms (SNPs) (marker SNPs) in each region in the meta-analysis of discovery and replication. $-\log_{10}p$ values (Y-axis) of the SNPs (within the regions spanning 500 kb on either side of the marker SNP) are presented according to the chromosomal positions of the SNPs (X-axis, hg19). SNPs are coloured according to their linkage disequilibrium (LD) with the marker SNP. The LD values were established based on the 1000 Genomes Asian (ASI) Data (March 2012). Estimated recombination rates with samples from the 1000 Genomes Project March 2012 release are shown as blue lines, and the genomic locations of genes within the regions of interest annotated from the University of California, Santa Cruz (UCSC) genome browser are shown as arrows.

In summary, we identified the first GWS risk loci and indicated an important role for both *HLA* class I and II regions in the genetic architecture of AOSD using a genome-wide scan in a Chinese sample. Further replication studies of other populations are necessary to confirm our findings and to investigate the possible ancestral differences.

Zhiqiang Li^{1,2,3,4}, Hong-Lei Liu⁵, Jianhua Chen^{2,3,6}, Ting Zeng⁷, Lin He^{2,4}, Meihang Li¹, Cainan Luo⁸, Shuang Liu⁹, Ting-Ting Ding¹⁰, Kuerbanjiang Yimaiti⁸, Jialin Teng⁵, Xingwang Li², Yonghe Ding¹, Xiaobing Cheng⁵, Juan Zhou², Junna Ye⁵, Jue Ji², Yu-Tong Su⁵, Hui Shi⁵, Yue Sun⁵, Chengwen Gao¹, Qiongyi Hu⁵, Huihui Chi⁵, Xuan Yuan¹¹, Zhuochao Zhou⁵, Dong Wang², Ke Wang², Chang-Gui Li¹¹, Yuanchao Sun¹, Yujuan Niu¹, Lin-Jie Chen¹⁰, Jian Xu⁹, Lijun Wu⁸, Zhaowei Zhou², Dun Pan², Haitao Niu^{1,12,13}, Yongyong Shi^{1,2,3,4,6,14,15}, Chengde Yang⁵

¹The Affiliated Hospital of Qingdao University & The Biomedical Sciences Institute of Qingdao University (Qingdao Branch of SJTU Bio-X Institutes), Qingdao University, Qingdao, China

²Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders (Ministry of Education), The Collaborative Innovation Center for Brain Science, Shanghai Jiao Tong University, Shanghai, China

³Shanghai Key Laboratory of Sleep Disordered Breathing, Shanghai, China
⁴Institute of Neuropsychiatric Science and Systems Biological Medicine, Shanghai Jiao Tong University, Shanghai, China
⁵Department of Rheumatology and Immunology, Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital, Shanghai, China
⁶Shanghai Key Laboratory of Psychotic Disorders, Shanghai Mental Health Center, Shanghai Jiao Tong University School of Medicine, Shanghai, China
⁷Xinhua Hospital Chongming Branch Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai, China
⁸Department of Rheumatology and Immunology, The People's Hospital of Xinjiang Uygur Autonomous Region, Urumqi, China
⁹Department of Rheumatology and Immunology, First Affiliated Hospital of Kunming Medical University, Kunming, China
¹⁰Department of Rheumatology and Immunology, First Affiliated Hospital of Bengbu Medical College, Bengbu, China
¹¹Shandong Provincial Key Laboratory of Metabolic Disease, The Affiliated Hospital of Qingdao University, Qingdao, China
¹²Department of Urology, The Affiliated Hospital of Qingdao University, Qingdao, China
¹³Key Laboratory, Department of Urology and Andrology, The Affiliated Hospital of Qingdao University, Qingdao, China
¹⁴Department of Psychiatry, The First Teaching Hospital of Xinjiang Medical University, Urumqi, China
¹⁵Changning Mental Health Center, Shanghai, China

Correspondence to Dr Chengde Yang; yangchengde@sina.com Dr Yongyong Shi; shiyongyong@gmail.com Dr Zhiqiang Li; liqzsjtu@163.com Dr Haitao Niu; niuht0532@126.com

Handling editor Josef S Smolen

Acknowledgements We appreciated to all the participants, research staffs, and students who took part in this study.

Contributors ZL, H-LL, YShi and CY designed the study. H-LL, TZ, CL, SL, T-TD, KY, JT, XC, JY, Y-TS, HS, Yue Sun, QH, HC, Zhuochao Zhou, L-JC, JX and LW collected the samples and organized the patient history information. JC, LH, ML, XL, YD, JZ, JJ, CG, XY, DW, KW, C-GL, Yuanchao Sun, YN, Zhaowei Zhou, DP and HN performed or contributed to the experiments. ZL and H-LL conducted data analyses. ZL and H-LL interpreted the results and drafted the manuscript. YShi and CY revised the manuscript. All authors critically reviewed and approved the manuscript.

Funding This study was supported by National Natural Science Foundation of China (81502016 for H-LL, 81671589 for CY, 81701321 and 81871055 for ZL, 31325014 for YShi, 81421061 for LH, 81501154 for JC, 81772713 for HN), the 973 Program (2015CB559100 for YShi), the National Key R&D Program of China (2016YFC0903402 for YShi and ZL), Taishan Scholar Program of Shandong Province (tsqn201812153 for ZL and tsqn20161077 for HN), Natural Science Foundation of Shandong Province (ZR2019YQ14 for ZL), Program of Shanghai Subject Chief Scientist (15XD1502200 for YShi), Shanghai Municipal Health Commission (ZK2015B01 and 20154011 for YShi), National Program for Support of Top-Notch Young Professionals for YShi.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Institutional review board and Institutional Ethical Committee of the AOSD-database partner (Shanghai Jiao Tong University School of Medicine Affiliated Ruijin Hospital, Xinhua Hospital Chongming Branch Affiliated to Shanghai Jiao Tong University School of Medicine, People's hospital of Xinjiang Uygur Autonomous Region, First Affiliated Hospital of Kunming Medical University, First Affiliated Hospital of Bengbu Medical College).

Provenance and peer review Not commissioned; externally peer reviewed.

OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2019-215239>).

ZL and H-LL contributed equally.

To cite Li Z, Liu H-L, Chen J, *et al.* *Ann Rheum Dis* 2020;**79**:161–163.

Received 14 February 2019

Revised 20 August 2019

Accepted 20 August 2019

Published Online First 30 August 2019

Ann Rheum Dis 2020;**79**:161–163. doi:10.1136/annrheumdis-2019-215239

ORCID iDs

Zhiqiang Li <http://orcid.org/0000-0002-0174-9254>

Huihui Chi <http://orcid.org/0000-0001-9587-1180>

REFERENCES

- 1 Giacomelli R, Ruscitti P, Shoenfeld Y. A comprehensive review on adult onset still's disease. *J Autoimmun* 2018;93:24–36.
- 2 Feist E, Mitrovic S, Fautrel B. Mechanisms, biomarkers and targets for adult-onset Still's disease. *Nat Rev Rheumatol* 2018;14:603–18.
- 3 Asano T, Furukawa H, Sato S, *et al.* Effects of HLA-DRB1 alleles on susceptibility and clinical manifestations in Japanese patients with adult onset Still's disease. *Arthritis Res Ther* 2017;19.
- 4 Ombrello MJ, Remmers EF, Tachmazidou I, *et al.* HLA-DRB1*11 and variants of the MHC class II locus are strong risk factors for systemic juvenile idiopathic arthritis. *Proc Natl Acad Sci U S A* 2015;112:15970–5.
- 5 Sighart R, Rech J, Hueber A, *et al.* Evidence for genetic overlap between adult onset Still's disease and hereditary periodic fever syndromes. *Rheumatol Int* 2018;38:111–20.
- 6 Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012;40:D930–D934.
- 7 Boyle AP, Hong EL, Hariharan M, *et al.* Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* 2012;22:1790–7.
- 8 Ardlie KG, DeLuca DS, Segre AV, *et al.* Human genomics. The Genotype-Tissue expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* 2015;348:648–60.
- 9 Hinks A, Cobb J, Marion MC, *et al.* Dense genotyping of immune-related disease regions identifies 14 new susceptibility loci for juvenile idiopathic arthritis. *Nat Genet* 2013;45:664–9.
- 10 Ombrello MJ, Arthur VL, Remmers EF, *et al.* Genetic architecture distinguishes systemic juvenile idiopathic arthritis from other forms of juvenile idiopathic arthritis: clinical and therapeutic implications. *Ann Rheum Dis* 2017;76:906–13.