Dysfunctional missense variant of *OAT10/SLC22A13* decreases gout risk and serum uric acid levels

Organic anion transporter 10 (OAT10), also known as SLC22A13, has hitherto been identified as a urate transporter by *in vitro* analyses.¹ Despite the reported expression of OAT10 on the apical membrane of the renal proximal tubular cells,¹ the physiological impact of OAT10 on urate handling in humans remains to be elucidated. Accumulating evidence suggests that functional variants of already-characterised, physiologically

important urate transporters—URAT1/SLC22A12, GLUT9/SLC2A9, BCRP/ABCG2 and NPT1/SLC17A1—affect serum uric acid (SUA) levels and susceptibility of gout,^{2–6} the most common form of inflammatory arthritis. However, there are no reports on the association between *OAT10* gene and either hyperuricaemia or gout. Here, for the first time, we reveal that a dysfunctional variant of *OAT10* decreases both gout risk and SUA levels, suggesting *OAT10* to be physiologically involved in urate reabsorption in the human kidney, as described below.

To explore exonic variants in *OAT10* potentially associated with gout susceptibility, we sequenced all exons of *OAT10* in 480 gout cases and 480 controls of Japanese male⁶ and conducted an association analysis (see online supplementary tables S1 and S2), followed by a replication study on 924 gout cases and 2113 controls (see online supplementary figure S1). In two identified *OAT10* variants with minor allele frequency (MAF) >0.5%, only rs117371763 (c.1129C>T; p.Arg377Cys [R377C]) was significantly associated with gout susceptibility after Bonferroni correction ($p=0.014$). The significant association between rs117371763 and gout susceptibility was replicated, and our meta-analysis showed a significant protective effect of rs117371763 on gout susceptibility (OR=0.67; 95% CI 0.53 to 0.85; $p_{\text{meta}}=7.8 \times 10^{-4}$) (table 1). In addition, a quantitative trait locus analysis focusing on SUA levels in 3208 individuals (see online supplementary table S3) showed that the minor allele of rs117371763 significantly decreases SUA levels ($\beta=-0.156$ mg/dL, 95% CI -0.295 to -0.018 mg/dL, $p=0.027$). Results were similar even after adjustment for age.

Furthermore, *via* a series of cell-based experiments, we identified the R377C variant as an almost null variant of *OAT10* (figure 1A–C). Immunoblotting and confocal microscopic observations showed the R377C variant to have little effect on *OAT10* protein levels (figure 1A) or its cellular localisation (figure 1B). Cell-based urate transport assay demonstrated that, consistent with a previous report,¹ *OAT10* wild-type can transport urate (figure 1C); however, the urate transport activity of R377C variant-expressing cells was close to that of mock cells, demonstrating that this variant disrupts *OAT10*'s function as a urate transporter. As it is conserved across different species (see online supplementary figure S2), R377 may be important for *OAT10* function.

Considering the following three points, we conclude that *OAT10* is a urate reabsorption transporter on the apical side of the renal proximal tubular cells (figure 1D). First, the R377C variant of *OAT10* was almost null as a urate transporter (figure 1C). Second, this dysfunctional variant decreased SUA levels (see online supplementary table S3), suggesting that functional *OAT10* is physiologically involved in a supply route of urate into the blood. Third, like URAT1/SLC22A12, which plays a pivotal role in urate transport from urine to the blood,² *OAT10* is reportedly expressed in the brush border membranes of the renal epithelium,¹ therefore making it a potential target for urate-lowering therapy like URAT1. Although rs117371763 of

Figure 1 Effects of Arg377Cys (R377C) on the expression, plasma membrane localisation, and function of the organic anion transporter 10 (*OAT10*) urate transporter transiently expressed in 293A cells. (A) (Upper) Immunoblot detection of *OAT10*/SLC22A13 protein in whole cell lysate samples. *OAT10* fused with EGFP was detected by an anti-EGFP antibody. Arrowhead, matured *OAT10* as a glycoprotein; arrow, non-glycosylated form of *OAT10*; α -tubulin, a loading control; (Lower) Relative protein levels of *OAT10* wild-type (WT) and Arg377Cys (R377C) variant. Data are expressed as the mean $\pm$ SD, $n=3$. N.S., not significantly different between groups (two sided t -test). (B) Confocal microscopic observation of cellular localisation. Nuclei were stained with TO-PRO-3 iodide (grey). Bars, 5 μ m. (C) Functional analysis. *OAT10*-expressing 293A cells were incubated with 10 μ M of [¹⁴C]-urate for 60 s, then the amount of urate incorporated into the cells was measured. Data are expressed as the mean $\pm$ SD, $n=7$. ** $p<0.01$ versus the other groups (Tukey-Kramer multiple-comparison test). All experiments were performed 48 hours after plasmid transfection. (D) Proposed physiological model of *OAT10* in human kidney. *OAT10* is expressed on the apical membrane of renal proximal tubules and mediates reabsorption of urate from urine to blood. Other previously characterised urate reabsorption transporters (URAT1/SLC22A12 and GLUT9/SLC2A9) and urate excretion transporters (BCRP/ABCG2 and NPT1/SLC17A1) are also described.

OAT10 is common in Japanese (see online supplementary table S2), this variant is rare in other populations, including European Caucasians (see online supplementary table S4). Such populations, in which most people have functional *OAT10*, may offer a greater potential for *OAT10* as a drug target for the treatment of gout/hyperuricaemia. Our findings will contribute to uncovering

Table 1 Association analysis of *OAT10*/SLC22A13 variant, rs117371763 [Arg377Cys (R377C)], with gout susceptibility

	Gout cases				Controls				p value	OR (95% CI)
	C/C	C/T	T/T	MAF (%)	C/C	C/T	T/T	MAF (%)		
Discovery phase	447	31	2	3.65	427	46	6	6.05	0.014	0.59 (0.38 to 0.90)
Replication phase	859	63	2	3.63	1900	203	5	5.02	0.015	0.71 (0.53 to 0.94)
Meta-analysis									7.8×10^{-4}	0.67 (0.53 to 0.85)

In the meta-analysis, no apparent heterogeneity was observed (p value for Cochran's Q test=0.48, $I^2=0\%$).

MAF, minor allele frequency.

the physiological role of OAT10 as a renal urate reabsorber and its pathophysiological importance in urate-related disorders such as gout/hyperuricaemia.

Toshihide Higashino,^{1,2} Keito Morimoto,³ Hirofumi Nakaoka,⁴ Yu Toyoda,³ Yusuke Kawamura,¹ Seiko Shimizu,¹ Takahiro Nakamura,⁵ Kazuyoshi Hosomichi,⁶ Akiyoshi Nakayama,¹ Keiko Ooyama,⁷ Hiroshi Ooyama,⁷ Toru Shimizu,⁸ Miki Ueno,⁹ Toshimitsu Ito,¹⁰ Takashi Tamura,¹¹ Mariko Naito,¹¹ Hiroshi Nakashima,¹² Makoto Kawaguchi,¹ Mikiya Takao,¹ Yosuke Kawai,¹³ Naoki Osada,² Kimiyoshi Ichida,¹⁴ Ken Yamamoto,¹⁵ Hiroshi Suzuki,³ Nariyoshi Shinomiya,¹ Ituro Inoue,⁴ Tappei Takada,³ Hirotaka Matsuo¹

¹Department of Integrative Physiology and Bio-Nano Medicine, National Defense Medical College, Tokorozawa, Japan

²Graduate School of Information Science and Technology, Hokkaido University, Sapporo, Japan

³Department of Pharmacy, The University of Tokyo Hospital, Tokyo, Japan

⁴Division of Human Genetics, Department of Integrated Genetics, National Institute of Genetics, Mishima, Japan

⁵Laboratory for Mathematics, National Defense Medical College, Tokorozawa, Japan

⁶Department of Bioinformatics and Genomics, Graduate School of Advanced Preventive Medical Sciences, Kanazawa University, Kanazawa, Japan

⁷Ryugoku East Gate Clinic, Tokyo, Japan

⁸Midorigaoka Hospital, Takatsuki, Japan

⁹Division of Nursing, National Defense Medical College, Tokorozawa, Japan

¹⁰Department of Internal Medicine, Self-Defense Forces Central Hospital, Tokyo, Japan

¹¹Department of Preventive Medicine, Nagoya University Graduate School of Medicine, Nagoya, Japan

¹²Department of Preventive Medicine and Public Health, National Defense Medical College, Tokorozawa, Japan

¹³Genome Medical Science Project (Toyama), National Center for Global Health and Medicine, Tokyo, Japan

¹⁴Department of Pathophysiology, Tokyo University of Pharmacy and Life Sciences, Tokyo, Japan

¹⁵Department of Medical Biochemistry, Kurume University School of Medicine, Kurume, Japan

Correspondence to Dr Hirotaka Matsuo, Department of Integrative Physiology and Bio-Nano Medicine, National Defense Medical College, Tokorozawa 359-8513, Japan; hmatsuo@ndmc.ac.jp and Dr Tappei Takada, Department of Pharmacy, The University of Tokyo Hospital, Tokyo, Japan; tappei-ty@umin.ac.jp

Handling editor Josef S Smolen

Acknowledgements We would like to thank all the participants for their generous involvement in this study. We also thank members of J-MICC Study Shizuoka Field and Daiko Field for supporting the study. We are indebted to K Gotanda, M Miyazawa, Y Aoyagi, Y Aoki and K Yokoi (National Defense Medical College) for genetic analysis. We are indebted to M Senda (Ryugoku East Gate Clinic), H Fujiwara (Midorigaoka Hospital), K Wakai and N Hamajima (Nagoya University) for sample collection.

Contributors TH, HNakaoka, YT, TTakada and HM conceived and designed this study. TN, KH, AN, MU, TI, KI, KY, HS, NS and II assisted with research design. SS, KO, HO, TS, NS and HM collected and analysed clinical data of cases. YKawamura, SS, MU, TI, TTamura, MN, HNakashima, MK, MT and HM collected and analysed clinical data of controls. TH, HNakaoka, SS, NS, II and HM performed genetic analysis. HNakaoka, YKawamura, HNakashima, TN and II performed statistical analyses. KM, YT, HS and TTakada performed functional analysis. TTakada and HM organised this collaborative study as corresponding authors. KM, TN, KH, AN, YKawai, NO, KI and KY provided intellectual input and assisted with the preparation of the manuscripts. TH, YT, YKawamura, TTakada and HM wrote the manuscript. TH, KM, HNakaoka, YT, YKawamura and SS contributed equally to this work. All authors have read and approved the final version of the manuscript.

Funding This study was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan (Nos 17H04128, 19K22786, 25293145, 15K15227, 17015018, 221S0001, 221S0002, 15H05610, 16H06277, 16H06279, 16H01808, 18KK0247 and 22136015), the Ministry of Defense, the Uehara Memorial Foundation, Mochida Memorial Foundation for Medical and Pharmaceutical Research, the Takeda Medical Foundation, MSD Life Science Foundation, Public Interest Incorporated Foundation, the Kawano Masanori Memorial Foundation for Promotion of Pediatrics and the Gout Research Foundation of Japan.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was approved by the institutions' Ethical Committees (National Defense Medical College, National Institute of Genetics, and Nagoya University). All procedures were performed in accordance with the Declaration of Helsinki, with written informed consent obtained from each participant.

Provenance and peer review Not commissioned; externally peer reviewed.

OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2019-216044>).

TH, KM, HNakaoka, YT, YKawamura and SS contributed equally.

To cite Higashino T, Morimoto K, Nakaoka H, *et al.* *Ann Rheum Dis* 2020;**79**:164–166.

Received 18 July 2019

Revised 13 September 2019

Accepted 30 September 2019

Published Online First 28 November 2019

Ann Rheum Dis 2020;**79**:164–166. doi:10.1136/annrheumdis-2019-216044

REFERENCES

- 1 Bahn A, Hagos Y, Reuter S, *et al.* Identification of a new urate and high affinity nicotinate transporter, hOAT10 (SLC22A13). *J Biol Chem* 2008;283:16332–41.
- 2 Enomoto A, Kimura H, Chairoungdua A, *et al.* Molecular identification of a renal urate anion exchanger that regulates blood urate levels. *Nature* 2002;417:447–52.
- 3 Matsuo H, Takada T, Ichida K, *et al.* Common defects of ABCG2, a high-capacity urate exporter, cause gout: a function-based genetic analysis in a Japanese population. *Sci Transl Med* 2009;1:5ra11.
- 4 Woodward OM, Köttgen A, Coresh J, *et al.* Identification of a urate transporter, ABCG2, with a common functional polymorphism causing gout. *Proc Natl Acad Sci U S A* 2009;106:10338–42.
- 5 Nakayama A, Nakaoka H, Yamamoto K, *et al.* GWAS of clinically defined gout and subtypes identifies multiple susceptibility loci that include urate transporter genes. *Ann Rheum Dis* 2017;76:869–77.
- 6 Higashino T, Takada T, Nakaoka H, *et al.* Multiple common and rare variants of ABCG2 cause gout. *RMD Open* 2017;3:e000464.