


## Online supplementary figure S2



**Figure S2** Flowchart showing the selection process

Abbreviations: YvE-H; convenor, BB; fellow, CIA; Chronic Inflammatory Arthritis, ACR/ARHP: The American College of Rheumatology and The Association of Rheumatology Health Professionals, EULAR; The European League Against Rheumatism.

\*Abstracts from EULAR 2016 and ACR/ARHP 2016 and 2017 were covered in the main search.

## References to the included records (n=51)

- Amity CL, Schlenk EA, Gold KN, et al. Agreement of Physicians and Nurses Performing Tender and Swollen Joint Counts in Rheumatoid Arthritis. *J Clin Rheumatol* 2016;22(1):30-4.
- Arends RY, Bode C, Taal E, et al. A mixed-methods process evaluation of a goal management intervention for patients with polyarthritis. *Psychol Health* 2017;32(1):38-60.
- Bala SV, Samuelson K, Hagell P, et al. The experience of care at nurse-led rheumatology clinics. *Musculoskeletal Care*. 2012;10(4):202-11.
- Beattie KA, Macintyre NJ, Cividino A. Screening for signs and symptoms of rheumatoid arthritis by family physicians and nurse practitioners using the Gait, Arms, Legs, and Spine musculoskeletal examination. *Arthritis Care Res (Hoboken)* 2012;64(12):1923-27.
- Bellerby T, Stevens D, Arnold T, et al. Audit of nurse-led DGH rheumatology advice line [abstract]. *Ann Rheum Dis* 2016;75:1302-3.
- Cheung PP, Dougados M, Andre V, et al. The learning curve of nurses for the assessment of swollen and tender joints in rheumatoid arthritis. *Joint Bone Spine* 2014;81(2):154-59.
- Cheung PP, Ruysen-Witrand A, Gossec L, et al. Reliability of patient self-evaluation of swollen and tender joints in rheumatoid arthritis: A comparison study with ultrasonography, physician, and nurse assessments. *Arthritis Care Res (Hoboken)* 2010;62(8):1112-19.
- Cheung YS, Lee PF, Tang SK. Starting care at the peep of rheumatoid arthritis-a patient education and empowerment program [abstract]. *Ann Rheum Dis* 2016;75:1293-94.
- Cottrell JED, Jonas M, Bergsten U, et al The Nurse's Role in Addressing Unmet Treatment and Management Needs of Patients with Rheumatoid Arthritis: Delphi-Based Recommendations. *Int J Nurs Knowl* 2013;24(2):66-76.
- De Thurah A, Esbensen BA, Roelsgaard IK, et al. Efficacy of embedded nurse-led versus conventional physician-led follow-up in rheumatoid arthritis: A systematic review and meta-Analysis. *RMD open* 2017;3(2):e000481.
- De Thurah A, Stengaard-Pedersen K, Axelsen M, et al. Tele-Health Followup Strategy for Tight Control of Disease Activity in Rheumatoid Arthritis: Results of a Randomized Controlled Trial. *Arthritis Care Res (Hoboken)* 2018;70(3):353-60.
- Dougados M, Soubrier M, Perrodeau E, et al. Impact of a nurse-led programme on comorbidity management and impact of a patient self-assessment of disease activity on the management of

- rheumatoid arthritis: results of a prospective, multicentre, randomised, controlled trial (COMEDRA). *Ann Rheum Dis* 2015;74(9):1725-33.
- Fitzgerald P, Hale C. Moving the injectable methotrexate service closer to home: a practice development initiative. *Musculoskeletal Care* 2014;12(3):182-89.
- Frolund JC, Primdahl J. Patients' Experiences of Nurse-Led Screening for Cardiovascular Risk in Rheumatoid Arthritis. *Musculoskeletal Care* 2015;13(4):236-47
- Fusama M, Higashi K, Maeda K, et al. Survey on the understanding and practice of T2T for nurses engaged in medical treatment of rheumatoid arthritis [abstract]. *Arthritis Rheumatol* 2016;68(Suppl 10):1497.
- Gerega S, Millson B, Bessette L, et al. Canadian study of outcomes in adalimumab (Humira) patients with support for adherence-results from the companion study [abstract]. *Arthritis Rheumatol* 2016;68:806-07.
- Gossec L, Soubrier M, Foissac F, et al. One third of patients with established rheumatoid arthritis (RA) are correctly vaccinated against influenza and pneumococcus and this is increasing: 3 year longitudinal assessment of 776 patients [abstract]. *Arthritis Rheumatol* 2016;68(Suppl 10):1874-76.
- Goulenok T SJ, Francois C, Van der Haegen M-C, et al. Nurse-led vaccination program dramatically improves pneumococcal vaccination coverage among patients with chronic inflammatory rheumatic diseases: a prospective pilot study [abstract]. *Ann Rheum Dis* 2017;76(Suppl 2):78-2.
- Grønning K, Midttun L, Steinsbekk A. Patients' confidence in coping with arthritis after nurse-led education; a qualitative study. *BMC Nurs* 2016;15:28.
- Grønning K, Rannestad T, Skomsvoll JF, et al. Long-term effects of a nurse-led group and individual patient education programme for patients with chronic inflammatory polyarthritis - a randomised controlled trial. *J Clin Nurs* 2014;23(7-8):1005-17.
- Grønning K, Skomsvoll JF, Rannestad T, et al. The effect of an educational programme consisting of group and individual arthritis education for patients with polyarthritis-A randomised controlled trial. *Patient Educ Couns* 2012;88(1):113-20.
- Helland Y, Garratt A, Kjekken I, et al. Current practice and barriers to the management of sexual issues in rheumatology: Results of a survey of health professionals. *Scand J Rheumatol* 2013;42(1):20-26.

Hoving JL, Zoer I, Van Der Meer M, et al. E-health to improve work functioning in employees with rheumatoid arthritis in rheumatology practice: A feasibility study. *Scand J Rheumatol* 2014;43(6):481-87.

Hulst LT, Creemers MC, Fransen J, et al. How to improve DAS28 use in daily clinical practice? - a pilot study of a nurse-led intervention. *Rheumatology (Oxford)* 2010;49(4):741-48.

Hurkmans EJ, De Gucht V, Maes S, et al. Promoting physical activity in patients with rheumatoid arthritis: Rheumatologists' and health professionals' practice and educational needs. *Clin Rheumatol* 2011;30(12):1603-09.

Knittle K, De Gucht V, Hurkmans E, et al. Targeting motivation and self-regulation to increase physical activity among patients with rheumatoid arthritis: a randomised controlled trial. *Clin Rheumatol* 2015;34(2):231-38.

Kwok SKG, Tang LMM, Chung HYJ, et al. Rapid access triage clinic: A nursing service that shortens rheumatology clinic waiting time and facilitates early treatment [abstract]. *Ann Rheum Dis* 2016;75:1291.

Larsson I, Bergman S, Fridlund B, et al. Patients' experiences of a nurse-led rheumatology clinic in Sweden: a qualitative study. *Nurs Health Sci* 2012;14(4):501-07.

Larsson I, Fridlund B, Arvidsson B, et al. Randomized controlled trial of a nurse-led rheumatology clinic for monitoring biological therapy. *J Adv Nurs*. 2014;70(1):164-75.

Larsson I, Fridlund B, Arvidsson B, et al. A nurse-led rheumatology clinic versus rheumatologist-led clinic in monitoring of patients with chronic inflammatory arthritis undergoing biological therapy: a cost comparison study in a randomised controlled trial. *BMC Musculoskelet Disord* 2015;16:354.

Libman B, Farley S, Edwards M, et al. Nurse telephone education for promoting a treat-to-target approach in recently diagnosed rheumatoid arthritis patients-a preliminary review. [abstract]. *Arthritis Rheumatol* 2017;69(Suppl 10). Later published as: Farley S, Libman B, Edwards M, et al. Nurse telephone education for promoting a treat-to-target approach in recently diagnosed rheumatoid arthritis patients: a pilot project. *Musculoskeletal Care* 2019;17:156-60

Lineker SC, Bell MJ, Badley EM. Evaluation of an inter-professional educational intervention to improve the use of arthritis best practices in primary care. *J Rheumatol* 2011;38(5):931-37.

Lineker SC, Husted JA, Brown KS. Influence of discipline of provider and model of care on an arthritis educational intervention in primary care. *Arthritis Care Res (Hoboken)* 2012;64(3):424-33.

Mawson M, Hardy S. Using a psychological approach in rheumatoid arthritis. *Practice Nursing* 2017;28(10):431-6.

Sousa FIM, Santos EJM, Cunha M, et al. Effectiveness of nursing consultations in people with rheumatoid arthritis: systematic review. *Revista de Enfermagem Referência* 2017;4(13):147-56.

Mourgues C, Blanquet M, Gerbaud L, et al. Economic analysis of a nurse-led programme for comorbidities management of rheumatoid arthritis patients. *Joint Bone Spine* 2018;85(5):573-76.

Munoz-Fernández S, Aguilar MD, Almodóvar R, et al. SCORE study: quality indicators for rheumatology nursing clinics. *Rheumatol Int* 2017;37(3):353-61.

Munoz-Fernández S, Aguilar MD, Rodriguez A, et al. Evaluation of the impact of nursing clinics in the rheumatology services. *Rheumatol Int* 2016;36(9):1309-17.

Ndosi M, Johnson D, Young T, et al. Effects of needs-based patient education on self-efficacy and health outcomes in people with rheumatoid arthritis: a multicentre, single blind, randomised controlled trial. *Ann Rheum Dis* 2016;75(6):1126-32.

Ndosi M, Lewis M, Hale C, et al. The outcome and cost-effectiveness of nurse-led care in people with rheumatoid arthritis: a multicentre randomised controlled trial. *Ann Rheum Dis* 2014;73(11):1975-82.

Nisar MK, Bergum J. Rheumatology nurse specialists and dmard prescription - where are we now? [abstract]. *Ann Rheum Dis* 2017;76(Suppl 2):1484-3.

Paek SI, Kim SH, Kim H, et al. Effects of an educational program using treat to target strategy in Korean patients with rheumatoid arthritis [abstract]. *Arthritis Rheumatol* 2016;68:1489

Primdahl J, Sorensen J, Horn HC, et al. Shared care or nursing consultations as an alternative to rheumatologist follow-up for rheumatoid arthritis outpatients with low disease activity-patient outcomes from a 2-year, randomised controlled trial. *Ann Rheum Dis*. 2014;73(2):357-64.

Primdahl J, Wagner L, Holst R, et al. The impact on self-efficacy of different types of follow-up care and disease status in patients with rheumatoid arthritis-A randomized trial. *Patient Educ Couns* 2012;88(1):121-8.

Primdahl J, Wagner L, Hørslev-Petersen K. Being an outpatient with rheumatoid arthritis - a focus group study on patients' self-efficacy and experiences from participation in a short course and one of three different outpatient settings. *Scand J Caring Sci* 2011;25(2):394-403.

- Robinson S, Hassell A, Ryan S, et al. A national survey of nurse training: Confidence and competence in educating patients commencing methotrexate therapy. *Musculoskeletal Care* 2017;15(3):281-92.
- Schuch F, Kern P, Kreher G, et al. [Curriculum for "Rheumatology health professionals DGRh-BDRh" as a way of qualification for clinical nursing specialists: Concept and results of an evaluation]. *Z Rheumatol* 2011;70(8):670-7.
- Solomon DH, Bitton A, Fraenkel L, et al. Roles of nurse practitioners and physician assistants in rheumatology practices in the US. *Arthritis Care Res (Hoboken)* 2014;66(7):1108-13.
- Sørensen J, Primdahl J, Horn H, et al. Shared care or nurse consultations as an alternative to rheumatologist follow-up for rheumatoid arthritis (RA) outpatients with stable low disease-activity RA: cost-effectiveness based on a 2-year randomized trial. *Scand J Rheumatol* 2015;44(1):13-21.
- Vinall-Collier K, Madill A, Firth J. A multi-centre study of interactional style in nurse specialist- and physician-led Rheumatology clinics in the UK. *Int J Nurs Stud* 2016;59:41-50.
- Watts RA, Mooney J, Barton G, et al. The outcome and cost-effectiveness of nurse-led care in the community for people with rheumatoid arthritis: a non-randomised pragmatic study. *BMJ open* 2015;5(8):e007696.